

Variable	Univariate HR (95% CI)	p-Value	Variable	Adjusted HR (95% CI)	p-Value
Age (continuous)	0.97 (0.94-1.01)	0.111	Age (continuous)	0.98 (0.94-1.01)	0.189
Visceral disease No Yes	Reference 1.07 (0.51-2.25)	0.866	Visceral disease No Yes	Reference 1.13 (0.53-2.41)	0.761
HR Negative Positive	Reference 1.64 (0.82-3.26)	0.161	HR Negative Positive	Reference 1.51 (0.76-3.03)	0.242
TILs High (>=5%) Low (<5%)	Reference 2.09 (1.05- 4.15)	0.036	TILs High (>=5%) Low (<5%)	Reference 1.88 (0.94-3.77)	0.075

Table 5 Cox Prognostic Models for TILs and PFS for the cohort of patients whose TILs were evaluated at the metastatic site. (= 57 patients)

At univariate analysis, women with low TILs had a worse prognosis (PFS) (HR=2.09, 95%CI: 1.05-4.15, p=0.036, and when adjusting for age, visceral disease and HR status, an association of TILs high vs low demonstrated a similar trend at limit of statistical significance (HR=1.88, 95%CI: 0.94-3.77)

Abbreviations: CI: confidence interval; HR: hormone receptor; PFS: progression-free survival; TILs: tumor-infiltrating lymphocytes.