

Variable	Univariate HR (95% CI)	p-Value	Variable	Adjusted HR (95% CI)	p-Value
Age (continuous)	0.98 (0.94-1.03)	0.386	Age (continuous)	0.99 (0.95-1.04)	0.679
Visceral disease No Yes	Reference 1.34 (0.47-3.78)	0.583	Visceral disease No Yes	Reference 1.74 (0.60-5.04)	0.310
HR Negative Positive	Reference 2.44 (0.91-6.51)	0.076	HR Negative Positive	Reference 2.46 (0.90-6.71)	0.078
TILs High (>=5%) Low (<5%)	Reference 3.18 (1.19- 8.50)	0.021	TILs High (>=5%) Low (<5%)	Reference 3.17 (1.17-8.65)	0.024

Table 6 Cox Prognostic Models for TILs and OS for the cohort of patients whose TILs were evaluated at the metastatic site. (= 57 patients)

At univariate analysis, women with low TILs had a worse prognosis (OS) (HR=3.18, 95%CI: 1.19-8.50, p=0.021, and when adjusting for age, visceral disease and hormonal receptor status significant association of TILs with OS was confirmed (HR=3.17,95%CI: 1.17-8.65, p=0.024)

Abbreviations: CI: confidence interval; HR: hormone receptor; OS: overall survival; TILs: tumour-infiltrating lymphocytes