

Table 8. Logistic regression analysis for positive cultured ascites

Variable	Univariate		Multivariate	
	OR (95% CI)	<i>P</i>	OR (95% CI)	<i>P</i>
Intercept			0.002 (0.001-0.378)	0.02
Age	1.090 (1.020-1.160)	0.005	1.050 (0.980-1.120)	0.18
BMI	0.901 (0.788-1.030)	0.13		
Male gender	2.220 (0.693-7.120)	0.18		
ASA	3.450 (1.500-7.950)	0.003	1.410 (0.514-3.890)	0.50
Previous abdominal/pelvic surgery	0.976 (0.380-2.510)	0.960		
Preoperative use of proton pump inhibitor	2.040 (0.735-5.640)	0.17		
Frailty	6.250 (2.570-15.20)	<0.001	3.910 (1.350-11.30)	0.01
Operative time	0.996 (0.990-1.000)	0.18		
Intestinal tract reconstruction time	1.000 (0.972-1.030)	0.91		
Diversion type (neobladder)	0.428 (0.116-1.570)	0.20		
EBL	1.000 (0.998-1.000)	0.72		
Intraoperative crystalloid administration	1.000 (0.999-1.000)	0.56		
Blood transfusion	1.300 (0.274-8.300)	0.76		
ERAS	0.762 (0.301-1.930)	0.56		
Postoperative Alb	0.457 (0.125-1.670)	0.23		
Postoperative CRP	0.987 (0.864-1.130)	0.84		

OR, odds ratio; CI, confidence interval; BMI, body mass index; ASA, American Society of Anesthesiologists; EBL, estimated blood loss; ERAS, enhanced recovery after surgery; Alb, albumin; CRP, c-reactive protein