

Establishment of a depression model using dexamethasone-treated three-dimensional cultured rat cortical cells

Mi Kyoung Seo[†], Sehoon Jeong[†], Woo Seok Cheon, Dong Yun Lee, Sumin Lee,
Gyu-Hui Lee, Deok-Gyeong Kang, Dae-Hyun Seog, Seong-Ho Kim,
Jung Goo Lee^{*}, and Sung Woo Park^{*}

Journal name: Cellular and Molecular Neurobiology

*** Corresponding authors**

Sung Woo Park, Ph.D. and Jung Goo Lee, M.D., Ph.D.


Department of Convergence Biomedical Science, College of Medicine, Inje University,
75, Bokji-ro, Busanjin-gu, Busan 47392, Republic of Korea. (S.-W. Park)

Department of Psychiatry, College of Medicine, Haeundae Paik Hospital, Inje University,
Busan 48108, Republic of Korea. (J.-G. Lee)


E-mail address: swpark@inje.ac.kr (S.-W. Park) / iybihwc@naver.com (J.-G. Lee)

Fig. 5


A. BDNF



B. Sirtuin 1




C. ERK1/2




The original images from which we constructed fig. 5 are shown.

Fig. 6 A


a) mTORC1



b) 4E-BP1




c) p70S6K




The original images from which we constructed fig. 6A are shown.

Fig. 6 B

a) PSD-95



b) GluA1



The original images from which we constructed fig. 6B are shown.