

Additional file 3. Hospitalization causes of the patient group of GP1 and the patient group of other GPs in 2018/19: data listed as counts (n) with percentages (%).

	Missing	Overall	GP1	Other GPs	P-Value
n		74	19	55	
Breathing difficulties/Dyspnoea, n (%)	0	7 (9.5)		7 (12.7)	0.180
Gastrointestinal problems, n (%)	0	7 (9.5)	1 (5.3)	6 (10.9)	0.669
Cardiovascular disorder, n (%)	0	3 (4.1)		3 (5.5)	0.565
Fall, n (%)	0	19 (25.7)	4 (21.1)	15 (27.3)	0.764
General health status deterioration (unclear genesis), n (%)	0	12 (16.2)	4 (21.1)	8 (14.5)	0.491
Urologic problems, n (%)	0	1 (1.4)	1 (5.3)		0.257
Nephrological problems, n (%)	0	2 (2.7)		2 (3.6)	1.000
Thoracic pain, n (%)	0	5 (6.8)	2 (10.5)	3 (5.5)	0.598
Abdominal pain, n (%)	0	1 (1.4)		1 (1.8)	1.000
Suspected apoplexy/TIA, n (%)	0	5 (6.8)	2 (10.5)	3 (5.5)	0.598
Suspected pneumonia, n (%)	0	3 (4.1)		3 (5.5)	0.565
Psychopathological anomaly, n (%)	0	3 (4.1)		3 (5.5)	0.565
Suspected thrombosis, n (%)	0	2 (2.7)		2 (3.6)	1.000
Oedema, n (%)	0	2 (2.7)		2 (3.6)	1.000
Inflammation signs of the lower extremities, n (%)	0	3 (4.1)	1 (5.3)	2 (3.6)	1.000
Exsiccosis n (%)	0	1 (1.4)	1 (5.3)		0.257
Dermatological anomaly, n (%)	0	1 (1.4)	1 (5.3)		0.257
Medication adjustment, n (%)	0	1 (1.4)	1 (5.3)		0.257
Suspected lung embolism, n (%)	0	1 (1.4)	1 (5.3)		0.257

Abbreviation. TIA: transient ischaemic attack.