



# Full wwPDB X-ray Structure Validation Report i

Jun 25, 2024 – 11:00 AM EDT

PDB ID : 9CD8  
Title : Crystal Structure of Acetyl-CoA synthetase from *Cryptococcus neoformans* H99 in complex with inhibitor HGN-1310 (dd3-027)  
Deposited on : 2024-06-24  
Resolution : 2.40 Å (reported)

This wwPDB validation report is for manuscript review

This is a Full wwPDB X-ray Structure Validation Report.

This report is produced by the wwPDB biocuration pipeline after annotation of the structure.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

The types of validation reports are described at  
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
 Validation Pipeline (wwPDB-VP) : 2.37.1

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	A1AV1	B	706	-	-	-	X

## 2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 15201 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acetyl-coenzyme A synthetase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	657	5157	3287	879	965	26	0	3	0

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-13	MET	-	initiating methionine	UNP J9VFT1
A	-12	HIS	-	expression tag	UNP J9VFT1
A	-11	HIS	-	expression tag	UNP J9VFT1
A	-10	HIS	-	expression tag	UNP J9VFT1
A	-9	HIS	-	expression tag	UNP J9VFT1
A	-8	HIS	-	expression tag	UNP J9VFT1
A	-7	HIS	-	expression tag	UNP J9VFT1
A	-6	HIS	-	expression tag	UNP J9VFT1
A	-5	HIS	-	expression tag	UNP J9VFT1
A	-4	GLU	-	expression tag	UNP J9VFT1
A	-3	ASN	-	expression tag	UNP J9VFT1
A	-2	LEU	-	expression tag	UNP J9VFT1
A	-1	TYR	-	expression tag	UNP J9VFT1
A	0	PHE	-	expression tag	UNP J9VFT1
A	1	GLN	-	expression tag	UNP J9VFT1


- Molecule 2 is a protein called Acetyl-coenzyme A synthetase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	536	4253	2712	724	794	23	0	3	0
2	C	656	5084	3247	867	944	26	0	2	0

There are 30 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	-13	MET	-	initiating methionine	UNP J9VFT1
B	-12	HIS	-	expression tag	UNP J9VFT1
B	-11	HIS	-	expression tag	UNP J9VFT1
B	-10	HIS	-	expression tag	UNP J9VFT1
B	-9	HIS	-	expression tag	UNP J9VFT1
B	-8	HIS	-	expression tag	UNP J9VFT1
B	-7	HIS	-	expression tag	UNP J9VFT1
B	-6	HIS	-	expression tag	UNP J9VFT1
B	-5	HIS	-	expression tag	UNP J9VFT1
B	-4	GLU	-	expression tag	UNP J9VFT1
B	-3	ASN	-	expression tag	UNP J9VFT1
B	-2	LEU	-	expression tag	UNP J9VFT1
B	-1	TYR	-	expression tag	UNP J9VFT1
B	0	PHE	-	expression tag	UNP J9VFT1
B	1	GLN	-	expression tag	UNP J9VFT1
C	-13	MET	-	initiating methionine	UNP J9VFT1
C	-12	HIS	-	expression tag	UNP J9VFT1
C	-11	HIS	-	expression tag	UNP J9VFT1
C	-10	HIS	-	expression tag	UNP J9VFT1
C	-9	HIS	-	expression tag	UNP J9VFT1
C	-8	HIS	-	expression tag	UNP J9VFT1
C	-7	HIS	-	expression tag	UNP J9VFT1
C	-6	HIS	-	expression tag	UNP J9VFT1
C	-5	HIS	-	expression tag	UNP J9VFT1
C	-4	GLU	-	expression tag	UNP J9VFT1
C	-3	ASN	-	expression tag	UNP J9VFT1
C	-2	LEU	-	expression tag	UNP J9VFT1
C	-1	TYR	-	expression tag	UNP J9VFT1
C	0	PHE	-	expression tag	UNP J9VFT1
C	1	GLN	-	expression tag	UNP J9VFT1

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	C	1	Total C O 6 3 3	0	0
3	C	1	Total C O 6 3 3	0	0
3	C	1	Total C O 6 3 3	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Cl 1 1	0	0

- Molecule 6 is (5P)-3-azido-N-[(5-cyclopropyl-1,2-oxazol-3-yl)methyl]-5-(2,3-dihydro-1-benzo-furan-5-yl)-N-methylbenzamide (three-letter code: A1AV1) (formula: C<sub>23</sub>H<sub>21</sub>N<sub>5</sub>O<sub>3</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
6	A	1	Total 28	C 23	N 2	O 3	0	0
6	B	1	Total 28	C 23	N 2	O 3	0	0
6	C	1	Total 28	C 23	N 2	O 3	0	0

- Molecule 7 is water.


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	204	Total 204	O 204		0	0
7	B	149	Total 149	O 149		0	0
7	C	195	Total 195	O 195		0	0

### 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Acetyl-coenzyme A synthetase

Chain A:




- Molecule 2: Acetyl-coenzyme A synthetase


Chain B:



- Molecule 2: Acetyl-coenzyme A synthetase

Chain C:





For Manuscript Review

## 4 Data and refinement statistics (i)

Property	Value			Source
Space group	P 41 21 2			Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	176.22Å 90.00°	176.22Å 90.00°	159.46Å 90.00°	Depositor
Resolution (Å)	124.61	–	2.40	Depositor
	124.61	–	2.40	EDS
% Data completeness (in resolution range)	100.0 (124.61-2.40) 100.0 (124.61-2.40)			Depositor EDS
$R_{merge}$	0.18			Depositor
$R_{sym}$	(Not available)			Depositor
$\langle I/\sigma(I) \rangle^1$	1.81 (at 2.40Å)			Xtriage
Refinement program	PHENIX 1.21rc1_5177			Depositor
$R$ , $R_{free}$	0.178 0.176	,	0.219 0.211	Depositor DCC
$R_{free}$ test set	4864 reflections (4.96%)			wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	46.1			Xtriage
Anisotropy	0.180			Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32, 40.5			EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$			Xtriage
Estimated twinning fraction	No twinning to report.			Xtriage
$F_o, F_c$ correlation	0.96			EDS
Total number of atoms	15201			wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	54.0			wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.94% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [\(i\)](#)

### 5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ALY, SO4, GOL, CL, A1AV1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.26	0/5284	0.52	0/7190
2	B	0.25	0/4379	0.51	0/5961
2	C	0.26	0/5221	0.51	0/7114
All	All	0.26	0/14884	0.52	0/20265

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5157	0	5006	19	0
2	B	4253	0	4093	17	0
2	C	5084	0	4886	13	0
3	A	18	0	24	1	0
3	B	18	0	24	0	0
3	C	18	0	24	0	0
4	A	10	0	0	0	0
4	B	10	0	0	1	0
5	A	1	0	0	0	0
6	A	28	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	28	0	0	0	0
6	C	28	0	0	0	0
7	A	204	0	0	1	0
7	B	149	0	0	0	0
7	C	195	0	0	0	0
All	All	15201	0	14057	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:390:ARG:NH1	1:A:418:GLU:OE2	2.27	0.68
1:A:642:MET:HE1	1:A:663:ILE:HD12	1.81	0.63
2:B:544:ASP:O	2:B:546:VAL:HG23	1.99	0.61
1:A:645:VAL:HG23	7:A:840:HOH:O	2.07	0.55
1:A:462:ALA:O	1:A:463:THR:OG1	2.19	0.54
2:B:214:ASP:OD1	2:B:215:GLU:N	2.42	0.53
1:A:565:LEU:HD12	1:A:617:VAL:HG21	1.92	0.52
2:C:570:ALA:HB2	2:C:593:LYS:HG3	1.92	0.51
1:A:663:ILE:HD13	1:A:666:PRO:HA	1.93	0.51
2:B:199:ARG:NH1	2:B:235:GLN:OE1	2.44	0.49
2:B:463:THR:OG1	2:B:464:PHE:N	2.38	0.49
2:B:517:LYS:N	2:B:518:PRO:CD	2.76	0.49
2:B:387:ARG:NE	4:B:704:SO4:O4	2.37	0.47
1:A:435[A]:VAL:O	1:A:435[A]:VAL:HG23	2.14	0.47
1:A:517:LYS:N	1:A:518:PRO:CD	2.79	0.46
1:A:642:MET:CE	1:A:663:ILE:HD12	2.45	0.45
1:A:170:TRP:HE1	3:A:702:GOL:H2	1.81	0.45
2:B:295:LYS:N	2:B:295:LYS:HD3	2.32	0.45
2:C:591:THR:HG22	2:C:630:VAL:HG23	1.99	0.45
1:A:14:HIS:HB2	1:A:15:PRO:HD2	1.99	0.45
2:B:295:LYS:O	2:B:295:LYS:HG2	2.17	0.44
2:B:314:LEU:HD22	2:B:345:PRO:HA	1.99	0.44
2:C:475:ASP:HB2	2:C:482:LEU:HD21	2.00	0.44
1:A:631:SER:OG	1:A:671:GLU:OE1	2.35	0.44
1:A:603:GLU:HG3	1:A:629:LEU:HD12	2.00	0.44
2:C:569:VAL:HG13	2:C:590:VAL:HG13	1.99	0.44
1:A:213:THR:HG23	1:A:225:THR:OG1	2.18	0.43
2:C:435[A]:VAL:HG23	2:C:435[A]:VAL:O	2.17	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:517:LYS:N	2:C:518:PRO:CD	2.81	0.43
1:A:606:LEU:HD23	1:A:629:LEU:CD1	2.48	0.43
2:B:425:ASP:O	2:B:429:LYS:HA	2.18	0.43
2:C:463:THR:OG1	2:C:464:PHE:N	2.51	0.43
2:B:405:LEU:HD13	2:B:408:LEU:HD21	2.01	0.43
2:B:380:TYR:CZ	2:B:411:VAL:HB	2.54	0.43
1:A:214:ASP:OD1	1:A:215:GLU:N	2.52	0.42
2:B:213:THR:HG23	2:B:225:THR:OG1	2.18	0.42
2:B:58:THR:HG22	2:B:66:TRP:CD2	2.54	0.42
2:C:58:THR:HG22	2:C:66:TRP:CD2	2.54	0.42
2:C:560:GLU:HG3	2:C:574:VAL:HG13	2.01	0.42
2:C:214:ASP:OD1	2:C:215:GLU:N	2.50	0.42
2:B:271:ALA:HB3	2:C:93:ASP:HB3	2.01	0.41
2:B:472:ASP:OD2	2:B:494:ARG:NH1	2.53	0.41
2:C:442:GLU:OE1	2:C:442:GLU:N	2.51	0.41
1:A:475:ASP:HB2	1:A:482:LEU:HD21	2.01	0.41
2:B:49:GLU:HA	2:B:52:VAL:HG12	2.03	0.41
1:A:561:SER:O	1:A:564:ILE:HG22	2.21	0.41
1:A:606:LEU:HD23	1:A:629:LEU:HD13	2.03	0.40
2:C:668:ILE:O	2:C:672:VAL:HG23	2.21	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles

#### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	653/694 (94%)	630 (96%)	22 (3%)	1 (0%)	47 62
2	B	535/694 (77%)	516 (96%)	18 (3%)	1 (0%)	47 62
2	C	648/694 (93%)	626 (97%)	21 (3%)	1 (0%)	47 62
All	All	1836/2082 (88%)	1772 (96%)	61 (3%)	3 (0%)	47 62

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	463	THR
2	C	463	THR
2	B	463	THR

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	539/575 (94%)	534 (99%)	5 (1%)	78 90
2	B	445/576 (77%)	438 (98%)	7 (2%)	62 79
2	C	523/576 (91%)	511 (98%)	12 (2%)	50 70
All	All	1507/1727 (87%)	1483 (98%)	24 (2%)	65 79

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	110	ASP
1	A	116	ASN
1	A	262	TRP
1	A	290	SER
1	A	472	ASP
2	B	110	ASP
2	B	116	ASN
2	B	199	ARG
2	B	290	SER
2	B	295	LYS
2	B	423	TYR
2	B	431	GLN
2	C	25	ASP
2	C	110	ASP
2	C	116	ASN
2	C	262	TRP
2	C	277[A]	ARG
2	C	277[B]	ARG

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type
2	C	290	SER
2	C	423	TYR
2	C	483	GLU
2	C	495	ARG
2	C	581	LEU
2	C	593	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	ALY	A	640	1	10,11,12	0.45	0	7,12,14	0.18	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	ALY	A	640	1	-	0/9/10/12	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [\(i\)](#)

Of 17 ligands modelled in this entry, 1 is monoatomic - leaving 16 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	A1AV1	B	706	-	30,32,35	0.91	2 (6%)	39,46,50	1.17	2 (5%)
3	GOL	C	703	-	5,5,5	0.32	0	5,5,5	0.34	0
6	A1AV1	A	707	-	30,32,35	0.90	2 (6%)	39,46,50	1.15	2 (5%)
4	SO4	A	704	-	4,4,4	0.59	0	6,6,6	0.07	0
3	GOL	C	701	-	5,5,5	0.30	0	5,5,5	0.35	0
4	SO4	B	705	-	4,4,4	0.61	0	6,6,6	0.06	0
3	GOL	B	703	-	5,5,5	0.34	0	5,5,5	0.17	0
3	GOL	A	702	-	5,5,5	0.28	0	5,5,5	0.50	0
3	GOL	A	701	-	5,5,5	0.31	0	5,5,5	0.31	0
3	GOL	B	701	-	5,5,5	0.31	0	5,5,5	0.30	0
3	GOL	B	702	-	5,5,5	0.30	0	5,5,5	0.37	0
3	GOL	A	703	-	5,5,5	0.30	0	5,5,5	0.32	0
4	SO4	B	704	-	4,4,4	0.60	0	6,6,6	0.07	0
6	A1AV1	C	704	-	30,32,35	0.88	2 (6%)	39,46,50	1.21	2 (5%)
3	GOL	C	702	-	5,5,5	0.32	0	5,5,5	0.43	0
4	SO4	A	705	-	4,4,4	0.61	0	6,6,6	0.06	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	A1AV1	B	706	-	-	6/17/28/31	0/5/5/5
3	GOL	C	703	-	-	2/4/4/4	-
6	A1AV1	A	707	-	-	3/17/28/31	0/5/5/5
3	GOL	C	701	-	-	2/4/4/4	-
3	GOL	B	703	-	-	1/4/4/4	-
3	GOL	A	702	-	-	2/4/4/4	-
3	GOL	A	701	-	-	4/4/4/4	-
3	GOL	B	701	-	-	4/4/4/4	-
3	GOL	B	702	-	-	0/4/4/4	-
3	GOL	A	703	-	-	4/4/4/4	-
6	A1AV1	C	704	-	-	6/17/28/31	0/5/5/5
3	GOL	C	702	-	-	2/4/4/4	-

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	706	A1AV1	C4-C5	-2.97	1.47	1.51
6	A	707	A1AV1	C4-C5	-2.92	1.47	1.51
6	C	704	A1AV1	C4-C5	-2.85	1.47	1.51
6	B	706	A1AV1	C8-C4	-2.50	1.36	1.39
6	C	704	A1AV1	C8-C4	-2.39	1.36	1.39
6	A	707	A1AV1	C8-C4	-2.38	1.36	1.39

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	707	A1AV1	C8-C4-C5	4.92	134.76	129.00
6	B	706	A1AV1	C8-C4-C5	4.80	134.62	129.00
6	C	704	A1AV1	C8-C4-C5	4.68	134.48	129.00
6	C	704	A1AV1	C19-C20-C17	-2.42	99.95	102.02
6	A	707	A1AV1	C19-C20-C17	-2.29	100.07	102.02
6	B	706	A1AV1	C19-C20-C17	-2.03	100.29	102.02

There are no chirality outliers.

All (36) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	701	GOL	O1-C1-C2-C3
3	A	701	GOL	C1-C2-C3-O3
3	A	702	GOL	O1-C1-C2-C3

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	703	GOL	O1-C1-C2-C3
3	A	703	GOL	C1-C2-C3-O3
3	A	703	GOL	O2-C2-C3-O3
3	B	701	GOL	C1-C2-C3-O3
3	C	702	GOL	O1-C1-C2-C3
3	C	703	GOL	O1-C1-C2-C3
6	A	707	A1AV1	C8-C4-C5-C6
6	B	706	A1AV1	C8-C4-C5-C6
6	B	706	A1AV1	C8-C4-C5-C7
6	C	704	A1AV1	C8-C4-C5-C6
3	A	701	GOL	O1-C1-C2-O2
3	B	701	GOL	O1-C1-C2-C3
3	C	701	GOL	O1-C1-C2-C3
3	A	701	GOL	O2-C2-C3-O3
3	A	702	GOL	O1-C1-C2-O2
3	A	703	GOL	O1-C1-C2-O2
3	B	701	GOL	O1-C1-C2-O2
3	B	701	GOL	O2-C2-C3-O3
3	C	703	GOL	O1-C1-C2-O2
3	C	702	GOL	O1-C1-C2-O2
6	B	706	A1AV1	C13-C14-C15-C22
6	B	706	A1AV1	C23-C14-C15-C16
6	B	706	A1AV1	C23-C14-C15-C22
3	B	703	GOL	C1-C2-C3-O3
6	C	704	A1AV1	C8-C4-C5-C7
6	B	706	A1AV1	C13-C14-C15-C16
6	C	704	A1AV1	C23-C14-C15-C22
6	C	704	A1AV1	C23-C14-C15-C16
6	C	704	A1AV1	C13-C14-C15-C22
6	C	704	A1AV1	C13-C14-C15-C16
3	C	701	GOL	O1-C1-C2-O2
6	A	707	A1AV1	C13-C14-C15-C22
6	A	707	A1AV1	C23-C14-C15-C22


There are no ring outliers.


2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	702	GOL	1	0
4	B	704	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	656/694 (94%)	-0.28	2 (0%) 94 93	32, 46, 83, 110	0
2	B	536/694 (77%)	-0.23	5 (0%) 84 82	35, 50, 84, 126	0
2	C	656/694 (94%)	-0.06	34 (5%) 27 26	32, 48, 115, 154	0
All	All	1848/2082 (88%)	-0.19	41 (2%) 62 60	32, 47, 103, 154	0

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	629	LEU	5.4
2	C	596	PHE	4.9
2	C	630	VAL	4.8
2	C	10	VAL	4.7
2	C	564	ILE	4.5
2	C	632	ASP	4.3
2	C	569	VAL	4.1
2	B	559	VAL	4.0
2	C	607	SER	3.9
2	C	592	MET	3.8
2	B	544	ASP	3.3
2	B	554	LEU	3.3
2	C	673	LYS	3.2
2	C	589	PHE	3.2
2	C	633	LEU	3.2
2	C	641	ILE	3.1
2	C	588	ALA	3.1
2	C	21	PRO	3.0
2	B	547	ILE	2.8
2	C	606	LEU	2.7
1	A	608	LYS	2.7
2	C	672	VAL	2.7
2	C	563	LEU	2.7

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	C	591	THR	2.5
2	C	594	PRO	2.5
2	C	574	VAL	2.4
2	C	676	VAL	2.4
2	C	609	GLU	2.3
2	C	566	HIS	2.3
2	C	593	LYS	2.3
2	C	590	VAL	2.3
2	C	568	GLY	2.3
2	B	558	GLU	2.3
2	C	651	ALA	2.2
2	C	25	ASP	2.2
2	C	294	GLY	2.1
2	C	628	TYR	2.1
2	C	613	GLN	2.1
2	C	8	PRO	2.1
2	C	612	ILE	2.0
1	A	665	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	ALY	A	640	12/13	0.96	0.14	49,57,62,64	0

## 6.3 Carbohydrates [\(i\)](#)


There are no monosaccharides in this entry.

## 6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	CL	A	706	1/1	0.64	0.13	81,81,81,81	0
6	A1AV1	B	706	28/31	0.78	0.52	54,85,93,98	0
3	GOL	A	702	6/6	0.80	0.15	61,63,66,70	0
3	GOL	B	703	6/6	0.81	0.21	59,61,65,68	0
6	A1AV1	C	704	28/31	0.84	0.35	50,66,78,80	0
3	GOL	B	701	6/6	0.85	0.34	57,60,62,66	0
3	GOL	B	702	6/6	0.85	0.15	66,67,68,69	0
3	GOL	C	702	6/6	0.86	0.19	50,60,64,69	0
3	GOL	C	701	6/6	0.89	0.17	54,61,65,68	0
6	A1AV1	A	707	28/31	0.89	0.20	46,60,65,68	0
3	GOL	A	703	6/6	0.90	0.15	52,55,62,64	0
3	GOL	A	701	6/6	0.90	0.15	49,54,60,67	0
3	GOL	C	703	6/6	0.91	0.18	52,55,57,58	0
4	SO4	B	705	5/5	0.96	0.13	79,81,93,98	0
4	SO4	B	704	5/5	0.97	0.18	80,84,87,92	0
4	SO4	A	704	5/5	0.97	0.17	61,64,73,83	0
4	SO4	A	705	5/5	0.97	0.15	58,62,79,80	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around A1AV1 C 704:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around A1AV1 A 707:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.

For Manuscript Review