

Detailed Methods

Group A and Group B received telerehabilitation (TR) via Tencent Meeting video software under real-time guidance from speech therapists, while Group C completed offline therapy in a clinical setting with speech therapists supervision. The TR groups differed only in the intensity of ORLA protocols. Speech therapists pre-assessed individual aphasia severity levels and curated personalized digital materials in advance. These materials, designed around daily life-related content (e.g., personal routines, familial interactions), were structured into a three-tiered difficulty progression: The difficulty coefficient ranged from 3-5 words for simple short sentences to 8-12 words for medium-long sentences or gradually expanded to complex sentences (15-30 words), short texts, etc.

The specific steps were as follows (ORLA):

- ① The therapist reads the training materials loudly and points out each word while reading;
- ② The therapist reads the training materials again and asks the patient to point out each word while listening;
- ③ The therapist randomly reads out a word or phrase in a sentence (either a real word or a function word) and asks the patient to identify and point it out;
- ④ The therapist randomly points out a word or phrase in the sentence (which can be a content word such as a noun or verb or a function word such as a pronoun, preposition, or conjunction) and asks the patient to read it aloud;
- ⑤ The therapist reads sentences aloud with the patient, mainly focusing on the patient. The therapist adjusts their speaking speed and volume based on the patient's specific ability level (for example, by slightly reading out the sound before the patient so that the patient can hear the initial phoneme of the word; as the patient needs fewer clues, the volume decreases), while reading aloud, asking the patient to recognize and point out each word and phrase;
- ⑥ Please ask the patient to read the complete sentence aloud, pointing out each word while reading;
- ⑦ Please ask the patient to read the complete sentence aloud again.

Supplemental Tables

Table S1. Subitems paired comparison 1

CCread	TIME	TIME	M.D	[95% CI]	Std.Error	df	P Value
CWread	1.00	2.00	-9.26*	[-11.00, -7.52]	0.87	42.05	0.000
		3.00	-9.98*	[-11.72, -8.23]	0.87	44.99	0.000
	2.00	3.00	-0.71*	[-1.38, -0.05]	0.34	26.64	0.037
CWname	1.00	2.00	-1.13*	[-1.35, -0.91]	0.11	48.82	0.000
		3.00	-1.52*	[-1.75, -1.29]	0.11	53.35	0.000
	2.00	3.00	-0.39*	[-0.60, -0.18]	0.10	15.34	0.001
CWfluen	1.00	2.00	-0.93*	[-1.26, -0.60]	0.16	42.05	0.000
		3.00	-0.98*	[-1.32, -0.65]	0.17	44.99	0.000
	2.00	3.00	-0.05	[-0.25, 0.14]	0.09	26.64	0.574
CWdisco	1.00	2.00	-1.10*	[-1.38, -0.81]	0.14	42.93	0.000
		3.00	-1.36*	[-1.64, -1.08]	0.14	44.17	0.000
	2.00	3.00	-0.26*	[-0.41, -0.12]	0.07	35.89	0.001
CWlisten	1.00	2.00	-0.98*	[-1.25, -0.70]	0.14	45.92	0.000
		3.00	-1.21*	[-1.49, -0.94]	0.14	46.60	0.000
	2.00	3.00	-0.24*	[-0.42, -0.06]	0.09	30.04	0.011

Bold font indicates *P value* less than 0.05; * indicates the data has statistical significance; MD,Mean difference;CCread, CRRCAE-reading subitem; CWread, WAB-reading subitem; CWname, WAB-naming subitem; CWfluen, WAB-fluency subitem; CWdisco, WAB-discourse subitem; CWlisten, WAB-listening comprehension subitem.

Table S2. Subitems paired comparison 2

CCread	GROUP	GROUP	M.D	[95% CI]	Std.Error	df	P Value
CWread	C	B	24.17*	[17.14, 31.19]	3.49	46.41	0.000
		A	0.37	[-6.66, 7.40]	3.49	46.41	0.916
	B	A	-23.80*	[-30.82, -16.77]	3.49	46.41	0.000
CWname							
CWname	C	B	2.09*	[1.50, 2.68]	0.29	25.95	0.000
		A	0.09	[-0.50, 0.67]	0.29	25.95	0.761
	B	A	-2.00*	[-2.59, -1.42]	0.29	25.95	0.000
CWfluen							
CWfluen	C	B	1.09*	[0.28, 1.90]	0.40	38.34	0.000
		A	0.24	[-0.57, 1.05]	0.40	38.34	0.761
	B	A	-0.85*	[-1.66, -0.04]	0.40	38.34	0.000
CWdisco							
CWdisco	C	B	1.21*	[0.38, 2.05]	0.42	41.58	0.006
		A	-0.38	[-1.22, 0.46]	0.42	41.58	0.364
	B	A	-1.60*	[-2.43, -0.76]	0.42	41.58	0.000
CWlisten							
CWlisten	C	B	1.34*	[0.47, 2.21]	0.43	37.34	0.004
		A	0.03	[-0.85, 0.90]	0.43	37.34	0.954
	B	A	-1.32*	[-2.19, -0.44]	0.43	37.34	0.004

Bold font indicates *P value* less than 0.05; * indicates the data has statistical significance; MD,Mean difference;CCread, CRRCAE-reading subitem; CWread, WAB-reading subitem; CWname, WAB-naming subitem; CWfluen, WAB-fluency subitem; CWdisco, WAB-discourse subitem; CWlisten, WAB-listening subitem; Group[A], moderate-to-high-intensity ORLA telerehabilitation group; Group[B], low-intensity ORLA telerehabilitation group;Group[C],moderate-to-high-intensity ORLA offline group.

Table S3. CWread interaction comparison

<i>predictors</i>	MD in CWread	[95% CI]	Std.Error	T Value	<i>P Value</i>
(Intercept)	3.89	[3.43, 4.34]	0.23	17.20	0.000
Group[C]	0.07	[-0.57, 0.71]	0.32	0.21	0.833
Group[B]	-2.29	[-2.93, -1.64]	0.32	-7.14	0.000
TIME[1]	-1.72	[-2.12, -1.33]	0.20	-8.74	0.000
TIME[2]	-0.49	[-0.86, -0.12]	0.17	-2.82	0.013
Group[C]*TIME[1]	0.01	[-0.55, 0.56]	0.28	0.02	0.984
Group[C]*TIME[2]	0.05	[-0.47, 0.57]	0.25	0.22	0.830
Group[B]*TIME[1]	0.60	[0.04, 1.16]	0.28	2.16	0.036
Group[B]*TIME[2]	0.25	[7.84, 24.12]	0.25	1.01	0.326

* indicates interaction; Bold font indicates *P value* less than 0.05; MD,Mean difference;CWread, WAB-reading subitem; TIME[1],Pre_Post=T0_T1; TIME[2],Pre_Post=T0_T2;T1, 4 weeks post-intervention; T2, 3 months post-intervention; T3, 6 months post-intervention; Group[B], low-intensity ORLA telerehabilitation group;Group[C],moderate-to-high-intensity ORLA offline group.

Table S4. CWname interaction comparison

<i>predictors</i>	MD in CWname	[95% CI]	Std.Error	T Value	<i>P Value</i>
(Intercept)	3.06	[2.47,3.64]	0.29	10.55	0.000
Group[C]	0.28	[-0.54,1.11]	0.41	0.69	0.492
Group[B]	-0.80	[-1.63,0.02]	0.41	-1.96	0.056
TIME[1]	-0.96	[-1.55, -0.38]	0.29	-3.32	0.002
TIME[2]	-0.02	[-0.32,0.35]	0.16	0.11	0.910

Bold font indicates *P value* less than 0.05; MD,Mean difference; CWname, WAB-naming subitem;TIME[1],Pre_Post=T0_T1; TIME[2],Pre_Post=T0_T2;T1, 4 weeks post-intervention; T2, 3 months post-intervention; Group[B], low-intensity ORLA telerehabilitation group;Group[C],moderate-to-high-intensity ORLA offline group.

Table S5. CWfluen interaction comparison

<i>predictors</i>	MD in CWfluen	[95% CI]	Std.Error	T Value	<i>P Value</i>
(Intercept)	3.36	[2.77,3.94]	0.29	11.58	0.000
Group[C]	-0.36	[-1.19,0.47]	0.41	-0.87	0.389
Group[B]	-1.71	[-2.54, -0.89]	0.41	-4.18	0.000
TIME[1]	-1.50	[-1.99, -1.01]	0.24	-6.23	0.000
TIME[2]	-0.21	[-0.47,0.04]	0.13	-1.71	0.097

Bold font indicates *P value* less than 0.05; MD,Mean difference;CWfluen, WAB-fluency subitem;TIME[1],Pre_Post=T0_T1; TIME[2],Pre_Post=T0_T2;T1, 4 weeks post-intervention; T2, 3 months post-intervention; Group[B], low-intensity ORLA telerehabilitation group;Group[C],moderate-to-high-intensity ORLA offline group.

Table S6. CWdisco interaction comparison

<i>predictors</i>	MD in CWdisco	[95% CI]	Std.Error	T Value	<i>P Value</i>
(Intercept)	2.64	[2.77, 3.94]	0.31	8.49	0.000
Group[C]	-0.07	[-1.19, 0.47]	0.44	-0.16	0.872
Group[B]	-1.43	[-2.54, -0.89]	0.44	-3.25	0.002
TIME[1]	-1.64	[-1.99, -1.01]	0.22	-7.30	0.000
TIME[2]	-0.29	[-0.47, 0.04]	0.16	-1.80	0.083
Group[C]*TIME[1]	0.57	[-0.62, 0.76]	0.32	1.80	0.079
Group[C]*TIME[2]	0.07	[-0.50, 0.22]	0.22	0.32	0.753
Group[B]*TIME[1]	0.71	[-0.33, 1.04]	0.32	2.25	0.030
Group[B]*TIME[2]	0.07	[-0.36, 0.36]	0.22	0.32	0.753

* indicates interaction;Bold font indicates *P value* less than 0.05; MD,Mean difference; CWdisco, WAB-discourse subitem;TIME[1],Pre_Post=T0_T1;TIME[2],Pre_Post=T0_T2; T1, 4 weeks post-intervention; T2, 3 months post-intervention; Group[B], low-intensity ORLA telerehabilitation group;Group[C],moderate-to-high-intensity ORLA offline group.

Table S7. CWlisten interaction comparison

<i>predictors</i>	MD in CWlisten	[95% CI]	Std.Error	T Value	<i>P Value</i>
(Intercept)	3.03	[2.25, 3.80]	0.38	7.89	0.000
Group[C]	0.47	[-0.63, 1.56]	0.54	0.86	0.395
Group[B]	-0.54	[-1.64, 0.55]	0.54	-1.00	0.322
TIME[1]	-0.96	[-1.55, -0.37]	0.29	-3.28	0.002
TIME[2]	-0.01	[-0.29, 0.26]	0.13	-0.11	0.916

Bold font indicates *P value* less than 0.05; MD,Mean difference; CWlisten, WAB-listening Comprehension subitem;TIME[1],Pre_Post=T0_T1;TIME[2],Pre_Post=T0_T2; T1, 4 weeks post-intervention; T2, 3 months post-intervention; Group[B], low-intensity ORLA telerehabilitation group;Group[C],moderate-to-high-intensity ORLA offline group.