

Table S3: Treatment options for different pathological types

	Total (N=1023)	DLBCL (N=668)	MALT (N=208)	FL (N=24)	MCL (N=38)	BL (N=35)	EALT (N=13)	NKT (N=16)	Other T (N=21)	P
Treatment,N (%)	< .001*									
Supportive care	75 (7%)	27 (5%)	37 (18%)	2 (8%)	2 (6%)	0 (0%)	0 (0%)	1 (6%)	6 (29%)	
Chemotherapy	474 (47%)	327 (50%)	75 (37%)	14 (58%)	23 (64%)	26 (74%)	1 (8%)	2 (12%)	6 (29%)	
Radiotherapy	27 (3%)	2 (0%)	25 (13%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	
Surgery	92 (9%)	49 (7%)	24 (12%)	2 (8%)	3 (8%)	3 (9%)	1 (8%)	5 (32%)	5 (24%)	
C+R	69 (7%)	45(7%)	20 (10%)	1 (4%)	1 (3%)	0 (0%)	0 (0%)	1 (6%)	1 (5%)	
R+S	1 (0%)	1 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	
C+S	255 (26%)	198 (30%)	20 (10%)	4 (17%)	7 (19%)	6 (17%)	10 (84%)	7 (44%)	3 (13%)	
C+R+S	8 (1%)	7 (1%)	0 (0%)	1 (5%)	0 (0%)	0 (0%)	0(0%)	0(0%)	0 (0%)	

*: Simulated p-value

C: chemotherapy, R: radiotherapy, S: surgery