

Table S2: Locations of disease corresponding to different pathological types

	Primarysite, N (%)						Total
	Esophagus	Stomach	Small intestine	Ileocecus	Large intestine	Multiregions	
B-cell lymphoma							973 (95.10)
DLBCL	2 (0.30)	369 (55.24)	77 (11.52)	66 (9.88)	123 (18.42)	31 (4.64)	668 (65.30)
MALT	1 (0.48)	148 (71.15)	22 (10.58)	4 (1.92)	21 (10.10)	12 (5.77)	208 (20.33)
FL	0	9 (37.5)	9 (37.5)	0	4 (16.67)	2 (8.33)	24 (2.34)
MCL	0	1 (26.32)	3 (7.89)	6 (15.79)	11 (28.95)	8 (21.05)	38 (3.71)
BL	0	1 (42.86)	5 (14.29)	5 (14.29)	5 (14.29)	5 (14.29)	35 (3.42)
T-cell lymphoma							50 (4.90)
EALT	0	1 (7.70)	9 (69.22)	0	2 (15.38)	1 (7.70)	13 (1.28)
NK/T	0	2 (12.50)	1 (6.25)	2 (12.50)	10 (62.5)	1 (6.25)	16 (1.56)
Other T-cell lymphoma	1 (4.76)	6 (28.57)	8 (38.10)	2 (9.52)	1 (4.76)	3 (14.29)	21 (2.06)
Total	4 (0.39)	560 (54.74)	134 (13.10)	85 (8.31)	177 (17.30)	63 (6.16)	1023