

Assessing the impact of migration-related stressors on refugees' gut microbiome

Giorgia Palladino^{1,2}, Marianna Marangi³, Daniel Scicchitano^{1,2}, Silvia Turroni^{1,4}, Simone Rampelli^{1,2}, Marco Candela^{1,2*}

¹Unit of Microbiome Science and Biotechnology, Department of Pharmacy and Biotechnology, Alma Mater Studiorum - University of Bologna, Via Belmeloro 6, 40126 Bologna, Italy

²Fano Marine Center, The Inter-Institute Center for Research on Marine Biodiversity, Resources and Biotechnologies, Viale Adriatico 1, 61032, Fano, Italy

³Department of Clinical and Experimental Medicine, University of Foggia, Foggia, Italy

⁴IRCCS Azienda Ospedaliero-Universitaria di Bologna, 40138 Bologna, Italy

*corresponding author: marco.candela@unibo.it

Supplementary information

Supplementary Table 1 - Overlapping gut microbiome compositional features between refugees and rural populations. Refugee gut microbiome genera that were similarly represented in rural populations but significantly decreased in urban populations are shown (Wilcoxon rank-sum test corrected with FDR, p-values ≤ 0.05).

Bacterial genera	P-values	
	Refugees vs rural	Refugees vs urban
Alloprevotella	> 0.05	< 0.001
Prevotella	> 0.05	< 0.001
Prevotella 7	> 0.05	< 0.01
Streptococcus	> 0.05	< 0.05
Ruminococcus gnavus group	> 0.05	< 0.001
Dorea	> 0.05	< 0.001
Lachnospiraceae	> 0.05	< 0.001
Tyzzerella	> 0.05	< 0.001
Romboutsia	> 0.05	< 0.05
Succinivibrio	> 0.05	< 0.001
Akkermansia	> 0.05	< 0.001