

TABLE 1 Baseline characteristics of the patients with FL (N=118)

Characteristic	n (%)
Age (year), median (range)	
<60	33
≥60	85
Gender	
Female	53
Male	65
Extranodal involvement ≥ 2	
Yes	47
No	71
B symptoms	
Yes	42
No	76
Bone marrow involvement	
Yes	55
No	63
Ann Arbor stage	
I/II	25
III/IV	93
PRIMA-PI	
low and intermediate-risk group	47
high-risk group	71
Histological grade	
1-3A	78
3B+composite	40
FLIPI-2	
<3	57
≥3	61

TABLE 1 Continued

Characteristic	n (%)
HB (g/L) <120	
Yes	59
No	59
ALB (g/L) <40	
Yes	70
No	48
LDH (U/L) >UNL	
Yes	58
No	60
β ₂ -MG (mg/L) >UNL	
Yes	62
No	56
CD4 ⁺ T cells	
<30%	62
≥30%	56
NK cells	
<8%	56
≥8%	62
Ki-67	
<30%	36
≥30%	82
Bcl-2	
positive	111
negative	7
Bcl-6	
positive	114
negative	4

TABLE 1 Continued

Characteristic	n (%)
CD5	
positive	57
negative	61
CD21	
positive	114
negative	4
CD10	
positive	68
negative	50
CD20	
positive	118
negative	0
MUM1	
positive	65
negative	53

TABLE 2 Overall survival and progression-free survival in univariate analysis in follicular lymphoma

Influencing factors	OS rate in 3 years with follow-up (%)	<i>P</i>	PFS rate in 3 years with follow-up (%)	<i>P</i>
Pathological grade		<0.001		<0.001
1-3A	91.2%		76.5%	
3B+composite	68.7%		26.8%	
B symptoms		<0.001		0.001
Yes	70.3%		36.5%	
No	90.5%		70.9%	
PRIMA-PI		0.006		<0.001
LR and IR	93.5%		84.6%	
HR	76.1%		41.1%	
FLIPI-2		<0.001		0.002
<3	96.1%		80.8%	
≥3	72.0%		39.0%	
HB		0.003		0.016
<120g/L	77.2%		42.9%	
≥120g/L	87.1%		71.9%	
ALB		<0.001		0.002
<40g/L	74.2%		43.9%	
≥40g/L	97.7%		83.3%	
LDH		0.049		0.003
Normal	89.7%		72.3%	
Higher	75.5%		44.7%	

TABLE 2 Continued

Influencing factors	OS rate in 3 years with follow-up (%)	<i>P</i>	PFS rate in 3 years with follow-up (%)	<i>P</i>
β ₂ -MG		0.038		0.031
Normal	89.4%		69.2%	
Higher	76.3%		47.5%	
CD4 ⁺ T cells		0.001		<0.001
<30%	74.8%		42.1%	
≥30%	92.6%		78.7%	
NK cells		<0.001		<0.001
<8%	69.5%		35.6%	
≥8%	95.3%		77.8%	
Ki-67		0.005		0.001
<30%	91.5%		87.6%	
≥30%	78.9%		46.2%	
CD5		<0.001		<0.001
positive	71.9%		34.3%	
negative	93.4%		81.6%	
CD10		<0.001		<0.001
positive	94.0%		77.0%	
negative	69.9%		35.7%	
MUM1		0.001		0.001
positive	73.3%		41.5%	
negative	94.5%		79.4%	

TABLE 3 Overall survival and progression-free survival in multivariate Cox analysis in follicular lymphoma

Influencing factors	OS			PFS		
	HR	95%CI	<i>P</i>	HR	95%CI	<i>P</i>
Pathological grade 3B+composite	1.777	0.528-5.984	0.353	2.933	1.080-7.963	0.035
NK cells<8%	0.067	0.010-0.435	0.005	0.705	0.149-3.340	0.659