

Cholinergic Basal Forebrain Atrophy Accelerates Cognitive Decline via Cortical Thinning: The Moderating Role of Amyloid- β Pathology in Preclinical Alzheimer's Disease

Wencai Ding

dingneurology@wnmc.edu.cn

The Second Affiliated Hospital of Wannan Medical College <https://orcid.org/0000-0002-7800-500X>

Cen Si

The Second Affiliated Hospital of Wannan Medical College

Lijuan Wang

Meiling Qiu

Zhongqiang Xu

Li Xu

Rui Bao

Xiaolei Tang

Juanyu Gong

Jinting Wu

Zhiding Shao

Tonghua Zhang

Fan Yang

Article

Keywords:

Posted Date: May 25th, 2026

DOI: <https://doi.org/10.21203/rs.3.rs-6220536/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Additional Declarations: The authors have declared there is **NO** conflict of interest to disclose

Abstract

Degeneration of the cholinergic basal forebrain (cBF) is a hallmark of early neurodegeneration in Alzheimer's disease (AD). While cBF atrophy has been linked to cognitive decline, the extent to which cortical thinning mediates this association, particularly in preclinical AD, remains unclear. Furthermore, the influence of amyloid- β (A β) pathology on these relationships warrants further investigation. This study analyzed longitudinal MRI and PIB-PET data from 230 cognitively normal older adults in the Harvard Aging Brain Study, with a mean follow-up of six years. FreeSurfer was used to quantify cBF volume and cortical thickness, while cognitive performance was assessed using the Preclinical Alzheimer Cognitive Composite-5 (PACC5). Linear mixed-effects models evaluated longitudinal associations between cBF atrophy, cortical thinning, and cognitive decline. Mediation analyses assessed whether cortical thinning mediated the relationship between cBF degeneration and cognitive decline, and the moderating effect of A β burden was examined. Progressive cortical thinning in multiple cognition-related regions was significantly associated with longitudinal cBF atrophy. Mediation analysis revealed that cortical thinning accounted for approximately 44% of the association between cBF degeneration and cognitive decline. These effects were more pronounced in A β -positive individuals, suggesting a synergistic interaction between amyloid pathology and cholinergic degeneration. These findings indicate that cBF atrophy accelerates cognitive decline by driving cortical thinning, with A β pathology further exacerbating these effects. This study enhances understanding of the structural mechanisms underlying cognitive decline in preclinical AD and highlights potential early intervention targets.

Introduction

Individuals with Alzheimer's disease (AD) typically experience an extended period without cognitive symptoms before noticeable decline emerges[1]. Although definitions for preclinical AD vary, the widely accepted criterion includes normal cognitive function accompanied by at least one biomarker indicative of AD pathology, such as amyloid-beta (A β) deposits or tau pathology[2, 3]. However, amyloid plaque accumulation alone does not necessarily lead to Alzheimer's disease, as it is also associated with the normal aging process. Therefore, relying solely on amyloid biomarkers is insufficient for fully characterizing preclinical AD[4, 5].

Brain atrophy, particularly in areas like the cholinergic basal forebrain (cBF), represents a hallmark pathological characteristic of Alzheimer's disease[6]. Recent advancements in automated MRI techniques have facilitated the identification of in vivo neuroimaging biomarkers for basal forebrain atrophy[7, 8]. In individuals with mild cognitive impairment (MCI), atrophy of the basal forebrain cholinergic system has been associated with cortical thinning, especially in projection regions like the parahippocampal gyrus and cBF [9]. Research on Parkinson's disease has demonstrated a correlation between basal forebrain atrophy, cortical hypometabolism, and cognitive decline[10]. In early stages of AD, basal forebrain atrophy combined with amyloid deposition significantly contributes to cognitive impairment[11, 12]. Therefore, imaging techniques such as MRI assessments of basal forebrain volume

and cortical atrophy might identify early neurodegenerative changes, potentially aiding early diagnosis and interventions for preclinical AD.

Cortical thinning is also closely linked to cognitive decline. For instance, decreased cortical thickness in the left parahippocampal cortex has been associated with various cognitive functions[13]. Likewise, cortical thinning is observed in Parkinson's disease patients at early stages, even before noticeable cognitive impairment develops, suggesting its potential role as an early indicator of cognitive decline[14]. Distinct patterns of cortical atrophy have also been observed across various MCI subtypes over three years. Memory-deficit MCI patients initially experience rapid atrophy in medial temporal areas, whereas naming/memory-deficit subtypes primarily show temporal lobe involvement, and mixed subtypes exhibit widespread cortical atrophy[15]. However, previous studies have been limited by methodological constraints, such as cross-sectional designs, small sample sizes, and single-center recruitment. These limitations have impeded a thorough investigation of the relationships among basal forebrain volume, cortical thinning, and cognitive function.

The current study addresses these limitations by utilizing a large multicenter cohort and longitudinal design to investigate these variables systematically over time. We hypothesize that cBF volume influences cognitive function through its effect on cortical thinning in specific brain regions. By exploring early-stage neurodegenerative changes, this study provides novel perspectives on the mechanisms that underlie cognitive decline in preclinical AD.

Materials

Study Design and Participants

This study enrolled 230 cognitively normal older adults from the Harvard Aging Brain Study (HABS), a prospective longitudinal cohort designed to investigate early neurodegenerative changes associated with preclinical Alzheimer's disease. Conducted in collaboration between Massachusetts General Hospital and Brigham and Women's Hospital in Boston, Massachusetts, the study aimed to enhance the understanding of early disease markers[16]. HABS integrates multimodal neuroimaging techniques to capture structural and functional brain alterations, including amyloid and tau pathology. These imaging assessments are complemented by detailed neuropsychological evaluations to track cognitive performance over time. The primary goal of these assessments is to identify biomarkers that could serve as early indicators of cognitive decline. The study received ethical approval from the Institutional Review Board (IRB) at Mass General Brigham, and all participants provided written informed consent. The research protocol complies with the ethical guidelines established in the Declaration of Helsinki.

Neuropsychological Evaluation

Participants were identified as cognitively normal based on comprehensive neuropsychological evaluations. Inclusion criteria required the absence of clinical depression, defined as a Geriatric

Depression Scale score of ≤ 10 , with no history of active psychiatric disorders. Additionally, cognitive performance was evaluated using the Mini-Mental State Examination (MMSE), with a minimum score threshold of 25. Episodic memory was assessed using the Wechsler Logical Memory II delayed recall test, with participants required to score within 1.5 standard deviations of age- and education-adjusted norms. The Clinical Dementia Rating (CDR) scale was administered by trained clinicians blinded to biomarker status[16]. Global cognitive function was measured using the Preclinical Alzheimer Cognitive Composite-5 (PACC5), a composite index that integrates assessments across multiple cognitive domains, including episodic and semantic memory, executive function, and global cognition. The composite consists of the Mini-Mental State Examination (MMSE), Wechsler Memory Scale-Revised Logical Memory Delayed Recall (WMS-R LMDR), Digit-Symbol Coding Test (DSC), Free and Cued Selective Reminding Test-Free Recall (FCSRT96), and Category Fluency (CAT)[17]. The PACC5 serves as a sensitive tool for identifying early cognitive decline, beyond assessments of individual cognitive domains, and has been specifically designed to track subtle cognitive changes associated with A β pathology[17]. Participants underwent yearly cognitive assessments, incorporating evaluations of episodic memory, executive functioning, and global cognition, in conjunction with CDR scoring. Baseline cognitive assessments were conducted within a year of the initial MRI session.

Imaging Acquisition and Processing

Structural MRI scans were acquired using a 3T scanner, following standardized imaging protocols to ensure consistency among participants. High-resolution T1-weighted images were obtained with a magnetization-prepared rapid gradient echo (MPRAGE) sequence. Imaging parameters included an isotropic voxel size of 1.0 mm³, an echo time (TE) of approximately 3.2 ms, a repetition time (TR) of ~ 2.2 s, and an inversion time (TI) of ~ 1.1 s. Parallel imaging techniques were employed to optimize signal quality and minimize scan duration. To assess amyloid- β (A β) burden, participants underwent positron emission tomography (PET) with Pittsburgh Compound B (¹¹C-PIB)[18]. The cerebellar gray matter served as a reference region for computing standardized uptake value ratios (SUVR). Data preprocessing included motion correction and partial volume correction (PVC), utilizing validated imaging pipelines to enhance quantification accuracy.

Basal Forebrain Volume

The cBF volume was used as an indicator of neuronal degeneration in cognitively normal older adults. Segmentation of the basal forebrain, was conducted on 3D T1-weighted MRI scans using the *ScLimbic* pipeline (<https://surfer.nmr.mgh.harvard.edu/fswiki/ScLimbic>)[19]. The total cBF volume was derived by summing the left and right basal forebrain volumes, as computed within each participant's native T1-weighted structural MRI space. Following the execution of the *ScLimbic* pipeline, we conducted a visual inspection of the segmented images to detect any substantial inaccuracies in structure labeling. Furthermore, Additionally, the `-write_qa_stats` function in FreeSurfer's *ScLimbic* pipeline was utilized to generate comprehensive quality assurance metrics.

Cortical Thickness

Cortical thickness was measured using FreeSurfer software (version 6.0.0). The software calculates cortical thickness by determining the average distance between the gray-white matter boundary and the pial surface across the cortex[20, 21]. Image processing included several standard preprocessing steps: removing non-brain tissues, spatial normalization to a standard brain template, atlas registration, and parcellation using participant-specific brain templates. After these steps, we computed cortical thickness maps for each participant[22]. We visually inspected all individual cortical maps to ensure accuracy before including them in further analyses. To adjust for variations in head size, we adjusted cortical thickness measures using intracranial volume (ICV) estimated from FreeSurfer. Specifically, we applied the residual adjustment method, which corrects thickness measures based on variations in ICV [21, 23, 24]. The cortical thickness maps were aligned to the fsaverage standard template and processed with spatial smoothing using a Gaussian kernel with a 10 mm full-width at half-maximum (FWHM). Smoothing reduced variability between individuals and improved the reliability of statistical comparisons. This approach is consistent with best practices recommended for neuroimaging analyses.

PIB-PET methods and quantification

Amyloid- β ($A\beta$) burden was assessed through positron emission tomography (PET) using Pittsburgh Compound B (^{11}C -PIB). The scans were acquired at Massachusetts General Hospital with an ECAT EXACT HR+ scanner (Siemens, Erlangen, Germany)[16]. After an initial transmission scan, participants received an intravenous bolus injection of approximately 10–15 mCi ^{11}C -PIB. A dynamic PET acquisition was conducted over a 60-minute period in three-dimensional mode, capturing 69 frames (12×15 s, 57×60 s) across 63 image planes with an axial field of view of 15.2 cm, trans-axial resolution of 5.6 mm, and slice thickness of 2.4 mm. To ensure alignment between PET images and anatomical structures, late-sum PIB-PET frames were co-registered to individual T1-weighted MRI scans using FreeSurfer's *mri_coreg* tool. Amyloid deposition was measured using Logan graphical analysis, with the cerebellar gray matter serving as the reference region for calculating the distribution volume ratio (DVR). Non-partial volume corrected (non-PVC) DVR maps were generated and projected onto each participant's cortical surface for regional analysis. $A\beta$ burden was derived from a composite cortical region of interest, including the frontal, lateral temporal, parietal, and retrosplenial cortices, collectively known as PIB-FLR. Participants were categorized as amyloid-positive ($A\beta+$) or amyloid-negative ($A\beta-$) at baseline based on a PIB-FLR DVR threshold of 1.2, previously determined through Gaussian mixture modeling[25]. See Fig. 1 for an overview of the study design and analytical procedures (Fig. 1).

Statistical analyses

Baseline differences in demographic, clinical, and cognitive characteristics between $A\beta+$ and $A\beta-$ groups were examined using independent-sample t-tests for continuous variables and chi-square tests

for categorical variables. The Shapiro–Wilk test was used to evaluate normality, and log transformation was applied to non-normally distributed data when necessary.

To investigate longitudinal changes in cBF volume, cortical thickness, and cognitive performance, we utilized linear mixed-effects models over the follow-up period. These models incorporated participant-specific random intercepts and slopes to account for repeated measurements. Age, gender, and educational attainment were incorporated as covariates in the subsequent analyses. Before processing cortical thickness data, the ComBat harmonization method was applied to mitigate potential confounding effects related to site variability, age, sex, and educational background. ComBat is an empirical Bayesian approach designed to minimize variability from these confounding factors while preserving relevant biological signals[26]. We conducted vertex-wise analyses separately for each cortical location, estimating the rate of change (slope) over time.

We conducted mediation analyses to determine whether regional cortical thinning serves as an intermediary in the relationship between cBF degeneration and cognitive decline in cognitively normal older adults. First, we examined the relationship between longitudinal cBF volume reduction and cognitive performance by computing individual slopes of change. We repeated this analysis for the relationship between cortical thinning and cognitive decline. We then identified cortical regions significantly associated with cognitive decline, weighting each region's cortical thinning slope according to its strength of association with cognition.

We conducted additional mediation analyses to further explore the relationships among basal forebrain atrophy, cortical thinning, and cognitive decline. First, we tested whether longitudinal reductions in basal forebrain volume were associated with cognitive decline. Next, we examined the relationship between cortical thinning in basal forebrain-related regions and cognitive performance using vertex-wise regression analyses. Rather than calculating a simple average cortical thinning across all significant regions, we weighted the cortical thinning slopes according to the strength of their association with cognitive decline. We conducted mediation analyses to assess whether cortical thinning acts as an intermediary in the association between cBF degeneration and cognitive decline. Specifically, we measured the average direct effect (ADE), representing the direct impact of basal forebrain degeneration on cognitive decline, and the average causal mediation effect (ACME), which captures the indirect effect mediated by cortical thinning. We used nonparametric bootstrapping with 10,000 resamples to determine the statistical significance of these effects and to estimate the proportion of mediation.

We conducted additional analyses to evaluate whether cortical thinning independently contributes to cognitive decline, separate from cBF degeneration. To investigate this relationship, we conducted vertex-wise regression analyses to evaluate the relationship between cortical thinning and longitudinal changes in PACC5 scores. We calculated the average cortical thinning effect across all significant vertices, both before and after adjusting for longitudinal changes in cBF volume. The primary measure of cognitive decline in these analyses was the slope of the PACC5 score change. Vertex-wise analyses were adjusted for multiple comparisons using permutation testing with 10,000 iterations, employing a two-tailed

cluster-forming threshold of $P < 0.05$. See the flowchart in Fig. 1 for the study design and analytical approach (Fig. 1). All statistical analyses were performed using R version 4.4.0. Linear mixed-effects models and mediation models were implemented utilizing the lme4 package and the mediation package, respectively.

Results

Demographic Characteristics

Table 1 summarizes the demographic, clinical, and cognitive characteristics of study participants. A total of 230 individuals were included, with 61 (26.5%) classified as A β + and 169 (73.5%) as A β - . The mean age of the A β + group was 74.7 years (SD = 5.37), and 55.7% were female. The frequency of the APOE- ϵ 4 allele was markedly greater in the A β + group (59.0%) than in the A β - group (16.6%, $P < 0.001$). Additionally, amyloid burden, measured by PIB-FLR DVR, was notably elevated in the A β + group (mean = 1.45, SD = 0.167) relative to the A β - group (mean = 1.08, SD = 0.0511, $P < 0.001$). Participants in the A β + group underwent MRI assessments for an average of 4.14 years (SD = 1.45), with approximately 2.69 follow-up scans (SD = 0.672). While PACC5 scores were marginally lower in the A β + group (mean = -0.0018, SD = 0.590) than in the A β - group, this difference did not reach statistical significance ($P = 0.42$). The A β - participants had a mean age of 72.9 years (SD = 6.32) and a slightly higher proportion of female participants (59.8%). Their PIB-FLR DVR values were significantly lower than those of the A β + group (mean = 1.08, SD = 0.0511, $P < 0.001$).

For the A β - group, the average MRI follow-up duration was 4.48 years (SD = 1.43), with an average of 2.80 MRI scans per participant (SD = 0.684). Mini-Mental State Examination (MMSE) scores were comparable between the two groups, with no significant difference (A β -: mean = 29.1, SD = 1.05; A β +: mean = 28.8, SD = 1.04, $P = 0.061$). Similarly, PACC5 scores did not significantly differ (A β -: mean = 0.0715, SD = 0.649; $P = 0.42$). Both groups underwent a similar number of cognitive assessments, averaging 5.77 assessments (SD = 0.500) in the A β - group.

Longitudinal Trajectories of cBF Atrophy, Cortical Thinning, and Cognitive Decline

At baseline, cBF volume and ICV showed no significant differences between A β + and A β - groups (Supplementary Fig. 1). However, all participants demonstrated significant declines in cBF volume over time ($\beta = -4.49$; 95% CI: [- 5.43, - 3.55]; $T = - 7.10$; $P < 0.001$). When stratified by amyloid status, both A β + and A β - groups exhibited significant cBF volume reductions, with a more pronounced decline observed in A β + individuals ($\beta = -5.93$; 95% CI: [- 8.21, - 3.66]; $T = - 5.15$; $P < 0.001$), whereas the A β - group demonstrated a comparatively smaller decrease ($\beta = -3.89$; 95% CI: [- 4.96, - 2.81]; $T = - 7.10$; $P < 0.001$). (Supplementary Table 1). Similarly, significant progressive cortical thinning occurred across extensive brain regions during follow-up. Regions exhibiting significant cortical thinning over the follow-

up period included the entorhinal cortex, fusiform gyrus, parahippocampal gyrus, inferior and middle temporal gyri, inferior parietal lobule, insula, isthmus cingulate, precuneus, and occipital cortex bilaterally (Supplementary Fig. 2). Cognitive performance, assessed by PACC5 scores, significantly declined over time across all participants ($\beta = -0.03$; 95% CI: $[-0.04, -0.01]$; $T = -2.90$; $P = 0.004$). When analyzed separately by amyloid status, the A β + group showed a pronounced cognitive decline ($\beta = -0.11$; 95% CI: $[-0.15, -0.06]$; $T = -4.50$; $P < 0.001$). In contrast, no significant cognitive decline was observed in the A β - group ($\beta = 0.003$; 95% CI: $[-0.01, 0.02]$; $T = 0.48$; $P = 0.635$) (Supplementary Table 2).

Longitudinal cBF Degeneration Associates with Parallel Cortical Thinning

We employed linear mixed-effects models to investigate the relationship between longitudinal cBF volume decline and cortical thinning. The models incorporated time as a fixed effect and participant as a random effect, adjusting for age, sex, and education using ComBat harmonization. To control for multiple comparisons, permutation testing with threshold-free cluster enhancement (TFCE) was applied. Significant associations were observed between cBF volume reduction and cortical thinning across various cortical areas (Fig. 2). Strong correlations were identified in bilateral temporal regions, including the inferior, middle, and superior temporal gyri, temporal poles, entorhinal cortex, and parahippocampal gyri. In the frontal lobes, notable associations were detected in the medial orbitofrontal cortex, anterior cingulate cortex, and precentral gyrus. The bilateral insular cortex also demonstrated significant thinning associated with declining cBF volume. Specifically, in the left hemisphere, significant effects appeared in the temporal pole, insula, precentral gyrus, medial orbitofrontal cortex, and cingulate cortex. In the right hemisphere, significant associations occurred primarily in the superior frontal gyrus, anterior cingulate cortex, and entorhinal cortex (Fig. 2).

Cortical Thinning as a Mediator of cBF Degeneration and Cognitive Decline

We further investigated whether cortical thinning acts as an intermediary in the association between cBF degeneration and cognitive decline in cognitively normal older adults. Cortical thinning in cBF-associated regions significantly correlated with cognitive decline. Specifically, cognitive decline was related to cortical thinning in several brain regions (Fig. 3A). In the left hemisphere, cortical thinning was observed in the temporal lobe, entorhinal cortex, parahippocampal gyrus, insular cortex, medial orbitofrontal cortex, cingulate cortex, and precentral gyrus. In the right hemisphere, it primarily affected the temporal lobe, entorhinal cortex, superior frontal gyrus, medial orbitofrontal cortex, anterior cingulate cortex, and precentral gyrus. Longitudinal cBF atrophy showed a significant correlation with declines in PACC5 scores ($r = 0.39$, $P = 1.06 \times 10^{-9}$) (Fig. 3B). Mediation analysis revealed that cortical thinning accounted for approximately 44.34% of the total effect of cBF degeneration on cognitive decline ($\beta = 0.0496$; 95% CI $[0.03, 0.07]$; $P = 1.06 \times 10^{-9}$). Both the direct effect of cBF degeneration (ADE: $\beta = 0.0276$; 95% CI $[0.01, 0.04]$; $P = 0.0016$) and the indirect effect mediated by cortical thinning (ACME: $\beta = 0.0220$; $P < 2 \times 10^{-16}$) were statistically significant (Fig. 3C-E).

Effects of Amyloid- β Pathology on the Observed Associations

Next, we examined how A β pathology influences the relationships among cBF atrophy, cortical thinning, and cognitive decline. Interestingly, longitudinal cBF atrophy did not significantly differ between A β + and amyloid-negative A β - participants (Supplementary Fig. 3). However, significant associations between cBF degeneration and cortical thinning were influenced by amyloid- β status. Specifically, regional cortical thinning associated with cBF degeneration was more pronounced in A β + participants compared to those without amyloid pathology. The regions showing these amyloid-dependent effects included the medial orbitofrontal cortex, temporal pole, entorhinal cortex, and parahippocampal cortex (Fig. 4A-B).

Effects of Amyloid- β Pathology on the Mediation of Cortical Thinning

We further investigated how A β pathology influences cortical thinning as a mediator between cBF degeneration and cognitive decline. In the A β + group, longitudinal cBF degeneration significantly correlated with declines in cognitive performance, measured by the PACC5 ($r = 0.26$, $P = 5.59 \times 10^{-4}$; Supplementary Fig. 4B). Mediation analyses revealed that cortical thinning served as a partial mediator in the association between cBF degeneration and cognitive decline (total effect: $\beta = 0.0902$; 95% CI [0.06, 0.12]; $P = 3.77 \times 10^{-16}$). Both the direct effect (ADE: $\beta = 0.0604$; $P = 0.017$) and indirect effect through cortical thinning (ACME: $\beta = 0.0299$; $P = 0.044$) were statistically significant, with cortical thinning mediating approximately 33.16% of the total effect (Supplementary Fig. 4C-E). Similarly, in the A β - group, longitudinal cBF degeneration strongly correlated with cognitive decline ($r = 0.39$, $P = 1.06 \times 10^{-9}$; Supplementary Fig. 5B). Mediation analyses indicated that cortical thinning played a mediating role in the link between cBF atrophy and cognitive impairment among A β - participants (total effect: $\beta = 0.0248$; 95% CI [0.01, 0.04]; $P = 0.0008$). Again, both direct (ADE: $\beta = 0.0148$; $P = 0.035$) and indirect effects (ACME: $\beta = 0.0100$; $P = 0.0004$) were significant, with cortical thinning accounting for approximately 40.39% of the total effect (Supplementary Fig. 5C-E).

Discussion

This study examined the longitudinal relationships among cBF degeneration, cortical thinning, and cognitive decline in cognitively normal older adults at the preclinical stage of AD. Additionally, we investigated whether A β pathology, as measured by PET imaging, modulates these associations. Our findings indicate that progressive cBF degeneration is significantly associated with cortical thinning in regions essential for cognitive function. Furthermore, cortical thinning serves as an intermediary factor linking cBF degeneration to cognitive decline. Notably, the association between cBF degeneration and cortical thinning was stronger in A β + individuals compared to A β - individuals, suggesting that A β pathology may exacerbate neurodegenerative processes linked to cholinergic system dysfunction.

In this study, we observed that even during the preclinical stage of AD—before the onset of cognitive decline—cortical thinning in regions associated with cognitive function closely corresponds to areas exhibiting cBF atrophy. This finding suggests a potential interrelationship between cortical thinning, cBF degeneration, and cognitive impairment. The loss of cholinergic neurons is a well-established neuropathological feature of AD-related dementia[27–29], and recent in vivo MRI morphometry studies have further explored the cognitive consequences of cBF degeneration in the preclinical phases of AD[30]. These studies demonstrate that reduced cBF volume, as measured by MRI, is strongly linked to cognitive impairment in AD patients. Moreover, our findings highlight the critical role of cBF atrophy in mediating the relationship between A β accumulation and early cognitive decline. Consistent with previous cross-sectional analyses, our longitudinal MRI data reveal that progressive cBF degeneration is accompanied by both global and domain-specific cognitive decline.

To further investigate this relationship, a longitudinal mediation analysis was performed to investigate the impact of cBF atrophy, cortical thinning in relevant regions, and cognitive decline. Our findings indicate that both cortical thinning and cBF atrophy are significantly associated with cognitive decline. Additionally, reduced cBF volume at baseline predicted future cognitive decline in individuals who were cognitively intact at study entry. Consistent with our results, previous MRI morphometry studies have reported associations between cortical atrophy and cognitive impairment across the AD continuum, particularly in the frontotemporal, insular, and posterior cortical regions[31, 32]. Temporal lobe atrophy, in particular, has been strongly linked to cognitive decline in AD, with studies demonstrating significant bilateral atrophy in the medial temporal lobes across various AD subtypes[33]. Recent evidence further suggests that basal forebrain degeneration not only precedes but also predicts the subsequent spread of AD pathology to cortical regions[34, 35]. Additionally, neuroimaging research has identified in vivo links between cBF degeneration and atrophy in its extensively innervated cortical projection areas[36]. Building upon these insights, our study is the first to establish a similar in vivo association between progressive cBF degeneration and simultaneous cortical thinning in preclinical AD. Furthermore, we observed that this association is linked to cognitive decline even in the earliest stages of the disease. However, similar findings have been reported in Parkinson's disease, indicating that cBF degeneration and cortical atrophy may be interrelated processes contributing to cognitive decline in neurodegenerative disorders beyond AD[37, 38]. While our results suggest that cBF degeneration influences cognitive decline primarily through its effect on cortical thinning, this does not exclude the possibility that cortical thinning independently contributes to cognitive deterioration in preclinical AD. Our supplementary analyses assessing the independent effects of cortical atrophy on cognitive decline support the notion that cortical thinning is the most immediate structural correlate of cognitive impairment. Furthermore, aspects of this process appear to be linked to the degeneration of cortically projecting cBF neurons, reinforcing its mediating role in cognitive decline.

Our findings on the cortical mediation of cognitive decline associated with cBF degeneration are consistent with similar results reported in Parkinson's disease[37, 39]. Research using selective cBF lesions in animal models has shown that these lesions lead to neuronal dysfunction in the corresponding cortical regions[40]. The severity of cognitive deficits appears to correlate with the extent

of cortical dysfunction. One plausible mechanism is that the loss of cholinergic innervation exacerbates pathological processes in affected cortical areas, potentially due to impaired clearance mechanisms. Recent studies emphasize the essential role of cholinergic innervation in maintaining metabolic and housekeeping functions in cortical neurons[41]. Furthermore, experimental AD models demonstrate that selective cBF lesions accelerate cortical A β accumulation[42, 43]. However, the directionality of these associations remains uncertain. An alternative explanation is that cortical atrophy induces retrograde degeneration of the cBF due to a lack of trophic support. Cholinergic neurons, with their extensive axonal projections to cortical and subcortical regions, rely heavily on trophic factors for maintenance and molecular transport. Disruptions in these processes may result from primary neurodegeneration in cortical target areas, further contributing to cBF atrophy.

To determine whether cBF degeneration is primarily driven by AD pathology, we conducted an explicit assessment using A β PET as a biomarker of AD pathology. Our findings indicate that cBF volume reduction correlates with cortical thinning, with more pronounced thinning observed in the A β + group compared to the A β – group. The relationship between progressive cBF degeneration and cortical thinning was partially influenced by A β status, indicating that these processes are integral to AD pathophysiology. Previous cross-sectional studies have shown that individuals with mild cognitive impairment exhibit reduced cBF volume, which is associated with cognitive deficits[30]. This reduction is strongly correlated with decreased metabolic activity across widespread cortical networks, particularly in the prefrontal cortex, medial temporal lobe, and temporoparietal junction [44]. The loss of cholinergic fibers is believed to contribute to cortical thinning in these regions by impairing synaptic plasticity and neuronal communication, thereby accelerating cognitive decline[45]. Additionally, disruptions in acetylcholine's role in regulating the blood-brain barrier may lead to abnormal metabolite transport between interstitial and cerebrospinal fluids. This dysfunction could impair A β clearance, exacerbating AD pathology by promoting A β accumulation, neurodegeneration, cBF atrophy, and cortical thinning[46]. These findings suggest that cholinergic deficits in the basal forebrain disrupt both neuronal function and cortical structure, potentially explaining the link between cBF volume loss and cortical thinning observed in our study.

This study has several limitations that warrant consideration. First, while we investigated the mediating role of cortical thinning in the relationship between cBF atrophy and cognitive decline in AD, our findings do not establish causality. The precise pathological mechanisms linking these factors remain uncertain. Second, the mediating effect of cortical thinning was relatively modest, suggesting that additional factors contribute to the relationship between cBF degeneration and cognitive function. Future research should explore the role of neuroinflammation, synaptic dysfunction, and other biomarkers to gain a more comprehensive understanding of this complex interplay. Finally, this study focused on individuals in the preclinical phase of AD over a relatively short follow-up period. Longer longitudinal studies are needed to determine whether these relationships evolve as individuals progress to clinically diagnosed AD. Additionally, future research should investigate whether early interventions targeting cBF atrophy, cortical thinning, or cholinergic dysfunction could help delay or prevent AD onset.

In conclusion, Our findings highlight the critical role of cholinergic system degeneration in the onset of cognitive deficits during the preclinical stage of AD. Moreover, they provide new insights into the relationship between cBF atrophy, regional cortical thinning, and A β pathology—two well-established biomarkers of cognitive decline in AD. These associations have important implications for biomarker-based disease prognosis and the stratification of participants in clinical trials. Additionally, our results suggest that cholinergic interventions may warrant re-evaluation, particularly in the predementia stages of AD, as targeting cholinergic dysfunction early in the disease process could offer therapeutic benefits.

Declarations

Conflict of interest

All authors declare no conflicts of interest.

Funding

Major Scientific Research Project of the Department of Education of Anhui Province (NO.2024AH040242). Natural Science Foundation of Heilongjiang Province (NO.LH2023H029), Outstanding Young Medical Talent Training Funding Project of the First Affiliated Hospital of Harbin Medical University (2024JQ06).

Author contributions

WCD, FY and THZ contributed to the conception and design of the study; SC, LJW, MLQ, ZQX, RB, LX, JTW, ZDS, XLT, JYG contributed to the qualification and analysis of data; SC and LJW contributed to drafting the text or preparing the figures.

Acknowledgements

Data used in the preparation of this article were obtained from the Harvard Aging Brain Study (HABS - P01AG036694; <https://habs.mgh.harvard.edu>). The HABS study was launched in 2010, funded by the National Institute on Aging. and is led by principal investigators Reisa A. Sperling MD and Keith A. Johnson MD at Massachusetts General Hospital/Harvard Medical School in Boston, MA. All neuroimaging and clinical data supporting this study's findings are available from <https://habs.mgh.harvard.edu/researchers>. The HABS investigators contributed data and/or participated in the design and implementation of the HABS; however, they were not involved in the analysis or authorship of this report. The data employed in this study originates from the HABS Data Release 2.0, obtained in May 2022 via the website <https://habs.mgh.harvard.edu>. Additionally, the manuscript has

received approval for publication from the HABS Data Committee, facilitated by the Chief Data Officer, currently Dr. Aaron P. Schultz (contact: apschultz@mgh.harvard.edu).

References

1. Dubois B, Hampel H, Feldman HH, Scheltens P, Aisen P, Andrieu S et al (2016) Preclinical Alzheimer's disease: Definition, natural history, and diagnostic criteria. *Alzheimers Dement* 12:292–323
2. Dubois B, Feldman HH, Jacova C, Cummings JL, Dekosky ST, Barberger-Gateau P et al (2010) Revising the definition of Alzheimer's disease: a new lexicon. *Lancet Neurol* 9:1118–1127
3. Jack CR Jr., Bennett DA, Blennow K, Carrillo MC, Dunn B, Haeberlein SB et al (2018) NIA-AA Research Framework: Toward a biological definition of Alzheimer's disease. *Alzheimers Dement* 14:535–562
4. Altomare D, de Wilde A, Ossenkoppele R, Pelkmans W, Bouwman F, Groot C et al (2019) Applying the ATN scheme in a memory clinic population: The ABIDE project. *Neurology* 93:e1635–e46
5. Delaère P, He Y, Fayet G, Duyckaerts C, Hauw JJ (1993) Beta A4 deposits are constant in the brain of the oldest old: an immunocytochemical study of 20 French centenarians. *Neurobiol Aging* 14:191–194
6. Rozalem Aranha M, Iulita MF, Montal V, Pegueroles J, Bejanin A, Vaqué-Alcázar L et al (2023) Basal forebrain atrophy along the Alzheimer's disease continuum in adults with Down syndrome. *Alzheimers Dement* 19:4817–4827
7. Labrador-Espinosa MA, Silva-Rodríguez J, Reina-Castillo MI, Mir P, Grothe MJ (2023) Basal Forebrain Atrophy, Cortical Thinning, and Amyloid- β Status in Parkinson's disease-Related Cognitive Decline. *Mov Disord* 38:1871–1880
8. Filippi M, Canu E, Donzuso G, Stojkovic T, Basaia S, Stankovic I et al (2020) Tracking Cortical Changes Throughout Cognitive Decline in Parkinson's Disease. *Mov Disord* 35:1987–1998
9. Kilimann I, Hausner L, Fellgiebel A, Filippi M, Würdemann TJ, Heinsen H et al (2017) Parallel Atrophy of Cortex and Basal Forebrain Cholinergic System in Mild Cognitive Impairment. *Cereb Cortex* 27:1841–1848
10. Labrador-Espinosa MA, Silva-Rodríguez J, Okkels N, Muñoz-Delgado L, Horsager J, Castro-Labrador S et al (2024) Cortical hypometabolism in Parkinson's disease is linked to cholinergic basal forebrain atrophy. *Mol Psychiatry*
11. Xia Y, Maruff P, Doré V, Bourgeat P, Laws SM, Fowler C et al (2023) Longitudinal trajectories of basal forebrain volume in normal aging and Alzheimer's disease. *Neurobiol Aging* 132:120–130
12. Xia Y, Dore V, Fripp J, Bourgeat P, Laws SM, Fowler CJ et al (2024) Association of Basal Forebrain Atrophy With Cognitive Decline in Early Alzheimer Disease. *Neurology* 103:e209626
13. Li R, Miao X, Han B, Li J (2023) Cortical thickness of the left parahippocampal cortex links central hearing and cognitive performance in aging. *Ann N Y Acad Sci* 1522:117–125

14. Pereira JB, Svenningsson P, Weintraub D, Brønnick K, Lebedev A, Westman E et al (2014) Initial cognitive decline is associated with cortical thinning in early Parkinson disease. *Neurology* 82:2017–2025
15. Edmonds EC, Weigand AJ, Hatton SN, Marshall AJ, Thomas KR, Ayala DA et al (2020) Patterns of longitudinal cortical atrophy over 3 years in empirically derived MCI subtypes. *Neurology* 94:e2532–e44
16. Dagley A, LaPoint M, Huijbers W, Hedden T, McLaren DG, Chatwal JP et al (2017) Harvard Aging Brain Study: Dataset and accessibility. *NeuroImage* 144:255–258
17. Papp KV, Rentz DM, Orlovsky I, Sperling RA, Mormino EC (2017) Optimizing the preclinical Alzheimer's cognitive composite with semantic processing: The PACC5. 3:668–677 *Alzheimers Dement (N Y)*
18. Dagley A, Lapoint M, Huijbers W, Hedden T, McLaren DG, Chatwal JP et al (2017) Harvard Aging Brain Study: Dataset and accessibility. *NeuroImage* 144:255–258
19. Greve DN, Billot B, Cordero D, Hoopes A, Hoffmann M, Dalca AV et al (2021) A deep learning toolbox for automatic segmentation of subcortical limbic structures from MRI images. *NeuroImage* 244:118610
20. Fischl B, Dale AM (2000) Measuring the thickness of the human cerebral cortex from magnetic resonance images. *Proc Natl Acad Sci U S A* 97:11050–11055
21. Fischl B (2012) FreeSurfer *Neuroimage* 62:774–781
22. Reuter M, Fischl B (2011) Avoiding asymmetry-induced bias in longitudinal image processing. *NeuroImage* 57:19–21
23. Reuter M, Schmansky NJ, Rosas HD, Fischl B (2012) Within-subject template estimation for unbiased longitudinal image analysis. *NeuroImage* 61:1402–1418
24. Voevodskaya O, Simmons A, Nordenskjöld R, Kullberg J, Ahlström H, Lind L et al (2014) The effects of intracranial volume adjustment approaches on multiple regional MRI volumes in healthy aging and Alzheimer's disease. *Front Aging Neurosci* 6:264
25. Mormino EC, Betensky RA, Hedden T, Schultz AP, Amariglio RE, Rentz DM et al (2014) Synergistic effect of β -amyloid and neurodegeneration on cognitive decline in clinically normal individuals. *JAMA Neurol* 71:1379–1385
26. Fortin J-P, Parker D, Tunç B, Watanabe T, Elliott MA, Ruparel K et al (2017) Harmonization of multi-site diffusion tensor imaging data. *NeuroImage* 161:149–170
27. Schumacher J, Ray NJ, Hamilton CA, Bergamino M, Donaghy PC, Firbank M et al (2023) Free water imaging of the cholinergic system in dementia with Lewy bodies and Alzheimer's disease. *Alzheimer's Dement* 19:4549–4563
28. Yoo HS, Jeon S, Cavedo E, Ko M, Yun M, Lee PH et al (2022) Association of β -Amyloid and Basal Forebrain With Cortical Thickness and Cognition in Alzheimer and Lewy Body Disease Spectra. *Neurology* 98:e947–e57

29. Yoo HS, Kim H-K, Lee J-H, Chun J-H, Lee HS, Grothe MJ et al (2024) Association of Basal Forebrain Volume with Amyloid, Tau, and Cognition in Alzheimer's Disease. *J Alzheimers Dis* 99:145–159
30. Xia Y, Dore V, Fripp J, Bourgeat P, Laws SM, Fowler CJ et al (2024) Association of Basal Forebrain Atrophy With Cognitive Decline in Early Alzheimer Disease. *Neurology*. ;103
31. Montal V, Vilaplana E, Alcolea D, Pegueroles J, Pasternak O, González-Ortiz S et al (2018) Cortical microstructural changes along the Alzheimer's disease continuum. *Alzheimer's Dement* 14:340–351
32. Sun P, He Z, Li A, Yang J, Zhu Y, Cai Y et al (2024) Spatial and temporal patterns of cortical mean diffusivity in Alzheimer's disease and suspected non-Alzheimer's disease pathophysiology. *Alzheimer's & Dementia*
33. Crane PK, Groot C, Ossenkoppele R, Mukherjee S, Choi SE, Lee M et al (2024) Cognitively defined Alzheimer's dementia subgroups have distinct atrophy patterns. *Alzheimers Dement* 20:1739–1752
34. Schmitz TW, Nathan Spreng R (2016) Basal forebrain degeneration precedes and predicts the cortical spread of Alzheimer's pathology. *Nat Commun* 7:13249
35. Fernández-Cabello S, Kronbichler M, Van Dijk KRA, Goodman JA, Spreng RN, Schmitz TW (2020) Basal forebrain volume reliably predicts the cortical spread of Alzheimer's degeneration. *Brain* 143:993–1009
36. Schmitz TW, Mur M, Aghourian M, Bedard M-A, Spreng RN (2018) Longitudinal Alzheimer's Degeneration Reflects the Spatial Topography of Cholinergic Basal Forebrain Projections. *Cell Rep* 24:38–46
37. Labrador-Espinosa MA, Silva-Rodríguez J, Reina-Castillo MI, Mir P, Grothe MJ (2023) Basal Forebrain Atrophy, Cortical Thinning, and Amyloid- β Status in Parkinson's disease-Related Cognitive Decline. *Mov Disord* 38:1871–1880
38. Crowley SJ, Kanel P, Roytman S, Bohnen NI, Hampstead BM (2024) Basal forebrain integrity, cholinergic innervation and cognition in idiopathic Parkinson's disease. *Brain* 147:1799–1808
39. Labrador-Espinosa MA, Silva-Rodriguez J, Okkels N, Muñoz-Delgado L, Horsager J, Castro-Labrador S et al (2024) Cortical hypometabolism in Parkinson's disease is linked to cholinergic basal forebrain atrophy. *Mol Psychiatry*
40. Browne SE, Lin L, Mattsson A, Georgievska B, Isacson O (2001) Selective Antibody-Induced Cholinergic Cell and Synapse Loss Produce Sustained Hippocampal and Cortical Hypometabolism with Correlated Cognitive Deficits. *Exp Neurol* 170:36–47
41. Ovsepian SV, O'Leary VB, Zaborszky L (2016) Cholinergic Mechanisms in the Cerebral Cortex: Beyond Synaptic Transmission. *Neuroscientist* 22:238–251
42. Duan L, Bhattacharyya BJ, Belmadani A, Pan L, Miller RJ, Kessler JA (2014) Stem cell derived basal forebrain cholinergic neurons from Alzheimer's disease patients are more susceptible to cell death. *Mol Neurodegener* 9:3
43. Baker-Nigh A, Vahedi S, Davis EG, Weintraub S, Bigio EH, Klein WL et al (2015) Neuronal amyloid- β accumulation within cholinergic basal forebrain in ageing and Alzheimer's disease. *Brain* 138:1722–1737

44. Grothe MJ, Heinsen H, Amaro E Jr., Grinberg LT, Teipel SJ (2016) Cognitive Correlates of Basal Forebrain Atrophy and Associated Cortical Hypometabolism in Mild Cognitive Impairment. *Cereb Cortex* 26:2411–2426
45. Sassin I, Schultz C, Thal DR, Rüb U, Arai K, Braak E et al (2000) Evolution of Alzheimer's disease-related cytoskeletal changes in the basal nucleus of Meynert. *Acta Neuropathol* 100:259–269
46. Hunter JM, Kwan J, Malek-Ahmadi M, Maarouf CL, Kokjohn TA, Belden C et al (2012) Morphological and pathological evolution of the brain microcirculation in aging and Alzheimer's disease. *PLoS ONE* 7:e36893

Tables

Table 1 is available in the Supplementary Files section.

Figures




Figure 1

Study design flowchart outlining the complete analytical approach (details in METHODS).




Figure 2

Association between longitudinal cBF volume atrophy and cortical thinning in all participants. The figure illustrates the regional effects of a regression analysis examining the relationship between the rate of change in cBF volume and the rate of change in vertex cortical thickness within a normal aging population. After adjusting for age, sex, and years of education, and applying permutation test correction, significant associations between cBF atrophy and cortical thinning were identified. Only clusters that met family-wise error correction at $P < 0.05$ are shown.




Figure 3

Mediation analysis of associations between cholinergic basal forebrain (cBF) degeneration, cortical thinning, and cognitive decline. **A.** Cortical clusters showing significant correlations between vertex-wise slopes of cortical thickness change and the slope of change in Preclinical Alzheimer Cognitive Composite-5 (PACC5) score. Vertex-wise regression analyses were restricted to cortical areas showing significant associations with cBF degeneration (as shown in Fig. 2) and corrected for multiple

comparisons using permutation testing. Only clusters that met family-wise error correction at $P < 0.05$ are shown. **B.** Individual slopes of change in PACC5 score (x-axis) are plotted against individual slopes of change in cBF volume (y-axis). The blue line represents the linear trend. **C.** Average individual slopes of thickness change in cognition-related cortical areas (x-axis) are plotted against individual slopes of change in PACC5 score (y-axis). The purple line represents the linear trend. **D.** Individual slopes of change in cBF volume (x-axis) are plotted against average individual slopes of thickness change in cognition-related cortical areas (y-axis). The yellow line represents the linear trend. **E.** Path diagram of the causal mediation model, where the effect of cBF degeneration on cognitive decline is plotted in path Z (yellow arrow), the effect of cBF degeneration on cortical thinning is plotted in path X (blue arrow), and the effect of cortical thinning on cognitive decline is plotted in path Y (purple arrow). Betas [95% confidence intervals] and P values of regression analyses are indicated for each association. The total effect of path Z is composed of the sum of the average direct effect (ZADE) and the average causal mediation effect (ZACME) (purple arrows).




Figure 4

Associations between longitudinal cholinergic basal forebrain (cBF) degeneration and regional cortical thinning in A β ⁺ and A β ⁻ groups. The figure shows regional effects of regression analyses of the slope of change in cBF volume on vertex-wise slopes of change in cortical thickness, separated by A β ⁺ and A β ⁻ groups. After adjusting for age, sex, and years of education, and applying permutation test correction, significant associations between cBF atrophy and cortical thinning were identified. Only clusters that met family-wise error correction at $P < 0.05$ are shown.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryTable2.docx](#)
- [SupplementaryFig.3.pdf](#)
- [SupplementaryTable1.docx](#)
- [SupplementaryFig.4.pdf](#)
- [SupplementaryFig.2.pdf](#)
- [SupplementaryFig.1.pdf](#)
- [SupplementaryFig.5.pdf](#)
- [Table.docx](#)
- [SUPPLEMENTALMATERIAL.docx](#)