

Characteristics (n = 100)	Count (Percent)
Age (years)	61 median (21-90 range)
Female	37 (37%)
Male	63 (63%)
Race	
White	67 (67%)
Asian	13 (13%)
Black	3 (3%)
Other/Unknown	17 (17%)
Education	
Less than high school	4 (4%)
High school graduate	26 (26%)
Associate degree	7 (7%)
Bachelor's degree	37 (37%)
Master's degree	14 (14%)
Doctorate degree	12 (12%)
Prior diagnosis of mood disorder	29 (29%)
Anxiety	25 (25%)
Depression	17 (17%)
PTSD, bipolar disorder, or other	7 (7%)
History of psychedelic use	25 (25%)
Psilocybin	19 (19%)
LSD	15 (15%)
MDMA	8 (8%)
Ketamine	3 (3%)
DMT/ayahuasca	2 (2%)
Other	9 (9%)
Primary Cancer	
Prostate	23 (23%)
Breast	16 (16%)
Head and neck	15 (15%)
Skin	11 (11%)
CNS	8 (8%)
Gynecologic	8 (8%)
Anorectal	7 (7%)
Lung	5 (5%)
Sarcoma	4 (4%)
Lymphoma	2 (2%)
Bladder	1 (1%)
Survey Timing	
Before radiation	48 (48%)
During radiation	24 (24%)
After radiation	28 (28%)
Radiation Therapy	
Definitive Intent	76 (76%)
Palliative Intent	24 (24%)