

The impact of climate change on the productivity of conservation agriculture

Yang Su (✉ yang.su@inra.fr)

French National Institute for Agricultural Research (INRAE) <https://orcid.org/0000-0002-4717-9971>

Benoît Gabrielle

AgroParisTech - INRA <https://orcid.org/0000-0002-9131-2549>

David Makowski

University of Paris-Saclay <https://orcid.org/0000-0001-6385-3703>

Article

Keywords: climate change, conservation agriculture, conventional tillage

Posted Date: September 15th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-63752/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Nature Climate Change on June 17th, 2021.
See the published version at <https://doi.org/10.1038/s41558-021-01075-w>.

The impact of climate change on the productivity of conservation agriculture

Authors

Yang Su^a, Benoit Gabrielle^a, David Makowski^{b,c}

Affiliations

^a UMR ECOSYS, INRAE AgroParisTech, Université Paris-Saclay, 78850 Thiverval-Grignon, France

^b UMR MIA 518 (applied mathematics and computer science), INRAE AgroParisTech, Université Paris-Saclay, 75231 Paris, France

^c CIRED, Centre international de recherche sur l'environnement et le développement, 45bis Avenue de la Belle Gabrielle, 94130 Nogent-sur-Marne, France

Corresponding Author: Yang Su (yang.su@inrae.fr)

Abstract

Conservation agriculture (CA) is being promoted as a set of management practices that can sustain crop production while providing positive environmental externalities. However, its impact on crop productivity is still hotly debated, and how this productivity will be affected by climate change remains uncertain. Here we compared the productivity of CA vs. conventional tillage (CT) systems under current and future climate conditions using a probabilistic machine-learning approach at the global scale. We reveal large differences in the probability of yield gains with CA across crop types, climate zones, and geographical regions. We show that, for most crops, CA performed better in continental, arid and temperate regions than in tropical ones. Under future climate conditions, the relative productive performance of CA is expected to increase for maize in almost all cropping areas within the tropical band, thus improving the competitiveness of CA for this major crop.

Conservation agriculture is a crop production system based on three principles: minimum soil disturbance (going as far as no tillage or NT), permanent soil cover with crop residue, and diversified crop rotation¹. In compliance with sustainability goals, CA is designed to ensure the long-term crop production while improving crop resilience to climate change and protecting the environment. It has been shown to enhance soil carbon sequestration, improve soil quality, reduce soil erosion, and increase biodiversity^{2,3}. However, since the productivity of crops depends on many interacting factors including local climatic conditions⁴, soil characteristics⁵, and agricultural management practices⁶, it is difficult to assess the potential of CA to increase agricultural productivity. Based on recent meta-analyses⁵⁻⁷ synthesizing current evidence on the effect of CA on crop yields, CA is expected to lead to a yield reduction⁶ compared to CT. However, the heterogeneity of the experimental results on CA vs. CT is very large, and their outcome varies as a function of climate conditions⁶ and management practices⁶⁻⁸. Most of the above studies relied on a synthetic aridity index or a large climate zone to characterize the impact of climate, which makes it hard to analyze the response of CA productivity to inter-annual weather variability or to predict the impact of future changes in climate. To date, a comprehensive, global synthesis of the impact of climate change on the productive performance of CA with respect to CT systems is still lacking.

In this paper, we compared the crop yields of CA and CT systems under current and future climate conditions based on a new, global database of paired yield observations of NT (including CA and NT without crop rotation and/or residue retention) vs. CT. We compiled the datasets collected from the previous meta-analyses⁵⁻⁷, and supplemented them by the latest literature references on this topic and the inclusion of a broader set of climatic variables from external databases, such as precipitation (P)⁹, minimum air temperature (Tmin)¹⁰, average air temperature (Tave)⁹, maximum air temperature (Tmax)¹⁰ and potential evapotranspiration (E)^{11,12} over the crop growing seasons¹³ considered in each experiment. As an indicator of water availability for crops, the precipitation balance (PB) was defined as (P – E) over the growing season⁵.

A machine learning model based on random forest¹⁴ was trained and cross-validated based on 4071 paired yield data of NT (including CA and NT without crop rotation and/or residue retention) vs. CT to map the probability of yield gain from CA (NT with crop rotation and residue retention), i.e. $\left(\frac{Yield_{CA}}{Yield_{CT}} > 1\right)$, at a spatial resolution of $0.5^\circ \text{ latitude} \times 0.5^\circ \text{ longitude}$ based on current (2011-2020) and future (2051-2060) average climatic conditions for eight major crops (barley, cotton, maize, rice, sorghum, soybean, sunflower, and wheat). Crop types, soil types and management practices were kept unchanged in all climate scenarios. The maps were then used to calculate the accumulated fractions of the cropping area as a function of the yield gain probability of CA vs. CT, and as a function of the level of change on the probability of yield gain from CA vs. CT in future climatic condition. These fractions show the proportions of cropping area with low to high probabilities of yield gain from CA vs. CT, and with low to high change level on the probability of yield gain from CA vs. CT under climate change. They were both computed at the global scale and in different climate zones¹⁵. The results were used to identify the favorable and unfavorable climate zones for CA, and to assess the climate change impact on the productivity of CA in different climate zones. The details of model setting for global projection are explained in the **Methods** section and further detailed in S1.

Our results show that, under current climate condition, CA is associated with a high chance of yield loss under tropical climates (Figure 1) and that, for most of the studied crops, the overall probability of yield gain is higher in continental, arid, and temperate regions than in tropical regions (Figure 1a, Figure 1b, Figure 1d-g). The risk of yield loss with CA is particularly high for rice, with a probability of yield gain with CA under 50% in about 80% of the global rice cropping area. This fraction rises to about 100% in the tropics (Figure 1d).

For several crops and climate regions, the estimated effect of climate change on the probability of yield gain with CA is relatively moderate. In approximately half of the cropping areas, a decrease of up to 10% in this probability is expected, while in the other half an increase of up to 10% may be anticipated (Figure 2). However, in some important cases the effect of climate change is stronger, especially for maize in tropical regions. Here, the probability of yield gain with CA increases in about 72% of the cropping area, with a larger than 10% gain for 30% of the maize cropping area (Figure 2c). An increase in yield gain is also estimated for more than 50% of the cropping area for rice and sorghum in tropical regions (Figure 2d, Figure 2f) and for sorghum in continental regions (Figure 2e).

To gain further insight into the conditions suitable for CA, partial dependence plots¹⁶ - a common approach used in machine learning to visualize the marginal effect of one or two inputs on the predicted outcome¹⁷ - were generated. The probability of yield gain with CA increases with decreasing PB, while the effect of temperature appears more complex (S2) since the relationship among yield gain, PB and Tave is not linear (Figure 3). Thus, the variation of CA performance in the future will depend on the current level of PB and Tave and the direction and magnitude of the change in these two drivers. The 2D partial dependence plot of Figure 3 can help project changes in the performance of CA for a given location or region based on their current levels of PB and Tave and their anticipated variations. For example for maize, the increased temperature and relatively stable PB anticipated under climate change

scenario RCP 4.5 in tropical regions (depicted by the continuous and dashed boxes in Figure 3) would lead to an increased probability of yield gain with CA, which is consistent with the results of Figure 2c.

Probabilities of yield gain with CA show important geographical variations under both current and future climate conditions, in particular for maize (Figure 4) but also for other crop species (S3a-g). Yield gains with CA are more likely in relatively higher latitude regions (> 40 Deg.) than lower latitudinal bands for barley, cotton, rice, sorghum, soybean, sunflower, and wheat (S3), in line with the results showed on Figure 1. For maize, the probability of yield gain from CA is less than 0.4 in Laurentian Plateau in Canada, northcentral and northeast of the US, Brazilian Highlands, north part of Western Europe and east part of Eastern Europe, Central Asia and northeast of China. For this crop, the probability of yield gain is higher than 0.5 in Arid West of the US, most of the regions in Mato Grosso and part of Amazon Basin in Brazil, the southern part of west Africa, north of India, and North China Plain (Figure 4). For other crop species, CA has a higher chance to lead to a yield loss compared to CT in the tropical regions, south of China (S3a-g), and north of India (S3a-f). For rice, sorghum, sunflower, wheat cropped in Atlantic and Gulf Coastal Plain in the US, CA is also likely to lead to a yield loss (S3c, S3d, S3f, S3g). Finally, CA has higher chance to outperform CT in several major cropping areas of barley, sunflower and wheat (S3a, S3f, S3g; S5a, S5g, S5h).

The maps reporting the differences of yield gain probabilities between current and future climate conditions (Figure 5 and S4) reveal important geographical disparities in the effects of climate change on the chances of yield gain with CA. Under climate change scenario RCP 4.5, the probability of yield gain is expected to increase in most of the northcentral and northeast of the US for barley, maize, sorghum, soybean, sunflower, and wheat (S4a, S4d-g, Figure 5); In most of the central-west region and Amazon Basin region in Brazil, Western Africa, and South Pacific Asia for maize, rice, and soybean (S4c, S4e, Figure 5); In India for cotton, maize, rice, sorghum, soybean, sunflower (S4b-f, Figure 5); In most of the north part of Central and Eastern Europe for barley, maize, sorghum, soybean, sunflower and wheat (S4a, S4d-g, Figure 5); In northeast China for rice, sorghum (S4c, S4d). While the overall performance of CA will decrease in the future in the US between 30- to 40-degree latitude for the barley, maize, cotton, rice, sorghum, soybean, and sunflower (S4a-f, Figure5); In most temperate regions in South America, including Uruguay, south of Brazil, and north of Argentina for barley, cotton, maize, rice, sorghum, and sunflower (S4a-d, S4f, Figure 5); In the southern part of east Europe and northwest Asia for Barley, soybean, sunflower and wheat (S4a, S4e-g); In south China for cotton, rice, sorghum, sunflower (S4b-d, S4f).

To assess the robustness of our conclusions, we analyzed the sensitivity of our results to four different climate models and RCP scenarios (S6). To summarize the response of our model to these choices, we plotted the fractions of global cropping areas corresponding to increasing levels of yield gain probability (from -0.1 to 0.2). The choice of the climate models had very little impact on the results (S6i-p), although the results obtained with Hadgem2-es, and Ipsl-cm5a-lr are somewhat more extreme than those obtained with Gfdl-esm2m and Miroc5. The sensitivity to the climate change scenarios was more important (S6a-h). Although the main conclusions remain similar across all RCP scenarios, the stronger changes of probability of yield gain are obtained under RCP 8.5 compared to RCP 6.0, RCP 4.5, and RCP 2.6 (S6a-h), in particular the changes of yield gains become higher for maize and rice under RCP 8.5.

Discussion

Compared to previous studies on the productivity of CA^{5-7,18}, this is the first time that the probabilities of yield gain with CA systems have been mapped in current and future climate scenarios. Thus, our results offer meaningful and new information for policymakers, agricultural extension services, and farmers. Relying on a global experimental dataset, we were able to identify favorable and unfavorable climatic conditions and geographical regions for the implementation of CA for eight major staple crops under current and future climate conditions. Some of the most promising geographical regions in our

analysis had also been identified in previous studies¹⁸, but we were able to report information on yield gains as probabilities instead of the simple categories of yield increase or decrease. More importantly, none of the previous meta-analyses addressed the effect of future climate change scenarios on the performance of CA and its geographical patterns. For most crops, CA outperformed CT in continental, arid and temperate regions, but proved less suitable in tropical regions. This overall pattern is in line with previous work⁶. We also show that there is higher chance of yield gain from CA under dryer conditions, compared to wetter environments^{5-7,18}. This is likely due to CA improving soil aggregate stability via crop residue returns to soils, which reduces soil evaporation and surface runoff, and maintains a higher level of soil moisture content compared to CT¹⁹⁻²⁷. This competitive advantage of CA comes into play for dryer climate conditions²⁸⁻³³, but does not apply in wetter and cooler environment (S2a-c, S2f-i) where the soil evaporation is less intense. In tropical regions, CA is commonly associated with waterlogging especially with poorly drained soils^{5,6,27} which can result in substantial yield losses. Air temperature also influences the performance of CA, but its effect appears complex variable across crop species. Previous studies reported that permanent soil cover or residue retention tends to lower soil temperature under warm conditions, but this trend can be reversed under cold conditions because of soil cover acting as an insulation barrier^{22,34,35}. As a result, CA may reduce the negative impact of extreme air temperature events on crop yields, but this effect can be either positive or negative depending on crop types, cultivars and development stages. Lower soil temperatures may delay crop establishment and growth^{27,36,37} and have a negative effect on crop yield³⁸⁻⁴⁰, but lower soil temperature has a positive impact on crop yield under arid conditions^{36,37}.

CA systems can dramatically improve biodiversity, increase the soil organic matter, and bring a lot of positive environmental externalities such as reduced soil erosion, improved soil quality and enhanced carbon sequestration^{2,3,41}. Moreover, CA can provide a buffer effect to mitigate the impacts from more intense rainstorms, more frequent droughts and increased daily temperature ranges, which can significantly increase the crop resilience towards the changing climate and increase the stability of crop yields^{42,43}. Although CA is associated with a high probability of yield loss in many regions, we also showed that, under future climate conditions, the relative productive performance of CA is expected to increase for several crop species. This is especially true for maize in tropical regions, which further strengthens the competitiveness of CA for this staple crop. Our results thus support the idea that CA will be a relevant option for cropping systems in the future, capable of ensuring a long-term, sustainable agricultural production for some key cropping areas^{44,45}.

Methods

Data collection

The literature search was done in February 2020 using the following keywords ‘Conservation agriculture / No-till / No tillage / Zero tillage’ & ‘Yield / Yield change’ in the websites ‘ScienceDirect’ and ‘Science Citation Index (web of science)’. A total of 1012 potentially relevant papers were identified by reviewing the title and abstract, these papers were then screened according to the procedure summarized in S7. In the end, 422 papers left (published between 1983 to 2020), and 4071 paired yield observations from NT (including CA and NT without crop rotation or/and residue retention) and CT were collected from those papers, including 8 major crop species (369 observations for barley, 94 observations for cotton, 1580 observations for maize, 169 observation for rice, 145 observations for sorghum, 552 observations for soybean, 60 observations for sunflower, 1122 observations for wheat) in 51 countries (S8) from the experiment year 1980 to 2017. We also retrieved from the papers the information of crop type, year and location of the experiments, and agricultural management activities including crop irrigation (crop irrigated vs. rainfed), the application of fertilizers (field fertilized vs. unfertilized), the management of weed and pest (controlled vs. non-controlled), crop rotation (rotated vs. monoculture) and the management of crop residues (retained vs. removed). Additional data were extracted from several external databases, pertaining to crop growing season¹³, soil texture⁴⁶ and climate factors such as precipitation (P)⁹, minimum temperature (Tmin)¹⁰, average temperature (Tave)⁹, maximum temperature (Tmax)¹⁰ and potential evapotranspiration (E)^{11,12} in the growing season¹³ in a particular year, and the precipitation balance (PB) was defined as precipitation minus total evapotranspiration, which indicated the water availability in the growing season. These data were obtained at a spatial resolution of $0.5^{\circ} \text{latitude} \times 0.5^{\circ} \text{longitude}$, and if the source data were in a finer spatial resolution, they were downscaled to the resolution of $0.5^{\circ} \text{latitude} \times 0.5^{\circ} \text{longitude}$.

Model training and cross-validation

The machine learning algorithm random forest¹⁴ was trained to analyze the yield ratios of NT (including CA and NT without crop rotation or/and residue retention) vs. CT as the function of climatic variables, crop types, soil textures, and agricultural management activities. The climatic variables during the growing season such as PB, Tmin/Tave/Tmax were defined as numerical inputs, while crop type, soil texture, and agricultural management activities including crop irrigation, field fertilization, control of pests and weeds, crop rotation and crop residue management were defined as categorical inputs. The model output was expressed as the probability of yield gain from NT (including CA and NT without crop rotation or/and residue retention) vs. CT. The performance of the algorithm was assessed by estimating the area under the ROC curve by leave-one-out cross validation (LOOCV) (AUC=0.782, see S9). Since the inputs related to crop rotation and crop residue management were included in the model, it was possible to map the probability of yield gain for CA systems when combining NT, crop rotation and crop residue management. Maps were generated for all crop species at a spatial resolution of $0.5^{\circ} \text{latitude} \times 0.5^{\circ} \text{longitude}$.

Importance and partial dependence plots

Precipitation balance and average temperature appeared to be the most influential input parameters in the importance ranking obtained from our random forest algorithm (S10). We thus drew 2D partial dependence plots relating the probability of yield gain with CA (NT with crop rotation and residue retention) to PB and Tave, on a crop-by-crop basis. The detailed procedure implemented to produce 2D partial dependence plots is described in the flow chart reported in S11.

Global projection

The fitted random forest model was used to estimate the probability of yield gain from CA (NT with crop rotation and residue retention) for each grid cell located in cropping regions under current (2011-

2020) and future (2051-2060) climate scenarios. The monthly-average values of the climatic variables (PB, Tmin/Tave/Tmax) were calculated in each grid cell over the two time periods considered, and then these data were used to calculate the climatic variables during the growing season based on the crop calendar database¹³ (assume no change in current and future scenario). All the climatic data in both current and future scenarios were obtained from four climate models: Gfdl-esm2m, Hadgem2-es, Ipsl-cm5a-lr, and Miroc5, and 4 RCP scenarios: RCP 2.6, RCP 4.5, RCP 6.0, and RCP 8.5. This results in 32 combinations (4 climate models x 4 RCP scenarios x 2 periods). We mainly focused on the Ipsl-cm5a-lr model and RCP 4.5 scenario in the baseline simulations, because of their importance and similar role in the protocol of ISIMIP2b project⁴⁷. However, results from all combinations were analyzed and shown in S6. All the climatic data can be downloaded through the website of Lawrence Livermore National Laboratory⁴⁸.


We did not change the categorical inputs describing cropping practices between current and future scenarios. The global soil texture was set based on HWSD dataset⁴⁶. To estimate the performance of CA vs. CT, we set the agricultural management activities as crop rotated, crop residue retained. As for other agricultural management practices, the field was set to be fertilized, and with the control of pest and weed. As for crop irrigation, it was set based on the crop irrigation mask from MIRCA2000 dataset⁴⁹. When more than 50% of the area in a grid cell were under rainfed conditions for a given crop in the MIRCA2000 database, then this cell was considered as non-irrigated for this crop, and vice versa. See S1 for the details of model settings. The model outputs were mapped at a spatial resolution of $0.5^{\circ} \text{ latitude} \times 0.5^{\circ} \text{ longitude}$ based on the MIRCA2000 crop mask database⁴⁹. Accumulated area fractions under different levels of yield gain probability and different levels of probability change between current and future scenarios were computed at the global scale and in different climate regions.

Climate regions

The “global” indicated the global cropping region for each crop²⁵. According to the Köppen-Geiger classification¹⁵ and its nomenclature. The “tropical climate” included the regions with the climate types Af, Am, As, Aw¹⁵. The “arid climate” included the regions with the climate types Bwk, Bwh, Bsk, Bsh¹⁵. The “temperate climate” included the regions with the climate types Cfa, Cfb, Cfc, Csa, Csb, Csc, Cwa, Cwb, Cwc¹⁵. The “continental climate” included the regions with the climate types Dfa, Dfb, Dfc, Dfd, Dsa, Dsb, Dsc, Dsd, Dwa, Dw b, Dwc, Dwd¹⁵.



Figures

Figure 1




Accumulated fraction of the cropping area as a function of the probability of yield gain from CA (with field fertilization, and the control of weed and pest) for eight major crops (a-h) and different climate zones. The results are based on the average climate conditions over 2021-2020 according to the *Ipsl-cm5a-lr* climate model and RCP 4.5 scenario. The cropping areas in the main climate zones are given in (i).


251 Figure 2




254 Accumulated fractions of the cropping area for different levels of change in the probability of yield gain from CA (with field
255 fertilization, and the control of weed and pest). The results are based on the mean climate conditions in 2021-2020 for current
256 scenario and 2051-2060 for future scenario (Ipsl-cm5a-lr climate model and RCP 4.5 scenario).



258
259 2D partial dependence plot for maize. The green color indicates that the probability of yield gain from CA is higher than 0.5,
260 while the red means that the probability is lower than or equal to 0.5. The ticks on x-axis and y-axis are the observations of
261 PB and Tave, a more reliable response can be expected in zones with high numbers of observations. The box plots show the
262 distribution of PB and Tave in maize cropping areas with tropical climate. "F." means future, "C." means current, "tr."
263 means tropical. The box with continuous lines in the partial dependence plot shows the 1st and 3rd quartile of the range of PB
264 and Tave under current climate conditions in the maize tropical area, while the box with dash lines shows the 1st and 3rd
265 quartile of the range of PB and Tave under future climate conditions in the maize tropical area.



268 Probability of yield gain with CA for maize under current (a) and future (b) climate conditions. Those regions with the
269 probability of yield gain lower than 0.5 were highlighted in red (and green shades when the probability was higher). The
270 results are based on the mean climate condition in 2011-2020 (current) and 2051-2060 (future) from the Ipsi-cm5a-lr climate
271 model and the RCP 4.5 scenario. The side plots showed the variations of 1st quartile (red), median (blue), and 3rd quartile
272 (green) yield gain probability with latitude and longitude. Non-cropping region indicated both the regions without maize and
273 the regions where climate data was unavailable.



275

276 The change of probability of yield gain with CA for maize in future climate conditions, regions with a decreasing trend are

277 depicted in red, while those with an increase in yield gain probability are depicted in green. The results are based on the mean

278 climate condition in 2011-2020 (current) and 2051-2060 (future) from Isp1-cm5a-lr climate model and RCP 4.5 scenario. The

279 side plots showed the variations of 1st quartile (red), median (blue), and 3rd quartile (green) change on the probability of

280 yield gain with latitude and longitude. Non-cropping region indicated both the regions without maize cropped and the regions

281 with unavailable climatic data.

282 Reference

- 283 1. Bisht, J. K., Meena, V. S., Mishra, P. K. & Pattanayak, A. *Conservation Agriculture*. (Springer
284 Singapore, 2016). doi:10.1007/978-981-10-2558-7.
- 285 2. Farooq, M. & Kadambot H. M. Siddique. *Conservation Agriculture*. *CONSERVATION*
286 *AGRICULTURE* (Springer, 2015). doi:10.1007/978-3-319-11620-4.
- 287 3. Giller, K. E. *et al.* Beyond conservation agriculture. *Frontiers in Plant Science* **6**, (2015).
- 288 4. Jat, S. L. *et al.* Differential response from nitrogen sources with and without residue
289 management under conservation agriculture on crop yields, water-use and economics in
290 maize-based rotations. *Field Crops Research* **236**, 96–110 (2019).
- 291 5. Steward, P. R. *et al.* The adaptive capacity of maize-based conservation agriculture systems to
292 climate stress in tropical and subtropical environments: A meta-regression of yields.
293 *Agriculture, Ecosystems and Environment* **251**, 194–202 (2018).
- 294 6. Pittelkow, C. M. *et al.* When does no-till yield more? A global meta-analysis. *Field Crops*
295 *Research* **183**, 156–168 (2015).
- 296 7. Pittelkow, C. M. *et al.* Productivity limits and potentials of the principles of conservation
297 agriculture. *Nature* **517**, 365–368 (2015).
- 298 8. Nichols, V., Verhulst, N., Cox, R. & Govaerts, B. Weed dynamics and conservation agriculture
299 principles: A review. *Field Crops Research* **183**, 56–68 (2015).
- 300 9. NOAA/OAR/ESRL PSD. University of Delaware Air Temperature & Precipitation.
301 https://www.esrl.noaa.gov/psd/data/gridded/data.UDel_AirT_Precip.html (2018).
- 302 10. NOAA/OAR/ESRL PSD. CPC Global Daily Temperature.
303 <https://www.esrl.noaa.gov/psd/data/gridded/data.cpc.globaltemp.html> (2020).
- 304 11. Martens, B. *et al.* GLEAM v3: satellite-based land evaporation and root-zone soil moisture.
305 *Geoscientific Model Development* **10**, 1903–1925 (2017).
- 306 12. Miralles, D. G. *et al.* Global land-surface evaporation estimated from satellite-based
307 observations. *Hydrology and Earth System Sciences* **15**, 453–469 (2011).
- 308 13. University of Wisconsin-Madison. Crop Calendar Dataset: netCDF 5 min | SAGE, UW-Madison.
309 <http://www.sage.wisc.edu/download/sacks/netCDF5min.html>.
- 310 14. Ho, T. K. Random decision forests. in *Proceedings of 3rd International Conference on Document*
311 *Analysis and Recognition* vol. 1 278–282 vol.1 (1995).
- 312 15. Kottek, M., Grieser, J., Beck, C., Rudolf, B. & Rubel, F. World Map of the Köppen-Geiger climate
313 classification updated. *Meteorologische Zeitschrift* **15**, 259–263 (2006).
- 314 16. Hastie, T., Tibshirani, R. & Friedman, J. *The Elements of Statistical Learning*. (Springer New York,
315 2009). doi:10.1007/978-0-387-84858-7.
- 316 17. Friedman, J. H. Greedy Function Approximation: A Gradient Boosting Machine. *The Annals of*
317 *Statistics* **29**, 1189–1232 (2001).

- 318 18. Laborde, J. P., Wortmann, C. S., Blanco-Canqui, H., Baigorria, G. A. & Lindquist, J. L. Identifying
319 the drivers and predicting the outcome of conservation agriculture globally. *Agricultural*
320 *Systems* **177**, 102692 (2020).
- 321 19. Su, Z. *et al.* Effects of conservation tillage practices on winter wheat water-use efficiency and
322 crop yield on the Loess Plateau, China. *Agricultural Water Management* **87**, 307–314 (2007).
- 323 20. van Wie, J. B., Adam, J. C. & Ullman, J. L. Conservation tillage in dryland agriculture impacts
324 watershed hydrology. *Journal of Hydrology* **483**, 26–38 (2013).
- 325 21. Shukla, M. K., Lal, R. & Ebinger, M. Tillage effects on physical and hydrological properties of a
326 typical argiaquoll in Central Ohio. *Soil Science* **168**, 802–811 (2003).
- 327 22. Shen, Y., McLaughlin, N., Zhang, X., Xu, M. & Liang, A. Effect of tillage and crop residue on soil
328 temperature following planting for a Black soil in Northeast China. *Scientific Reports* **8**, 4500
329 (2018).
- 330 23. Zuazo, V. H. D. & Pleguezuelo, C. R. R. Soil-erosion and runoff prevention by plant covers. A
331 review. *Agronomy for Sustainable Development* **28**, 65–86 (2008).
- 332 24. Shaxson, F. & Barber, R. *Optimizing soil moisture for plant production, The significance of soil*
333 *porosity*. (FOOD AND AGRICULTURE ORGANIZATION OF THE UNITED NATIONS, 2003).
- 334 25. Derpsch, R., Sidiras, N. & Roth, C. H. Results from Studies made from 1977-1984 to Control
335 Erosion by No-Tillage Techniques in Parana, Brazil. *Soil & Tillage Research* **8**, 253–263 (1986).
- 336 26. TerAvest, D., Carpenter-Boggs, L., Thierfelder, C. & Reganold, J. P. Crop production and soil
337 water management in conservation agriculture, no-till, and conventional tillage systems in
338 Malawi. *Agriculture, Ecosystems & Environment* **212**, 285–296 (2015).
- 339 27. Page, K. L., Dang, Y. P. & Dalal, R. C. The ability of conservation agriculture to conserve soil
340 organic carbon and the subsequent impact on soil physical, chemical, and biological properties
341 and yield. *Frontiers in Sustainable Food Systems* **4**, (2020).
- 342 28. Halvorson, A. D. *et al.* Spring Wheat Response to Tillage and Nitrogen Fertilization in Rotation
343 with Sunflower and Winter Wheat. *Agronomy Journal* **92**, 136–144 (2000).
- 344 29. Radford, B. J. & Thornton, C. M. Effects of 27 years of reduced tillage practices on soil
345 properties and crop performance in the semi-arid subtropics of Australia. *International Journal*
346 *of Energy Environment and Economics* **19**, 565 (2011).
- 347 30. García-Palacios, P., Alarcón, M. R., Tenorio, J. L. & Moreno, S. S. Ecological intensification of
348 agriculture in drylands. *Journal of Arid Environments* **167**, 101–105 (2019).
- 349 31. Scopel, E. *et al.* Conservation agriculture cropping systems in temperate and tropical
350 conditions, performances and impacts. A review. *Agronomy for Sustainable Development* **33**,
351 113–130 (2013).
- 352 32. Thierfelder, C. & Wall, P. C. Investigating Conservation Agriculture (CA) Systems in Zambia and
353 Zimbabwe to Mitigate Future Effects of Climate Change. *Journal of Crop Improvement* **24**, 113–
354 121 (2010).
- 355 33. Lal, R. The Role of Residues Management in Sustainable Agricultural Systems. *Journal of*
356 *Sustainable Agriculture* **5**, 51–78 (1995).

34. Chen, S., Zhang, X., Pei, D. & Sun, H. Effects of corn straw mulching on soil temperature and soil evaporation of winter wheat field. **21**, 171–173 (2005).
35. Moreno, M. M., Moreno, A. & Mancebo, I. Comparison of different mulch materials in a tomato (*Solanum lycopersicum* L.) crop. *Spanish Journal of Agricultural Research* 454–464 (2009).
36. Muñoz-Romero, V., Lopez-Bellido, L. & Lopez-Bellido, R. J. Effect of tillage system on soil temperature in a rainfed Mediterranean Vertisol. *International Agrophysics* **29**, 467–473 (2015).
37. Hatfield, J. L. & Prueger, J. H. Temperature extremes: Effect on plant growth and development. *Weather and Climate Extremes* **10**, 4–10 (2015).
38. Rusinamhodzi, L. *et al.* A meta-analysis of long-term effects of conservation agriculture on maize grain yield under rain-fed conditions. *Agronomy for Sustainable Development* **31**, 657–673 (2011).
39. Hatfield, J. L., Sauer, T. J. & Prueger, J. H. Managing soils to achieve greater water use efficiency. *Agronomy Journal* **93**, 271–280 (2001).
40. Wang, X. B., Cai, D. X., Hoogmoed, W. B., Oenema, O. & Perdok, U. D. Developments in conservation tillage in rainfed regions of North China. *Soil and Tillage Research* **93**, 239–250 (2007).
41. Smith, Pete., Powlson, D. S., Glendining, M. J. & Smith, Jo. U. Preliminary estimates of the potential for carbon mitigation in European soils through no-till farming. *Global Change Biology* **4**, 679–685 (1998).
42. Williams, A. *et al.* A regionally-adapted implementation of conservation agriculture delivers rapid improvements to soil properties associated with crop yield stability. *Scientific Reports* **8**, 8467 (2018).
43. Kassam, A., Friedrich, T., Shaxson, F. & Pretty, J. The spread of Conservation Agriculture: justification, sustainability and uptake. *International Journal of Agricultural Sustainability* **7**, 292–320 (2009).
44. Hobbs, P. R., Sayre, K. & Gupta, R. The role of conservation agriculture in sustainable agriculture. *Philosophical Transactions of the Royal Society B: Biological Sciences* **363**, 543–555 (2008).
45. Govaerts, B. *et al.* Conservation agriculture as a sustainable option for the central Mexican highlands. *Soil and Tillage Research* **103**, 222–230 (2009).
46. University of Tokyo. Soil Texture Map. <http://hydro.iis.u-tokyo.ac.jp/~sujan/research/gswp3/soil-texture-map.html>.
47. Frieler, K. *et al.* Assessing the impacts of 1.5 °C global warming – simulation protocol of the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP2b). *Geoscientific Model Development* **10**, 4321–4345 (2017).
48. Lawrence Livermore National Laboratory. ESGF project. <https://esgf-node.llnl.gov/projects/esgf-llnl/>.

396 49. Portmann, F. T., Siebert, S. & Döll, P. MIRCA2000-Global monthly irrigated and rainfed crop
397 areas around the year 2000: A new high-resolution data set for agricultural and hydrological
398 modeling. *Global Biogeochemical Cycles* **24**, (2010).
399

Figures





Figure 1

Accumulated fraction of the cropping area as a function of the probability of yield gain from CA (with field fertilization, and the control of weed and pest) for eight major crops (a-h) and different climate zones. The results are based on the average climate conditions over 2021-2020 according to the Ipsi-cm5a-lr climate model and RCP 4.5 scenario. The cropping areas in the main climate zones are given in (i).



Accumulated fractions of the cropping area for different levels of change in the probability of yield gain from CA (with field fertilization, and the control of weed and pest). The results are based on the mean climate conditions in 2021-2020 for current scenario and 2051-2060 for future scenario (Ipsl-cm5a-lr climate model and RCP 4.5 scenario).

Figure 2

Accumulated fractions of the cropping area for different levels of change in the probability of yield gain from CA (with field fertilization, and the control of weed and pest). The results are based on the mean climate conditions in 2021-2020 for current scenario and 2051-2060 for future scenario (Ipsl-cm5a-lr climate model and RCP 4.5 scenario).




Figure 3

2D partial dependence plot for maize. The green color indicates that the probability of yield gain from CA is higher than 0.5, while the red means that the probability is lower than or equal to 0.5. The ticks on x-axis and y-axis are the observations of PB and Tave, a more reliable response can be expected in zones with high numbers of observations. The box plots show the distribution of PB and Tave in maize cropping areas with tropical climate. "F." means future, "C." means current, "tr." means tropical. The box with continuous lines in the partial dependence plot shows the 1st and 3rd quartile of the range of PB and Tave under current climate conditions in the maize tropical area, while the box with dash lines shows the 1st and 3rd quartile of the range of PB and Tave under future climate conditions in the maize tropical area.




Figure 4

Probability of yield gain with CA for maize under current (a) and future (b) climate conditions. Those regions with the probability of yield gain lower than 0.5 were highlighted in red (and green shades when the probability was higher). The results are based on the mean climate condition in 2011-2020 (current) and 2051-2060 (future) from the Ipsl-cm5a-Ir climate model and the RCP 4.5 scenario. The side plots showed the variations of 1st quartile (red), median (blue), and 3rd quartile (green) yield gain probability

with latitude and longitude. Non-cropping region indicated both the regions without maize and the regions where climate data was unavailable.




Figure 5

The change of probability of yield gain with CA for maize in future climate conditions, regions with a decreasing trend are depicted in red, while those with an increase in yield gain probability are depicted in green. The results are based on the mean climate condition in 2011-2020 (current) and 2051-2060 (future) from Ipsi-cm5a-lr climate model and RCP 4.5 scenario. The side plots showed the variations of 1st quartile (red), median (blue), and 3rd quartile (green) change on the probability of yield gain with latitude and longitude. Non-cropping region indicated both the regions without maize cropped and the regions with unavailable climatic data.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementarymaterialsTheimpactofclimatechangeontheprobabilityofconservationagriculture.pdf](#)