

Long COVID symptoms among Japanese adults by age group.

Symptoms	Period prevalence (up to 1 year), n (%)		
	Younger (n=195)	Middle-aged (n=1,186)	Older (n=216)
Fatigue	84 (43.1)	460 (38.8)	30 (25.9)
Headache	73 (37.4)	337 (28.4)	14 (12.1)
Cough	67 (34.4)	434 (36.6)	33 (28.4)
Runny nose	58 (29.7)	419 (35.3)	34 (29.3)
\Nasal congestion	55 (28.2)	298 (25.1)	15 (12.9)
Sore throat	47 (24.1)	312 (26.3)	19 (16.4)
Anxiety	33 (16.9)	143 (12.1)	4 (3.4)
Dizziness	32 (16.4)	144 (12.1)	10 (8.6)
Depression	28 (14.4)	104 (8.8)	3 (2.6)
Diarrhoea	26 (13.3)	169 (14.2)	6 (5.2)
Blurry vision	24 (12.3)	208 (17.5)	18 (15.5)
Injection site reaction	24 (12.3)	192 (16.2)	32 (27.6)
Aches & pains	23 (11.8)	215 (18.1)	20 (17.2)
Chills	21 (10.8)	108 (9.1)	3 (2.6)
Fever	19 (9.7)	154 (13.0)	7 (6.0)
Eye sore	18 (9.2)	122 (10.3)	6 (5.2)
Insomnia	18 (9.2)	101 (8.5)	12 (10.3)
Shortness of breath	17 (8.7)	114 (9.6)	6 (5.2)
Shortness of breath	17 (8.7)	114 (9.6)	6 (5.2)
Heart palpitation	15 (7.7)	86 (7.3)	5 (4.3)
Nausea	14 (7.2)	81 (6.8)	4 (3.4)
Rash	10 (5.1)	69 (5.8)	2 (1.7)
Persistent chest pain/pressure	9 (4.6)	66 (5.6)	5 (4.3)
Muscle twitching	8 (4.1)	67 (5.6)	10 (8.6)
Decreased sense of smell	8 (4.1)	51 (4.3)	5 (4.3)
Decreased sense of test	7 (3.6)	48 (4.0)	5 (4.3)