Research Square

Preprints are preliminary reports that have not undergone peer review.
They should not be considered conclusive, used to inform clinical practice,
or referenced by the media as validated information.

The Influence of the Menstrual Cycle on Muscle Injuries - A Systematic Review and Meta-Analysis

Yannik Guthardt
University of Münster
Debby Sargent
Heriot-Watt University
Ross Julian
ross.julian@uni-muenster.de

University of Münster

Article

Keywords: Female Athlete, Follicular Phase, Injury Occurrence, Luteal Phase, Oestrogen, Team Sport

Posted Date: August 4th, 2025

DOI: https://doi.org/10.21203/rs.3.rs-6965721/v1

License: This work is licensed under a Creative Commons Attribution 4.0 International License.
Read Full License

Additional Declarations: No competing interests reported.

Version of Record: A version of this preprint was published at Scientific Reports on January 21st, 2026.
See the published version at https://doi.org/10.1038/s41598-026-36763-0.

Abstract

This systematic review and meta-analysis examined the relationship between menstrual cycle phases and the incidence of muscle injuries in female team sport athletes, following PRISMA 2020 and PERSiST guidelines. A comprehensive search was conducted in PubMed, Scopus, and SPORTDiscus from inception to mid-January 2024. Studies were included if they examined female team sport athletes of reproductive age with regular menstrual cycles and compared the occurrence of muscle injuries across at least two menstrual phases. Studies involving hormonal contraceptive use, medications affecting the menstrual cycle or musculoskeletal system, or menstrual dysfunction were excluded. Three studies met the inclusion criteria, involving 318 participants. Meta-analysis yielded a pooled Risk Ratio of 1.18 (95% CI: 0.75 to 1.86, p = 0.46) for injury risk between the luteal and follicular phases, suggesting no statistically significant association. However, the certainty of the cumulative evidence was rated as very low due to methodological limitations, including inconsistent phase classifications and reliance on imprecise methods for identifying menstrual phases. Future research employing standardised, physiologically accurate methods for classifying and detecting menstrual cycle phases is necessary to better understand the potential links between hormonal fluctuations and injury risk.

1 INTRODUCTION

Over the past few decades, women's sport has seen rapid growth in professionalism and commercialisation due to investments, structured development, and strategic planning [1–4]. This evolution has inherently led to an increase in the physical demands on athletes [5] as well as a rise in the frequency, intensity, and competitiveness of training and competitions [6], potentially elevating the risk of injury [7]. Given the significant health and performance-related impacts of injuries [8–12], there is a pressing need to design and investigate effective injury-mitigation strategies.

Despite the significant underrepresentation of female participants in sport medicine research [13, 14], numerous studies have highlighted notable differences in the predominant injury types, incidence, and burden between male and female athletes [15–18]. These disparities have prompted calls for research on female-specific injury prevention programmes, including the effects of sex-specific biological factors [1, 17, 19].

Arguably, one of the most prominent factors in eumenorrheic female athletes is the menstrual cycle, which has been shown to influence physiological functions and systems [20–22]. These cyclical changes in ovarian hormone concentrations, which can be observed in Fig. 1, lead to distinct hormonal profiles that can be used to identify and differentiate menstrual phases [23, 24]. Based on an idealised 28-day cycle, the menstrual cycle can be divided into the early follicular phase (days 1–5), mid follicular phase (days 6–8), late follicular phase (days 9–13), ovulation (day 14), early luteal phase (days 15–20), mid-luteal phase (days 21–24), and the late luteal phase (days 25–28) [21].

< Fig. 1 **>**

Systematic reviews and meta-analyses examining potential menstrual cycle-related injury risks seem to have primarily focused on the anterior cruciate ligament (ACL) [25–28]. This focus is likely due to previously demonstrated effects of ovarian hormones and the menstrual cycle on collagen formation[29, 30], the structural integrity of the ACL [31, 32], and knee laxity [31, 32]. Though rare, ACL injuries often sideline athletes for several months [33, 34], and jeopardise careers [34].

In contrast, muscle injuries are generally less severe but occur significantly more frequently. For example, in elite-level women's football, muscle injuries in the thigh alone have been observed to occur approximately 16 times more often than ACL injuries [34], with 15–20% of cases sidelining athletes for longer than a month[34]. Muscle strains are also the most common type of injury in both male and female athletes across various sport disciplines [34–37]. Given evidence suggesting that menstrual cycle hormones may affect soft tissue compliance [38–40] and neuromuscular ability [41–43], it seems plausible to assume a possible connection between the menstrual cycle and muscle injury risk. This possibility is further supported by research on changes in proprioception and pain perception during the menstrual cycle. While some studies dismiss menstrual cycle-related variations in proprioception and dynamic stability as major contributors to injury events [44], others suggest that under high athletic demands, these factors could indeed affect injury risk [45].

To the best of the authors' knowledge, this is the first attempt to provide a systematic and comprehensive overview of the influence of the menstrual cycle on muscle injury occurrence. The insights from this review may be valuable for athletes, coaches, practitioners, researchers, and other stakeholders working to understand and prevent muscle injuries.

2 METHODS

This review was pre-registered on the Open Science Framework (OSF), adheres to the PRISMA 2020 statement guidelines [46] and covers all the items of the PRISMA checklist (https://osf.io/bys84/). Further, the Prisma in Exercise, Rehabilitation, Sport medicine and SporTs science (PERSiST) guidance was followed [47].

2.1 Study Inclusion and Exclusion Criteria

The PICOS framework (consisting of Population, Intervention, Comparator, Outcomes, and Study Design) guided the process of determining this review’s relevant parameters.

2.1.1 Population

Participants included female team sport athletes who fulfilled the following criteria: (a) were of reproductive age (post-menarche and premenopausal), (b) had regular menstruations and ovulation cycles, (c) were non-users of any hormonal contraceptives or medications that affect the menstrual cycle or musculoskeletal system, and (d) were free from any menstrual dysfunctions (such as amenorrhea or anovulatory cycles) or other conditions that can influence their menstrual cycle and

hormone profile or musculoskeletal system (such as pregnancy and relative energy deficiency syndrome). No restrictions were imposed regarding the athletes' competition level.

2.1.2 Intervention

No particular intervention was investigated, but participants had to meet the population criteria above. Studies had to verify the participants' menstrual cycle phases through established means, and the classification used for the menstrual cycle phases had to be consistent with the existing literature.

2.1.3 Comparator

To understand the potential association between the menstrual cycle and the occurrence of muscle injuries, included studies were required to compare an outcome measure (i.e., injury number or injury rate incidence) at a minimum of two menstrual cycle phases. Injuries were defined as an occurrence that prevents an athlete from participating in training or match-play for a minimum of a day.

2.1.4 Outcomes

The primary outcome was the occurrence of non-contact muscle injuries. Further, related outcomes such as injury incidence rate and muscular injuries with unclear differentiation between contact and non-contact origin were considered.

2.1.5 Study Design

Studies were considered for inclusion if they met the following criteria: (a) fully published in a peer-reviewed journal, (b) written in English or translated and published in English, and (c) had the primary or secondary objective of assessing the incidence of muscle injuries within the phases of the menstrual cycle. Any reviews, case reports, editorials, conference abstracts, clinical commentaries, dissertations, and unpublished studies were excluded. No restrictions were placed on the date of publication.

2.2 Search Strategy

A systematic electronic literature search was conducted independently in mid-January 2024 by two reviewers (YG and DS), each using the three databases PubMed (including MEDLINE), Scopus and SPORTDiscus to identify all relevant articles. The following search string was used: ("muscle*" OR "muscular") AND (injur* OR strain* OR tear* OR rupture* OR "injury incidence") AND ("menstrual cycle" OR "menstrual phase" OR "menstrual" OR "menstruation" OR "follicular phase" OR "luteal phase" OR "ovulation" OR "ovulatory" OR "sex hormone*"). Databases were searched from inception onwards. The reference lists of relevant articles obtained were hand-searched to identify further potential studies that could be added manually.

2.3 Study Selection, Data Extraction and Quality Assessment

2.3.1 Screening and Selection of Eligible Studies

All search results were saved and managed in the systematic review software ‘Rayyan’ [48]. This tool was utilised to sort through, screen, and include qualified records. Any duplicates were automatically identified by comparing title, year, volume, and authorship. Afterwards, two reviewers (YG and DS) independently verified the accuracy of duplicates before removing them from consideration.

All the remaining articles underwent an independent two-phase screening strategy by two reviewers (YG and DS). In phase 1, the titles and abstracts were examined against the predetermined eligibility criteria. If neither the title nor the abstract of an article showed indications of meeting the inclusion criteria or met at least one of the exclusion criteria, the article was excluded in phase 1.

The full-text versions of the remaining articles were then read as phase 2 to confirm eligibility. In cases where studies were reported in multiple publications, all reports were collated. In case of disagreement on eligibility between the two reviewers, the most experienced reviewer (RJ) was consulted, and his decision was deemed final.

2.3.2 Data Extraction

Two reviewers (YG and DS) independently extracted the data using a standardised template. Any discrepancies were identified and addressed through a consensus-based discussion and reviewing the original article. The matter was referred to the most experienced researcher (RJ) for consultation in unresolved disagreements. A comprehensive list of the extracted data items is provided on OSF (https://osf.io/bys84).

2.3.3 Quality Assessment

The quality assessment was conducted independently by two reviewers (YG and DS) following the Grading of Recommendations Assessment Development and Evaluation (GRADE) system of rating quality of evidence [49]. This approach evaluates the certainty in the cumulative evidence based on five domains: risk of bias, indirectness, inconsistency, imprecision, and evidence of publication bias. Each study was individually assessed for risk of bias with the QUIPS (Quality In Prognosis Studies) appraisal tool [50]. Based on the QUIPS tool results, each study was assigned an adequate a priori quality rating of either ‘high’, ‘moderate’, ‘low’ or ‘very low’.

Following McNulty et al. [22] and based on the recommendations of De Jonge et al. [51], the initial rating was either maintained or downgraded based on two questions considered vital to assessing the indirectness of the research studies:

(*Q1) Was the menstrual cycle phase confirmed using blood samples or urinary ovulation detection kits?* The initial rating was maintained if the study confirmed the menstrual cycle phase using these methods. If not, the study was downgraded by one level. For example, a study initially rated as ‘high’ would be downgraded to ‘moderate’ if it did not use blood samples or urinary detection kits for confirmation.

(*Q2) Was the injury medically diagnosed by qualified experts or means?* The rating was downgraded by one level if the study did not report that qualified medical personnel recorded and diagnosed injuries. If

medical staff were involved in diagnosing injuries, the Q1 rating was maintained.

Consistency was ascertained through meta-analysis, visual inspection of effect size estimates, and overlap of confidence intervals, supplemented by statistical tests for heterogeneity. Precision was judged by closely examining the number of data points supporting the relevant outcome (with outcomes based on < 5 data points being downgraded) and visual analysis of the width of the confidence intervals. The assessment of publication bias included visual examination of result patterns (funnel plot), Egger's test and the recommendations of the GRADE workgroup [52].

These procedures collectively led to a final certainty rating for the cumulative evidence as 'high', 'moderate', 'low', or 'very low' [53]. Disagreements were resolved through discussion. If no consensus was reached, a third reviewer (RJ) made the final decision. Based on this appraisal strategy, no studies were excluded.

2.4 Data Synthesis

To facilitate a consistent analysis across studies, each study's menstrual cycle phases were aligned according to a predefined two-phase classification scheme commonly employed by prior menstrual cycle research [51]. The follicular phase was defined as extending from the onset of menstruation up to ovulation, and the luteal phase was considered to encompass ovulation and the remaining days until the start of the next menstrual period. The extracted data was utilised to align the phases of each eligible study to the predetermined classification for statistical analysis. This alignment was carried out based on the days of a 28-day idealised cycle length or the contextual information provided by the study in cases where the precise duration of each classified phase was ambiguous. Ovulation was deemed to start on day 13 of a 28-day idealised menstrual cycle in line with McNulty et al. [22]. Since none of the eligible studies focused exclusively on muscle injuries, the absolute number of muscle injuries was selected as the primary outcome for statistical analysis, as it was the only consistent and comparable metric across all studies.

A frequentist approach was employed for this meta-analysis, using the Risk Ratio (RR) as the primary effect size. This choice allowed for a direct comparison of the injury occurrence between the follicular and luteal phases, the latter of which acted as the 'control condition' across the selected studies. For each study, the RR was calculated as:

$$RR = \frac{Number\ of\ Injuries\ in\ Luteal\ Phase}{Number\ of\ Injuries\ in\ Follicular\ Phase}$$

The log-transformed Risk Ratio (logRR) and its associated standard error (SE) were computed to standardise the effect sizes across studies, facilitating pooling across studies. The meta-analysis was conducted using both fixed-effect (common-effect) and random-effects models to explore the impact of potential heterogeneity among studies. The fixed-effect model assumes that all studies estimate a common underlying effect size, attributing any observed differences solely to within-study variation. In

contrast, the random-effects model accounts for both within and between-study heterogeneity, assuming that the true effect size may vary across studies.

In both models, weights were assigned to each study based on the inverse of the logRR's variance. The random-effects model further adjusted the weights to account for between-study variability.

Heterogeneity across the studies was assessed using the I^2 statistic, which quantifies the proportion of total variation due to between-study heterogeneity. An I^2 value of 100% indicates maximal inconsistency.

Further, tau^2 was calculated using the Restricted Maximum Likelihood (REML) method to estimate the between-study variance, providing a refined measure of variability beyond what is captured by I^2 alone. Cochran's Q statistic was used to test the null hypothesis that all studies evaluate the same effect. A leave-one-out sensitivity analysis was conducted to ensure the robustness of the findings.

The results were visually summarised using a forest plot, which displays each study's individual RR and 95% confidence intervals (CIs) and the pooled effect sizes from both models. The forest plot also illustrates the weight assigned to each study, emphasising each contribution to the overall effect estimate. All analyses were conducted in the statistical software R Version 4.2.2 [54], including the use of the R package meta [55].

3 RESULTS

3.1 Literature Search

Figure 2 illustrates the search and selection of studies in a flow chart.

< Fig. 2 **>**

3.2 Study Characteristics

The final analysis included data from three studies [56–58]. The combined total number of participants observed in two [56, 58] of the three studies was 205, while the total number of participants observed in the third study was not ascertainable. However, all relevant injury data in this study were obtained from 113 participants [57]. Two studies were conducted in professional football and one in professional futsal. All studies assessed the menstrual data via self-reporting of the players through either a mobile menstrual tracking application [58], a combination of self-reported menstrual cycle length and a regression equation [57], or a mixture of calendar-based counting and different mobile applications [56]. Details of the study characteristics of the included studies can be found in Table 1.

Table 1
Study Characteristics of Included Studies

Study	Sample	Menstrual Cycle Classification	Method of MC Phase Verification	Method of Injury Diagnosis	Findings
Lago-Fuentes et al. [56]	N = 179 Players of the Spanish First and Second National Futsal League	*Follicular Phase* (Day 1–12) *Ovulatory Phase* (Day 13–15) *Luteal Phase* (Day 16–28)	Self-report via calendar-based counting and different apps	No Information	No statistical differences in the injury occurrence between the MC phases. Tendencies of higher frequencies in the *Follicular Phase*.
Martin et al. [57]	N = eight playing squads over four years, comprised of 3,947 individual player camp attendances (Injury data was obtained from n = 113) Players selected for the England National Football Team Under 15s – Senior Level	*Follicular Phase* (Time between the first day of the menses and the late Follicular Phase) *Late Follicular Phase* (Day of luteinising hormone peak and the two preceding days) *Luteal Phase* (Any time point following the *Late Follicular Phase*)	Self-report via typical cycle length and regression equation	Recorded by each team's medical support staff and classified using the Orchard Sports Injury Classification System by a medical professional	Muscle injuries were approximately twice as common in the *Late Follicular Phase* compared to the *Follicular Phase* and *Luteal Phase* per 1,000 person-days
Barlow et al. [58]	N = 26 24.1 ± 4.6 years of age Players of a professional Women's Super League Football Club	*Phase 1* (Menstruation) *Phase 2* (Remainder of the predicted follicular phase) *Phase 3*	Self-report via a mobile tracking application that recorded menstruation days and intensity of flow	Recorded by the football club's medical support staff and classified using the Orchard Sports Injury Classification System	Muscle injuries occurred more commonly in *Phase 3* and *Phase 4* than in *Phase 1* and *2.* Compared to *Phase 1*, muscle injuries occurred three times more likely in *Phase 2*, five times more

Study	Sample	Menstrual Cycle Classification	Method of MC Phase Verification	Method of Injury Diagnosis	Findings
		(Majority of the luteal phase) *Phase 4* (Premenstrual window, defined as the five days before the onset of menstruation)			likely in *Phase 3*, and over six times more likely in *Phase 4* per 1,000 person-days

3.3 Study Findings

Muscle injury occurrence across the studies showed variation between menstrual cycle phases, with inconsistent results reported. One study [57] observed a considerable increase in muscular injuries during the late follicular phase, as defined in their research. In contrast, another study reported the highest injury rates during the luteal phase, particularly in the premenstrual window [58]. A third study [56] indicated a tendency toward higher injury frequencies in the follicular phase compared to the other phases they defined, although these differences were not statistically significant. Detailed findings from each study are presented in Table 1.

3.4 Quality Assessment of Included Studies

Three studies (100%) were considered to be at high risk of bias after the risk of bias assessment of the individual study level obtained from the QUIPS tool and additional questions regarding the menstrual cycle phase and injury detection and verification. All studies were allocated an a priori rating of ‘moderate’ but downgraded based on the additional questions (Q1) and (Q2) regarding the method of menstrual phase verification and injury diagnosis and recording. Details of the domain-based risk of bias assessment from the QUIPS tool and (Q1) and (Q2) can be found on OSF (https://osf.io/bys84).

No asymmetry was observed following a visual examination of the funnel plot. Egger’s test also indicated no detectable publication bias ($p = 0.92$). However, because all included papers are observational studies, the recommendations of the GRADE workgroup were followed, and publication bias was considered to be inherently substantial [52]. Based on the GRADE approach, the certainty of the cumulative evidence of this review was assessed to be ‘very low’ [53].

3.5 Meta-Analysis Results

The common-effect model meta-analysis revealed a pooled Risk Ratio of 1.18 (95% CI: 0.86 to 1.60, $z = 1.03$, $p = 0.30$), suggesting no statistically significant difference in injury occurrence between the luteal and follicular phases. The random-effects model, which accounts for between-study heterogeneity, produced a similar pooled RR of 1.18 (95% CI: 0.75 to 1.86, $z = 0.73$, $p = 0.46$), further indicating the

absence of a significant association between the menstrual cycle phase and muscle injury occurrence. The results of the meta-analysis are visualised in Fig. 3.

The heterogeneity across studies was moderate, with an I^2 of 49.5% (95% CI: 0.0–85.3%) and a tau^2 of 0.08 (95% CI: 0.00 to 7.21). Cochran's Q test for heterogeneity yielded a Q value of 3.96 (d.f. = 2, p = 0.14), indicating that the observed variation across studies could be due to chance rather than true heterogeneity.

< Fig. 3 **>**

The leave-one-out sensitivity analysis to examine each study's influence on the overall results elicited varying RRs from 1.01 to 1.40, with corresponding 95% CIs that consistently overlapped and included the null value (RR = 1.0), indicating no significant effect. The pooled RR increased slightly to 1.22 (95% CI: 0.50 to 2.98, z = 0.43, p = 0.67), and the highest heterogeneity was observed (I^2 = 74.6%, Q = 3.93, d.f. = 1, p = 0.05) with the exclusion of the Lago-Fuentes et al. paper [56], suggesting a significant impact on the overall results. In contrast, the lowest heterogeneity was observed with an I^2 of 3.6% (Q = 1.04, d.f. = 1, p = 0.31) with the exclusion of the Barlow et al. study [58], resulting in a pooled RR of 1.01 (95% CI: 0.71 to 1.45, z = 0.07, p = 0.94).

4 DISCUSSION

This systematic review and meta-analysis sought to elucidate whether the menstrual cycle and its phases have a significant impact on the occurrence of muscle injuries in eumenorrheic female team sport athletes. The premise that there might be variability in muscle injuries across the menstrual cycle is grounded in previous findings on the effects of the inherent hormonal fluctuations on musculoskeletal function introduced at the beginning of this review, including altered soft tissue plasticity [38–40], impacted collagen metabolism [29, 30], and potentially distorted neuromuscular control [41–43] and proprioception [45].

Despite the biological and theoretical plausibility, the current meta-analysis did not find a statistically significant difference in the distribution of muscle injuries between the follicular and luteal phases of the menstrual cycle. The pooled Risk Ratio of 1.18, derived from both fixed-effect and random-effects models, indicates a non-significant association between the menstrual cycle phase and the instance of muscular injuries. This suggests that while hormonal fluctuations may influence musculoskeletal properties, these changes may not translate into a measurable difference in injury risk across the menstrual cycle. Notably, this conflicts with the athletes' perception, as over half of female athletes report a changed perceived risk of injury throughout the menstrual cycle [59]. When examining the RRs and conclusions from the individual studies, discordant and conflicting results become apparent, mirroring previous research predominantly centred on ACL injuries with similarly inconsistent findings [27, 28]. While these findings provide valuable information and appear to correspond loosely with existing evidence from similar research, they should be interpreted with caution due to the limited available literature and several notable limitations in the current review and the included studies.

The two paramount limitations are (a) the inconsistency and dissimilarity in the classification of the menstrual cycle phases across scientific literature and (b) the methods used by the included studies to detect and verify the menstrual cycle phases.

The inconsistent classification of menstrual phases across studies poses significant challenges for comparing and interpreting findings. There is considerable variation in how the menstrual cycle is divided, with differences in the number of phases, terminology, duration, and specific time points assigned to each phase. This variety is visualised in Fig. 4. As a result of this, events occurring at the same specific time point in the menstrual cycle might be interpreted and reported in completely unrelated and conflicting phases across the scientific literature. This issue is exacerbated when attempting to synthesise evidence and data across studies, as the heterogeneity introduced by different phase definitions complicates meta-analytical approaches, increasing the risk of biased, erroneous, or misleading findings.

< Fig. 4 **>**

In order to statistically synthesise and analyse the data of the included studies for this review, all respective phase classifications were aligned to a two-phase system, creating a follicular phase and a luteal phase. While this allowed for statistical analysis across the findings of the included studies and aligns with the majority of previous menstrual cycle research, which typically distinguishes and compares these two phases [51], it is vital to note that it does not accurately reflect the complex dynamic of the typical menstrual cycle, involving multiple fluctuations in oestrogen and progesterone levels at various stages.

Given the non-linear relationship between circulating hormones and their physiological effects, these distinct changes in the hormonal milieu and specific hormonal peaks may have substantial impacts on key biological factors related to injury risk, including muscle proteostasis and tissue stiffness [60]. This may be further magnified by evidence that mRNA and protein levels of oestrogen and progesterone receptors vary across the course of the menstrual cycle [61]. Thus, reducing the menstrual cycle to unsuitably broad and generalised phases, as was possibly done for the analysis, may overlook critical hormonal fluctuations and their potential effects.

Furthermore, the identification and verification of the menstrual cycle phases in the included studies were substantially based on self-reported data and calendar-based counting methods, which are inherently prone to inaccuracies. These methods, while convenient and considerably more cost-effective compared to hormonal assays, rely on temporal estimates and are susceptible to miscalculations and fallacious reflections of an individual's menstrual cycle due to a wide range of factors, including inter- and intra-individual variability in the cycle length and ovulation timing, as well as further unrecognised variability in the cycle induced through stress, exercise, illness, and changes in body weight [62–64]. Additionally, these methods typically define menstrual cycle phases based on self-reported data regarding the onset of menstruation. However, it is essential to recognise that the occurrence of regular menstrual bleeding does not necessarily indicate an ovulatory cycle with the corresponding typical

hormonal course [51, 65]. As a result, the sole use of calendar-based methods for phase identification is considered insufficiently accurate for reliably determining the menstrual cycle phases [51].

Several confounding factors may have obscured potential associations between menstrual cycle phases and muscle injuries. These possible confounders include, but are not limited to, variations in training load, cumulative training and game exposure, psychophysiological influences such as stress and fatigue, and, most notably, the athletes' previous injury history, given that previous muscle injury is widely recognised as the most significant risk factor for future muscle injury [66]. This complexity is exemplified by Lago-Fuentes et al. [56], who observed a higher incidence of injuries during the season's first quarter, attributing this to increased training load during that period. Further, they reported that over 60% of injuries occurred in the later stages of training sessions, highlighting the possible role of fatigue.

Moreover, the Lago-Fuentes et al. study [56] collected data over two full seasons but only excluded and controlled for hormonal contraceptives in the second season. The relatively low percentage (5%) of excluded hormonal contraceptive users in the second season might suggest a limited impact on the data from the first season and ultimately led to the study's inclusion after consensus-based discussion. However, it is important to acknowledge that a small, albeit unknown, proportion of data from participants using hormonal contraceptives was included in this review's final analysis.

Similarly, the authors cannot guarantee that all injury data were derived from non-contact events. While muscle injuries, particularly strains and tears, are typically attributed to intrinsic factors, the injury mechanism, stemming from excessive tensile or shear forces leading to the failure of muscle fibres and their surrounding connective tissue, can also result from external forces such as traumatic contact [66]. Any muscle injuries caused by external contact included in this review might have skewed the results. Furthermore, a recent scoping review [67] emphasised significant inconsistencies in how injuries are defined and measured across studies investigating the association between menstrual cycle phases and musculoskeletal injuries, thereby further hindering comparative research efforts.

The findings and limitations of this review highlight the need for more robust and well-designed research to explore the relationship between the menstrual cycle and muscle injury risk. Future research should prioritise the use of hormonal assays, such as serum measurements of oestrogen and progesterone, as well as urinary ovulation detection kits, to accurately verify menstrual cycle phases. Adopting the methodological recommendations, guidelines, and protocols established by DeJonge et al. [51] to strengthen quality across the scientific body. In addition, the classification scheme of the menstrual cycle needs to be standardised to simplify evidence synthesis and mitigate the loss of intricate data details through necessary phase alignment. The authors recommend using the classification system previously used by McNulty et al. [22] (visualised in Fig. 4), given that it includes the three most distinctive hormonal profiles of the menstrual cycle, namely, the early follicular phase with low oestrogen and low progesterone, the late follicular phase with high oestrogen and low progesterone, and the mid-luteal phase with high oestrogen and high progesterone, as reported by DeJonge and colleagues [51].

This approach may uncover more subtle associations between hormonal fluctuations and injury risk that were not detectable in the current analysis. It could also clarify the extent to which hormonal effects, such as progesterone's commonly associated stabilising influence on collagen [68, 69] and oestrogen's regulatory role in tissue metabolism [60, 70], contribute to injury susceptibility in this context. By elucidating these relationships, this approach could serve as a foundation for investigating potential causative links, rather than mere associations, between the hormonal fluctuations of the menstrual cycle and muscular injuries.

The authors acknowledge that these recommendations for future research pose substantial challenges, including the invasiveness of procedures, associated costs, and the need for specialised facilities and personnel, which may seem discouraging. However, research in this area is critically needed, and finding a compromise is essential for advancing the field. Based on DeJonge and colleagues' recommendations [51], this review suggests that studies unable to incorporate direct hormone measurements may still achieve sufficient accuracy by employing large sample sizes and using urinary detection kits to measure luteinising hormone.

It is important to recognise that categorising injury events into specific menstrual cycle phases will inevitably involve some degree of approximation, estimation, and retrospective day-counting. However, the absence of direct physiological measurements significantly amplifies the risk of inaccuracies, misrepresentations, and misclassifications, thereby undermining the ability to draw robust conclusions about the influence of menstrual cycle phases on injury occurrence. Ultimately, the accuracy of scientific findings must be solid and reliable, ensuring that their applicability to real-world settings is not compromised by a foundation built on uncertainty.

5 CONCLUSION

This meta-analysis did not reveal a statistically significant association between menstrual cycle phases and muscle injury incidence in female team sport athletes. However, the divergence between this finding and athletes' reports of perceived cyclical vulnerability, along with the limitations of the included studies, suggest that the true relationship may be more complex than currently understood. Although the theoretical basis for hormonal influences on tissue properties is compelling, methodological limitations, most notably inconsistent phase classification and the use of imprecise, self-reported measures for menstrual phase detection, may have masked subtle but clinically relevant effects. Future research in this field is needed and should be obliged to adopt standardised, physiologically accurate methods for detecting and classifying menstrual phases to enhance the comparability of studies and ultimately lead to more effective injury prevention strategies tailored to female athletes.

Statements and Declarations

Contribution YG designed the research, conducted the systematic literature search and screening of the search results (including the two-phase screening process), extracted the data, performed all statistical

analyses, and drafted the manuscript with critical input from the other authors. DS conducted the systematic literature search and screening of the search results (including the two-phase screening process), extracted the data, and provided critical input on the manuscript. RJ designed the research, provided critical input on the manuscript, and serves as the corresponding author, attesting that all listed authors meet authorship criteria and that no others meeting the criteria have been omitted. All authors had full access to all the data in this review, take full responsibility as guarantors for the integrity of the data and the accuracy of the data analyses, contributed to data interpretation, critically revised the manuscript, and have read and approved the final version of the manuscript.

Registration The study was registered at Open Science Framework (https://osf.io/mr6y8)

Data Availabilty The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Funding This research received no grant from any funding agency in the public, commercial or not-for-profit sectors.

Competing Interests Yannik Guthardt, Ross Julian, and Debby Sargent declare no competing interests

Ethics Approval Not applicable.

Consent to Participate Not applicable.

Consent for Publication Not applicable.

Code Availability Not applicable.

References

1. Fédération Internationale de Football Association. Women's Football Strategy. FIFA Strasse 20, P.O Box 8044 Zurich, Switzerland 2018.
2. International Olympic Committee. Factsheet: Women in the Olympic Movement 2023.
3. The Football Association. Inspiring Positive Change - The FA Strategy for Women's and Girl's Football: 2020-2024 2020.
4. Women’s Sport Trust. Visibility Uncovered 2023 - The Year In Review 2024.
5. Martínez-Lagunas V, Niessen M, Hartmann U. Women's football: Player characteristics and demands of the game. *Journal of Sport and Health Science* 2014;3(4):258–72. https://www.sciencedirect.com/science/article/pii/S2095254614000982.
6. Klein M-L. Women’s Football Leagues in Europe: Organizational and Economic Perspectives. In: Pfister G, Pope S, eds. Female Football Players and Fans. London: Palgrave Macmillan UK 2018:77–101.

7. Orejel Bustos A, Belluscio V, Camomilla V, et al. Overuse-Related Injuries of the Musculoskeletal System: Systematic Review and Quantitative Synthesis of Injuries, Locations, Risk Factors and Assessment Techniques. *Sensors (Basel)* 2021;21(7). doi:10.3390/s21072438 [published Online First: 1 April 2021].
8. Lohmander LS, Ostenberg A, Englund M, et al. High prevalence of knee osteoarthritis, pain, and functional limitations in female soccer players twelve years after anterior cruciate ligament injury. *Arthritis Rheum* 2004;50(10):3145–52.
9. Drew MK, Raysmith BP, Charlton PC. Injuries impair the chance of successful performance by sportspeople: a systematic review. *Br J Sports Med* 2017;51(16):1209–14. doi:10.1136/bjsports-2016-096731 [published Online First: 26 April 2017].
10. Haugen E. Athlete Mental Health & Psychological Impact of Sport Injury. *Operative Techniques in Sports Medicine* 2022;30(1).
11. Eliakim E, Morgulev E, Lidor R, et al. Estimation of injury costs: financial damage of English Premier League teams' underachievement due to injuries. *BMJ Open Sport Exerc Med* 2020;6(1):e000675. doi:10.1136/bmjsem-2019-000675 [published Online First: 20 May 2020].
12. Pulici L, Certa D, Zago M, et al. Injury Burden in Professional European Football (Soccer): Systematic Review, Meta-Analysis, and Economic Considerations. *Clin J Sport Med* 2022. doi:10.1097/JSM.0000000000001107 [published Online First: 22 November 2022].
13. Emmonds S, Heyward O, Jones B. The Challenge of Applying and Undertaking Research in Female Sport. *Sports Med Open* 2019;5(1):51. doi:10.1186/s40798-019-0224-x [published Online First: 12 December 2019].
14. Costello JT, Bieuzen F, Bleakley CM. Where are all the female participants in Sports and Exercise Medicine research? *Eur J Sport Sci* 2014;14(8):847–51. doi:10.1080/17461391.2014.911354 [published Online First: 25 April 2014].
15. Hollander K, Rahlf AL, Wilke J, et al. Sex-Specific Differences in Running Injuries: A Systematic Review with Meta-Analysis and Meta-Regression. *Sports Med* 2021;51(5):1011–39. doi:10.1007/s40279-020-01412-7 [published Online First: 12 January 2021].
16. Klein C, Henke T, Platen P. Injuries in football (soccer)—a systematic review of epidemiology and aetiological aspects. *Ger J Exerc Sport Res* 2018;48(3):309–22. https://link.springer.com/content/pdf/10.1007/s12662-018-0530-3.pdf.
17. Larruskain J, Lekue JA, Diaz N, et al. A comparison of injuries in elite male and female football players: A five-season prospective study. *Scand J Med Sci Sports* 2018;28(1):237–45. doi:10.1111/sms.12860 [published Online First: 27 March 2017].
18. Lin CY, Casey E, Herman DC, et al. Sex Differences in Common Sports Injuries. *PM R* 2018;10(10):1073–82. doi:10.1016/j.pmrj.2018.03.008 [published Online First: 14 March 2018].
19. Edouard P, Feddermann-Demont N, Alonso JM, et al. Sex differences in injury during top-level international athletics championships: surveillance data from 14 championships between 2007 and

2014. *Br J Sports Med* 2015;49(7):472–77. doi:10.1136/bjsports-2014-094316 [published Online First: 24 January 2015].

20. Driver HS, Werth E, Dijk D-J, et al. The Menstrual Cycle Effects on Sleep. *Sleep Medicine Clinics* 2008;3(1):1–11.
21. Pitchers G, Elliott-Sale K. Considerations for coaches training female athletes. *Prof Strength Cond.* 2019;55:19–30.
22. McNulty KL, Elliott-Sale KJ, Dolan E, et al. The Effects of Menstrual Cycle Phase on Exercise Performance in Eumenorrheic Women: A Systematic Review and Meta-Analysis. *Sports Med* 2020;50(10):1813–27.
23. Janse de Jonge XAK. Effects of the menstrual cycle on exercise performance. *Sports Med* 2003;33(11):833–51.
24. Mihm M, Gangooly S, Muttukrishna S. The normal menstrual cycle in women. *Anim Reprod Sci* 2011;124(3-4):229–36. doi:10.1016/j.anireprosci.2010.08.030 [published Online First: 3 September 2010].
25. Hewett TE, Zazulak BT, Myer GD. Effects of the menstrual cycle on anterior cruciate ligament injury risk: a systematic review. *Am J Sports Med* 2007;35(4):659–68. doi:10.1177/0363546506295699 [published Online First: 9 February 2007].
26. Moriceau J, Fevre A, Domínguez-Balmaseda D, et al. The Influence of the Menstrual Cycle and Oral Contraceptives on Knee Laxity or Anterior Cruciate Ligament Injury Risk: A Systematic Review. *Applied Sciences* 2022;12(24):12627.
27. Somerson JS, Isby IJ, Hagen MS, et al. The Menstrual Cycle May Affect Anterior Knee Laxity and the Rate of Anterior Cruciate Ligament Rupture: A Systematic Review and Meta-Analysis. *JBJS Rev* 2019;7(9):e2.
28. Dos'Santos T, Stebbings GK, Morse C, et al. Effects of the menstrual cycle phase on anterior cruciate ligament neuromuscular and biomechanical injury risk surrogates in eumenorrheic and naturally menstruating women: A systematic review. *PLoS One* 2023;18(1):e0280800. doi:10.1371/journal.pone.0280800 [published Online First: 26 January 2023].
29. Yu WD, Panossian V, Hatch JD, et al. Combined effects of estrogen and progesterone on the anterior cruciate ligament. *Clin Orthop Relat Res* 2001(383):268–81.
30. Yu WD, Liu SH, Hatch JD, et al. Effect of estrogen on cellular metabolism of the human anterior cruciate ligament. *Clin Orthop Relat Res* 1999(366):229–38.
31. Park S-K, Stefanyshyn DJ, Loitz-Ramage B, et al. Changing hormone levels during the menstrual cycle affect knee laxity and stiffness in healthy female subjects. *Am J Sports Med* 2009;37(3):588–98. doi:10.1177/0363546508326713 [published Online First: 27 January 2009].
32. Park S-K, Stefanyshyn DJ, Ramage B, et al. Relationship between knee joint laxity and knee joint mechanics during the menstrual cycle. *Br J Sports Med* 2009;43(3):174–79. doi:10.1136/bjsm.2008.049270 [published Online First: 26 August 2008].

33. Ekstrand J, Krutsch W, Spreco A, et al. Time before return to play for the most common injuries in professional football: a 16-year follow-up of the UEFA Elite Club Injury Study. *Br J Sports Med* 2020;54(7):421–26. doi:10.1136/bjsports-2019-100666 [published Online First: 10 June 2019].
34. Hallén A, Tomás R, Ekstrand J, et al. UEFA Women's Elite Club Injury Study: a prospective study on 1527 injuries over four consecutive seasons 2018/2019 to 2021/2022 reveals thigh muscle injuries to be most common and ACL injuries most burdensome. *Br J Sports Med* 2024;58(3):128–35. doi:10.1136/bjsports-2023-107133 [published Online First: 7 February 2024].
35. Ekstrand J, Hägglund M, Waldén M. Injury incidence and injury patterns in professional football: the UEFA injury study. *Br J Sports Med* 2011;45(7):553–58. doi:10.1136/bjsm.2009.060582 [published Online First: 23 June 2009].
36. Jacobsson J, Timpka T, Kowalski J, et al. Injury patterns in Swedish elite athletics: annual incidence, injury types and risk factors. *Br J Sports Med* 2013;47(15):941–52. doi:10.1136/bjsports-2012-091651 [published Online First: 29 March 2013].
37. Panagodage Perera NK, Kountouris A, Kemp JL, et al. The incidence, prevalence, nature, severity and mechanisms of injury in elite female cricketers: A prospective cohort study. *J Sci Med Sport* 2019;22(9):1014–20. doi:10.1016/j.jsams.2019.05.013 [published Online First: 25 May 2019].
38. Ham S, Kim S, Choi H, et al. Greater Muscle Stiffness during Contraction at Menstruation as Measured by Shear-Wave Elastography. *Tohoku J Exp Med* 2020;250(4):207–13.
39. Sung E-S, Kim J-H. The difference effect of estrogen on muscle tone of medial and lateral thigh muscle during ovulation. *J Exerc Rehabil* 2018;14(3):419–23. doi:10.12965/jer.1836110.055 [published Online First: 30 June 2018].
40. Yim J, Petrofsky J, Lee H. Correlation between Mechanical Properties of the Ankle Muscles and Postural Sway during the Menstrual Cycle. *Tohoku J Exp Med* 2018;244(3):201–07.
41. Weidauer L, Zwart MB, Clapper J, et al. Neuromuscular performance changes throughout the menstrual cycle in physically active females. *J Musculoskelet Neuronal Interact* 2020;20(3):314–24.
42. Graja A, Kacem M, Hammouda O, et al. Physical, Biochemical, and Neuromuscular Responses to Repeated Sprint Exercise in Eumenorrheic Female Handball Players: Effect of Menstrual Cycle Phases. *J Strength Cond Res* 2022;36(8):2268–76. doi:10.1519/JSC.0000000000003556 [published Online First: 10 March 2020].
43. Ansdell P, Brownstein CG, Škarabot J, et al. Menstrual cycle-associated modulations in neuromuscular function and fatigability of the knee extensors in eumenorrheic women. *J Appl Physiol (1985)* 2019;126(6):1701–12. doi:10.1152/japplphysiol.01041.2018 [published Online First: 7 March 2019].
44. Legerlotz K, Bey ME, Götz S, et al. Constant performance in balance and proprioception tests across the menstrual cycle - a pilot study in well trained female ice hockey players on hormonal contraception. *Health Sci Rep* 2018;1(1):e18. doi:10.1002/hsr2.18 [published Online First: 24 November 2017].

45. Frizziero A, Demeco A, Oliva F, et al. Changes in Proprioceptive Control in the Menstrual Cycle: a Risk Factor for Injuries? A Proof-of-Concept Study. *Muscle Ligaments and Tendons J* 2023;13(03):360.
46. Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021;372:n71. doi:10.1136/bmj.n71 [published Online First: 29 March 2021].
47. Ardern CL, Büttner F, Andrade R, et al. Implementing the 27 PRISMA 2020 Statement items for systematic reviews in the sport and exercise medicine, musculoskeletal rehabilitation and sports science fields: the PERSiST (implementing Prisma in Exercise, Rehabilitation, Sport medicine and SporTs science) guidance. *Br J Sports Med* 2022;56(4):175–95. doi:10.1136/bjsports-2021-103987 [published Online First: 8 October 2021].
48. Ouzzani M, Hammady H, Fedorowicz Z, et al. Rayyan-a web and mobile app for systematic reviews. *Syst Rev* 2016;5(1):210. doi:10.1186/s13643-016-0384-4 [published Online First: 5 December 2016].
49. Guyatt G, Oxman AD, Akl EA, et al. GRADE guidelines: 1. Introduction-GRADE evidence profiles and summary of findings tables. *J Clin Epidemiol* 2011;64(4):383–94. doi:10.1016/j.jclinepi.2010.04.026 [published Online First: 31 December 2010].
50. Hayden JA, van der Windt DA, Cartwright JL, et al. Assessing bias in studies of prognostic factors. *Ann Intern Med* 2013;158(4):280–86.
51. Janse DE Jonge X, Thompson B, Han A. Methodological Recommendations for Menstrual Cycle Research in Sports and Exercise. *Med Sci Sports Exerc* 2019;51(12):2610–17.
52. Guyatt GH, Oxman AD, Montori V, et al. GRADE guidelines: 5. Rating the quality of evidence--publication bias. *J Clin Epidemiol* 2011;64(12):1277–82. doi:10.1016/j.jclinepi.2011.01.011 [published Online First: 30 July 2011].
53. Balshem H, Helfand M, Schünemann HJ, et al. GRADE guidelines: 3. Rating the quality of evidence. *J Clin Epidemiol* 2011;64(4):401–06. doi:10.1016/j.jclinepi.2010.07.015 [published Online First: 5 January 2011].
54. R Core Team. R: A Language and Environment for Statistical. Vienna, Austria: R Foundation for Statistical Computing.
55. Balduzzi S, Rücker G, Schwarzer G. How to perform a meta-analysis with R: a practical tutorial. *Evid Based Ment Health* 2019;22(4):153–60. doi:10.1136/ebmental-2019-300117 [published Online First: 28 September 2019].
56. Lago-Fuentes C, Padrón-Cabo A, Fernández-Villarino M, et al. Follicular phase of menstrual cycle is related to higher tendency to suffer from severe injuries among elite female futsal players. *Phys Ther Sport* 2021;52:90–96. doi:10.1016/j.ptsp.2021.08.008 [published Online First: 21 August 2021].
57. Martin D, Timmins K, Cowie C, et al. Injury Incidence Across the Menstrual Cycle in International Footballers. *Front Sports Act Living* 2021;3:616999. doi:10.3389/fspor.2021.616999 [published Online First: 1 March 2021].
58. Barlow A, Blodgett JM, Williams S, et al. Injury Incidence, Severity and Type across the Menstrual Cycle in Female Footballers: A Prospective Three Season Cohort Study. *Med Sci Sports Exerc* 2024.

doi:10.1249/MSS.0000000000003391 [published Online First: 8 February 2024].

59. Hayward E, Akam L, Hunter D, et al. Role of the Menstrual Cycle on Performance and Injury Risk: A Survey of Female Professional Rugby Players in the United Kingdom. *Int J Environ Res Public Health* 2024;21(2). doi:10.3390/ijerph21020150 [published Online First: 29 January 2024].
60. Chidi-Ogbolu N, Baar K. Effect of Estrogen on Musculoskeletal Performance and Injury Risk. *Front Physiol* 2018;9:1834. doi:10.3389/fphys.2018.01834 [published Online First: 15 January 2019].
61. Ekenros L, Papoutsi Z, Fridén C, et al. Expression of sex steroid hormone receptors in human skeletal muscle during the menstrual cycle. *Acta Physiol (Oxf)* 2017;219(2):486–93. doi:10.1111/apha.12757 [published Online First: 9 August 2016].
62. Barrett ES, Thune I, Lipson SF, et al. A factor analysis approach to examining relationships among ovarian steroid concentrations, gonadotrophin concentrations and menstrual cycle length characteristics in healthy, cycling women. *Hum Reprod* 2013;28(3):801–11. doi:10.1093/humrep/des429 [published Online First: 18 December 2012].
63. Harlow SD, Matanoski GM. The association between weight, physical activity, and stress and variation in the length of the menstrual cycle. *Am J Epidemiol* 1991;133(1):38–49.
64. Reilly T. The Menstrual Cycle and Human Performance: An Overview. *Biological Rhythm Research* 2000;31(1):29–40.
65. Tourville TW, Shultz SJ, Vacek PM, et al. Evaluation of an Algorithm to Predict Menstrual-Cycle Phase at the Time of Injury. *J Athl Train* 2016;51(1):47–56. doi:10.4085/1062-6050-51.3.01 [published Online First: 25 January 2016].
66. Brukner P, Khan K. CLINICAL SPORTS MEDICINE. Sydney: McGraw-Hill Education 2017.
67. MacMillan C, Olivier B, Viljoen C, et al. The Association Between Menstrual Cycle Phase, Menstrual Irregularities, Contraceptive Use and Musculoskeletal Injury Among Female Athletes: A Scoping Review. *Sports Med* 2024;54(10):2515–30. doi:10.1007/s40279-024-02074-5 [published Online First: 31 August 2024].
68. Shultz SJ, Wideman L, Montgomery MM, et al. Changes in serum collagen markers, IGF-I, and knee joint laxity across the menstrual cycle. *J Orthop Res* 2012;30(9):1405–12. doi:10.1002/jor.22093 [published Online First: 2 March 2012].
69. Iwańska D, Kęska A, Dadura E, et al. The effect of the menstrual cycle on collagen metabolism, growth hormones and strength in young physically active women. *Biol Sport* 2021;38(4):721–28. doi:10.5114/biolsport.2021.107314 [published Online First: 30 June 2021].
70. Spangenburg EE, Geiger PC, Leinwand LA, et al. Regulation of physiological and metabolic function of muscle by female sex steroids. *Med Sci Sports Exerc* 2012;44(9):1653–62.

Figures




Figure 1

Schematic Hormonal Course of an Idealised 28-Day Cycle. Solid line: Oestrogen. Dashed line: Progesterone. Adapted from McNulty et al. [22]




Figure 2

PRISMA Flow Diagram of the Literature Search




Figure 3

Forest Plot of the Meta-Analysis Results




Figure 4

Variety in Menstrual Cycle Classifications Across Included Studies and Scientific Literature. The first three studies were included in the meta-analysis. *Classification scheme for the statistical analysis of this review. **Classification scheme of McNulty et al. [22]

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- PRISMAChecklist.pdf