

**Supplementary Table 1**

Correlation between normalized CD47-L - CD47-S and clinicopathological characteristics  
(Part A)

| Clinicopathological Feature  | Number | Dominant isoform |        | p value        |
|------------------------------|--------|------------------|--------|----------------|
|                              |        | CD47-L           | CD47-S |                |
| <b>FIGO Staging</b>          |        |                  |        |                |
| I - II                       | 49     | 34               | 15     | 0.7117         |
| III-IV                       | 136    | 99               | 37     |                |
| <b>Relapse</b>               |        |                  |        |                |
| Yes                          | 85     | 59               | 26     | 0.6218         |
| No                           | 40     | 26               | 14     |                |
| <b>Platinum status</b>       |        |                  |        |                |
| Sensitive                    | 46     | 34               | 12     | 0.3844         |
| Resistance                   | 13     | 8                | 5      |                |
| <b>ER</b>                    |        |                  |        |                |
| ++/+++                       | 27     | 19               | 8      | 0.9466         |
| -/+                          | 45     | 32               | 13     |                |
| <b>PR</b>                    |        |                  |        |                |
| ++/+++                       | 36     | 26               | 10     | 0.3173         |
| -/+                          | 36     | 22               | 14     |                |
| <b>p53</b>                   |        |                  |        |                |
| ++/+++                       | 28     | 23               | 5      | <b>0.0291*</b> |
| -/+                          | 42     | 24               | 18     |                |
| <b>Lymph node metastasis</b> |        |                  |        |                |
| Present                      | 28     | 21               | 7      | 0.3702         |
| Absent                       | 79     | 52               | 27     |                |
| <b>Omentum metastasis</b>    |        |                  |        |                |
| Present                      | 116    | 82               | 34     | 0.8934         |
| Absent                       | 53     | 38               | 15     |                |

The chi-square test was used to analyze the differences in clinical characteristics. \*p < 0.05  
CD47-L (Normalized CD47-L- CD47-S>-0.1), CD47-S (Normalized CD47-L- CD47-S<-0.1)

**Supplementary Table 2**

Correlation between normalized CD47-L - CD47-S and clinicopathological characteristics  
(Part B)

| Clinicopathological Feature | Dominant isoform |        | p value        |
|-----------------------------|------------------|--------|----------------|
|                             | CD47-L           | CD47-S |                |
| <b>Age</b>                  |                  |        |                |
| Mean                        | 54.73            | 54.21  | 0.7615         |
| Counts                      | 138              | 57     |                |
| <b>CA125 (U/ml)</b>         |                  |        |                |
| Mean                        | 1318             | 984.2  | 0.1564         |
| Counts                      | 129              | 54     |                |
| <b>CA199 (U/ml)</b>         |                  |        |                |
| Mean                        | 51.14            | 34.36  | 0.4675         |
| Counts                      | 55               | 27     |                |
| <b>hCG (IU/L)</b>           |                  |        |                |
| Mean                        | 2.510            | 1.607  | 0.3334         |
| Counts                      | 90               | 38     |                |
| <b>Ferritin (ng/ml)</b>     |                  |        |                |
| Mean                        | 162.8            | 325.2  | <b>0.0257*</b> |
| Counts                      | 41               | 14     |                |
| <b>Ascites volume (ml)</b>  |                  |        |                |
| Mean                        | 2034             | 1397   | <b>0.0419*</b> |
| Counts                      | 108              | 46     |                |

The unpaired, student's t-test was used to analyze the differences in clinical characteristics.

\*p < 0.05

CD47-L (Normalized CD47-L - CD47-S > -0.1), CD47-S (Normalized CD47-L - CD47-S < -0.1)