

Supplementary Materials: Investigation of the significance and role of androgen receptor expression in basal/squamous type urothelial carcinoma cell line J82 cells

Cellular Oncology

Ryotaro Sakai^{a,d}, Hiroki Shimada^{a,b,d}, Shuko Hata^a, Atsushi Yokoyama^b and Yasuhiro Nakamura^{a,c*}




Figure S1. Uncropped scans of western blots used in Figure. (A) Uncropped scans of Figure 1. (B) Uncropped scans of Figure 2. (C) Uncropped scans of Figure 4. Molecular weights of marker proteins are indicated on the left side.

^a Division of Pathology, Faculty of Medicine, Tohoku Medical and Pharmaceutical University, 1-15-1 Fukumuro, Miyagino-ku, Sendai, Miyagi 983–8536, Japan; 22023050@is.tohoku-mpu.ac.jp (R.S); shimadahis@tohoku-mpu.ac.jp (H.S); hatashu@tohoku-mpu.ac.jp (S. H.)

^b Department of Molecular Endocrinology, Tohoku University Graduate School of Medicine, 2-1 Seiryō-machi, Aoba-ku, Sendai, Miyagi, 980-8575, Japan; atsushi.yokoyama.c6@tohoku.ac.jp (A.Y)

^c Division of Pathology, Tohoku Medical and Pharmaceutical University Hospital, 1-12-1 Fukumuro, Miyagino-ku, Sendai, Miyagi 983–8512, Japan;

^d Contributed equally to this work.

*Correspondence: yasu-naka@tohoku-mpu.ac.jp (Y.N.)