

**The prevalence and development of anemia in community-acquired pneumonia patients and its association with the duration of hospitalization and mortality: questionnaire.**

<b>Patient Name</b>	<b>Patient sex</b>	<b>Patient age</b>
<b>Smoking</b> Smoking type: <ul style="list-style-type: none"> <li>- Cigarettes</li> <li>- Hookah</li> </ul> Smoking duration: .....packet/year	<b>Comorbidities:</b> Diabetes Meletus's Pulmonary Diseases <ul style="list-style-type: none"> <li>- Asthma</li> <li>- COPD</li> <li>- Lung Fibrosis</li> </ul> Hypertension \ Chronic Heart Failure Liver Diseases Kidney Diseases Tumors	
<b>Antibiotic Use Prior to Admission</b>	<b>Symptoms</b> Cough Dyspnea Fever Chest Pain Hemoptysis	
<b>Pneumonia Severity Index (PSI)</b> <ul style="list-style-type: none"> <li>- Points less than 70 (I)</li> <li>- Points 70 to 90 (II)</li> <li>- Points more than 91 (III – IV)</li> </ul>		
<b>Hemoglobin values</b> <ul style="list-style-type: none"> <li>- Day one</li> <li>- Day four</li> <li>- Day seven</li> <li>- Discharge day</li> </ul>	<b>Anemia severity</b> <ul style="list-style-type: none"> <li>- No Anemia</li> <li>- Mild Anemia</li> <li>- Moderate Anemia</li> <li>- Severe Anemia</li> <li>- Life Threatening Anemia</li> </ul>	
<b>In Hospital Complications</b> <ul style="list-style-type: none"> <li>- Treatment Failure</li> <li>- Death</li> <li>- ICU Admission</li> <li>- Septic Shock</li> <li>- Blood Transfusion</li> <li>- Mechanical Ventilation</li> </ul>	<b>Length of Hospital Stay (LOHS)</b>	