

Effects of Multifrequency Electromagnetic Pulses on tumor growth in immunocompetent mice

Roberta Piredda¹, Luis G Rodríguez Martínez², Jorge Martinez-Ortega¹,
Alejandro López Ferráz⁴, Sandra Villatoro-Gómez³, Elena Martín-García³,
María Laura García-Bermejo³, José M Almendral¹, Konstantinos
Stamatakis^{1,3, *}, and Yolanda Revilla^{1, *}

1. *Centro de Biología Molecular Severo Ochoa (CSIC-UAM), Universidad Autónoma de Madrid, Cantoblanco, Madrid, 28049, Spain.*
2. *Paso Alto Biotechnology INC, FL, EE.UU*
3. *Instituto Ramón y Cajal de Investigación Sanitaria (IRYCIS), Madrid, Spain.*
4. *Universidad de La Laguna, Santa Cruz de Tenerife, Spain.*

*Corresponding authors: Yolanda Revilla and Konstantinos Stamatakis. Centro de Biología Molecular "Severo Ochoa" (CSIC-UAM). Universidad Autónoma de Madrid. 28049 Cantoblanco, Madrid. Spain. Phone: 34-91-1964559/4570. E-mail: yrevilla@cbm.csic.es, k.s@csic.es.

Table S3. Immunohistochemical (IHC) staining for CD4, CD8, and CD68.

		CD4+ cells	CD8+ cells	CD68+ cells
Control	c0r0	++	++	++
	c1r0	++	+	++++
	c1r1	+	++	+++
MEMP	c1r4	++	+++	+++
	c0r2	++	++++	++++
	c0r1	++++	+++	++++

Staining was semi-quantitatively scored on a four-tiered scale by three independent, blinded observers. The consensus score for each sample is shown.