

Multilayered validation of graph attention autoencoder–derived drug repurposing candidates for COVID-19

Yusuke Nakayama

yusuke.nakayama@gexval.com

GEXVal Inc.

Shingo Tsuji

The University of Tokyo

Koji Yamamoto

Independent Researcher

Juran Kato-Suzuki

GEXVal Inc.

Kouichi Hosomi

Kindai University

Research Article

Keywords: drug repurposing; drug repositioning; COVID-19; SARS-CoV-2; graph attention autoencoder; knowledge graph; disproportionality analysis; real-world evidence; gene expression; hypothesis generation

Posted Date: July 2nd, 2026

DOI: <https://doi.org/10.21203/rs.3.rs-8274293/v2>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Additional Declarations: The authors declare potential competing interests as follows: J.K.-S. is the founder and representative director of GEXVal Inc. Y.N. is an employee of GEXVal Inc. S.T., K.Y. and K.H. are external consultants providing scientific and technical advice under service agreements with GEXVal Inc. for this academic study. The algorithms and methodologies reported in this study will be utilized by GEXVal Inc. for internal research and development activities and external collaborations.

Version of Record: A version of this preprint was published at Pharmaceuticals on September 25th, 2026. See the published version at <https://doi.org/10.3390/ph19101522>.

1 ***Title and authorship information***

2 ***Manuscript title:***

3 Multilayered validation of graph attention autoencoder-derived drug repurposing
4 candidates for COVID-19

5

6 ***Full author names:***

7 Yusuke Nakayama* † ¹, Shingo Tsuji † ², Koji Yamamoto³, Juran Kato-Suzuki¹, and
8 Kouichi Hosomi⁴

9

10 ***Full institutional mailing addresses:***

11 1: GEXVal Inc., Shonan iPark 2-26-1 Muraoka Higashi, Fujisawa-shi, Kanagawa, 251-
12 8555, Japan

13 2: Research Center for Advanced Science and Technology (RCAST), The University of
14 Tokyo, 4-6-1 Komaba, Meguro-ku, Tokyo, 153-8904, Japan.

15 3: Independent Researcher, Ibaraki, Japan

16 4: Faculty of Pharmacy, Kindai University, 3-4-1, Kowakae, Higashi-osaka, Osaka 577-
17 8502, Japan.

18

19 ***Email addresses:***

20 Yusuke Nakayama: yusuke.nakayama@gexval.com

21

22 ***Affiliations:***

23 1: GEXVal Inc.

24 2: Research Center for Advanced Science and Technology (RCAST), The University of
25 Tokyo

26 3: Independent Researcher

27 4: Faculty of Pharmacy, Kindai University

28 *: Corresponding author

29 † : Contributed equally

30

31

32 ***Abstract***

33 **Background/Objectives:** The COVID-19 pandemic highlighted the critical need for drug
34 repurposing to rapidly identify therapeutic options. While computational graph-based
35 approaches show promise, conventional single analytical methods often fail to capture
36 the complex pathological mechanisms needed for clinical translation. We aimed to
37 prioritize COVID-19 repurposing candidates by combining computational prediction
38 with orthogonal validation.

39 **Methods:** We developed a Graph Attention Autoencoder (GATE) framework with
40 multilayered validation integrating computational prediction, real-world data, and
41 phenome analyses. Network-based embeddings identified candidates through latent-
42 space similarity within biomedical knowledge graphs. Clinical relevance was assessed
43 by disproportionality analysis (DPA) of an adverse event reporting system, and
44 biological plausibility through gene expression and pathway profiling.

45 **Results:** Our GATE framework identified 17 drug candidates, including 13 already used
46 or recognized in the context of COVID-19 or its symptoms, consistent with the model
47 recovering known drug–disease associations. Four candidates with minimal prior
48 COVID-19 associations were identified (cilastatin, megestrol, drotrecogin alfa, and
49 ethacrynic acid). DPA prioritized two with significant inverse reporting associations for
50 COVID-19-related adverse event terms: cilastatin (reporting odds ratio [ROR]: 0.19,
51 95% confidence interval [CI]: 0.09–0.43) and megestrol (ROR: 0.63, 95% CI: 0.42–0.94).
52 Pathway profiling indicated that both drugs share molecular signatures with drugs
53 investigated in COVID-19 clinical trials; gene expression analyses suggest cilastatin

54 may exert anti-viral and anti-inflammatory effects through DPEP1 inhibition.

55 **Conclusions:** By integrating GATE-based prediction, real-world evidence, and phenome-
56 level pathway profiling, this study enabled multidimensional candidate validation that
57 single analytical methods do not provide. Prioritizing clinically approved drugs with
58 established safety profiles may facilitate accelerated clinical evaluation, and the
59 analyses provide mechanistic working hypotheses to streamline preclinical validation.
60 Although experimental confirmation remains essential, the framework offers a
61 generalizable strategy for rapid candidate identification during pandemics and diseases
62 with unmet therapeutic needs.

63

64

65 **Keywords:** drug repurposing; drug repositioning; COVID-19; SARS-CoV-2; graph
66 attention autoencoder; knowledge graph; disproportionality analysis; real-world
67 evidence; gene expression; hypothesis generation

68

69

70 ***1. Introduction***

71 The spread of infectious diseases and the emergence of new pathologies have
72 particularly increased the demand for treatments targeting rare diseases. The COVID-
73 19 pandemic highlighted critical gaps in rapid therapeutic development, demonstrating
74 the urgent need for drug repurposing strategies [1]. While traditional drug development
75 requires over 10 years and billions of dollars and enormous time commitments [2-4],
76 with approximately 10% of success rates in clinical development [5], drug repurposing
77 offers accelerated timelines and higher success probabilities due to established safety
78 and pharmacological profiles [6, 7]. Indeed, several computational approaches have
79 successfully identified COVID-19 therapeutics, including baricitinib through network-
80 based repositioning [8], validating the potential of systematic computational drug

81 discovery.

82 Computational drug repurposing leverages public databases and bioinformatics to
83 propose novel therapeutic candidates [6]. Graph structure-based approaches – which
84 model relationships between diseases, drugs, and genes as interconnected networks –
85 have emerged as powerful tools for capturing this biological complexity. Ritonavir (a
86 component of Paxlovid) and atorvastatin were successfully identified as potential
87 COVID-19 treatments through network-derived relationship analysis, highlighting how
88 graph-based representations can serve as powerful foundation for systematically
89 uncovering repurposing opportunities [9].

90 In graph structure-based drug repurposing, it is common to convert graph nodes and
91 edges into vector representations through embedding. Graph embedding methods have
92 been applied to recommendation systems in web stores and other applications [10].
93 Building upon these foundations, Graph Neural Networks (GNNs), which employ
94 neural networks for graph embedding, have been proposed [11] and have become widely
95 used.

96 While various GNN methods exist, Graph Attention Networks (GATs) that utilize
97 attention mechanisms have been widely adopted [12]. The attention mechanism is a
98 technology used in Transformer Architecture, which forms the foundation of generative
99 AI using large language models such as ChatGPT, and supports today's revolutionary
100 AI evolution [13]. Recently, these GATs have increasingly been applied to drug
101 repurposing research [14]. However, existing GNN approaches face critical limitations
102 when confronting complex pathological mechanisms; moreover, predictions from any
103 single computational method may be insufficient on their own and require
104 complementary, orthogonal validation before they can guide therapeutic development.

105 In fact, COVID-19 pathogenesis involves complex, multifaceted mechanisms
106 encompassing viral replication, host immune responses, and cellular metabolic

107 processes. Approximately 200 existing drugs may target SARS-CoV-2-associated
108 pathways, spanning diverse targets from viral proteins (spike protein, papain-like
109 protease (PLpro), 3-chymotrypsin-like protease (3CLpro), RNA-dependent RNA
110 polymerase (RdRp), and helicase [15]) and host cellular machinery (mTOR-PI3K
111 pathway [16], autophagy [17], vesicular transport systems [18], and immune response
112 [19]). This molecular complexity presents a fundamental challenge; single analytical
113 approaches may be insufficient for adequate exploration of therapeutic candidates.
114 To address this challenge, we developed a comprehensive multilayered analytical
115 framework that integrates a Graph Attention Autoencoder (GATE) with validation
116 using real-world evidence and phenome data. Rather than relying on a single line of
117 evidence, the framework triangulates each candidate across independent,
118 heterogeneous data sources. This integrated strategy prioritized candidate drugs that
119 have received little prior attention as COVID-19 therapeutics, such as cilastatin and
120 megestrol, and generated actionable mechanistic hypotheses for therapeutic
121 development.

122


123

124 ***2. Results***

125 **2.1. GATE-based identification of candidate drugs for COVID-19**

126 To systematically identify and prioritize novel drug repurposing candidates, we
127 developed a comprehensive multilayered analytical framework that complements
128 single-method computational prediction with orthogonal validation. Our methodology
129 consists of three integrated phases: (1) computational identification of candidate drugs
130 through a graph attention autoencoder (GATE) utilizing attention mechanisms and
131 data obtained from Open Targets [20], (2) assessment of potential clinical relevance
132 using real-world evidence, and (3) molecular biological assessment for mechanism of
133 action (MOA) hypothesis generation using phenome data (Figure 1). This multilayered

134 evaluation strategy integrates complementary lines of evidence, providing more
 135 actionable insights for therapeutic development than network-based prediction alone.
 136 To demonstrate the utility of this framework, we applied it to identify and prioritize
 137 novel drug repurposing candidates for COVID-19.
 138



139 **Figure 1.** Workflow of analytical framework for COVID-19
 140 AI network analysis using a graph attention autoencoder (GATE) selects candidate
 141 drugs for COVID-19, real-world data analysis (disproportionality signal detection)
 142 narrows them to drugs with inverse reporting signals for COVID-19-related AE terms,
 143 and phenome analysis (pathway profiling) evaluates their molecular plausibility and
 144 mechanisms of action.
 145
 146 Based on the latent space representation of GATE, we identified drug nodes showing
 147 high similarity to the COVID-19 node (Table 1). In interpreting the results, we
 148 considered different salt or derivative forms of the same compound (e.g., cilastatin and
 149 cilastatin sodium) as representing a single drug entity. Among the 17 drugs identified,
 150 13 are already used in the clinical management of COVID-19 or its symptoms, or are
 151 otherwise recognized in the COVID-19 literature, consistent with the model recovering
 152 established, low-novelty drug–disease relationships. The remaining 4 drugs (cilastatin,

153 megestrol, drotrecogin alfa, ethacrynic acid) had minimal prior association with COVID-
154 19, representing higher-novelty candidates that have received little prior attention as
155 COVID-19 therapeutics. Notably, these candidate drugs and COVID-19 were not
156 directly connected in the original knowledge graph (Figure 2, Figure S1-3), consistent
157 with the embedding capturing latent associations that are not represented by direct
158 edges.


160 **Table 1.** Top-ranked drug candidates based on embedding similarity to COVID-19 in the latent space learned by GATE model

ID	Name	Description	Cosine Similarity	OpenTargets Association Score	OpenTargets Rank of Score
CHEMBL1201057	CILASTATIN SODIUM	Antibiotic, Dipeptidyl Peptidase I Inhibitor	0.86067	0.01	2672
CHEMBL766	CILASTATIN	Antibiotic, Dipeptidyl Peptidase I Inhibitor	0.86006	0.01	2672
CHEMBL1201014	PREDNISOLONE SODIUM PHOSPHATE	Steroid Drug	0.85693	0.609	4
CHEMBL1201231	PREDNISOLONE PHOSPHORIC ACID	Steroid Drug	0.85526	0.609	4
CHEMBL679	EPINEPHRINE	Adrenergic Receptor Agonist	0.85111	0.372	169
CHEMBL2134724	IPRATROPIUM BROMIDE	Anticholinergic Drug, COPD and Bronchial Asthma Treatment	0.84696	0.076	737
CHEMBL1621597	IPRATROPIUM	Anticholinergic Drug, COPD and Bronchial Asthma Treatment	0.84621	0.076	737
CHEMBL1200689	NITRIC OXIDE	Nitric Oxide, Pulmonary Hypertension Treatment	0.84535	0.345	397
CHEMBL1201335	GLYCOPYRRONIUM	Anticholinergic Drug, COPD	0.84439	0.076	737
CHEMBL1437	NOREPINEPHRINE	Adrenergic Receptor Agonist	0.84266	0.372	169
CHEMBL1215	PHENYLEPHRINE	Adrenergic Receptor Agonist	0.84121	0.372	169
CHEMBL2108429	MEPOLIZUMAB	Anti-IL-5 Antibody	0.83772	0.068	799
CHEMBL1201139	MEGESTROL ACETATE	Antineoplastic Drug, Contraceptive, Progesterone Receptor Agonist	0.83667	0.086	636
CHEMBL1370	BUDESONIDE	Steroid Drug	0.83639	0.609	4
CHEMBL1364144	METHYLPREDNISOLONE ACETATE	Steroid Drug	0.83551	0.609	4
CHEMBL1201027	GLYCOPYRROLATE	Anticholinergic Drug	0.83493	0.076	737
CHEMBL2109065	DROTRECIGIN ALFA (ACTIVATED)	Antithrombotic, Anti-inflammatory, Fibrinolysis-promoting Human Activated Protein C	0.83448	0.026	1339
CHEMBL52440	DEXTROMETHORPHAN	Cough Suppressant and Expectorant (Medicon)	0.83335	0.365	334
CHEMBL1473	FLUTICASONE PROPIONATE	Steroid Drug	0.83256	0.609	4
CHEMBL3707243	GLYCOPYRRONIUM TOSYLATE	Anticholinergic Drug, Primary Axillary Hyperhidrosis Treatment	0.83239	0.076	737
CHEMBL456	ETHACRYNIC ACID	Loop Diuretic Drug	0.83171	0.004	4822

161 Drug candidates are ranked by cosine similarity scores in GATE latent space. Rows shown in bold indicate drugs not previously used

162 for COVID-19 treatment, representing candidates with minimal prior COVID-19 associations. The OpenTargets association scores

163 range from 0 to 1, with higher values indicating stronger relevance.



165 **Figure 2.** Subnetwork connecting COVID-19 and cilastatin in multimodal knowledge
166 graph


167 Subnetwork extracted from a multimodal knowledge graph consisting of drugs, diseases,
168 and genes. Nodes are colored by type: orange for diseases, blue for drugs, and green for
169 genes.

170

171 To further evaluate the findings, we investigated the association scores between
172 COVID-19 and the target genes of each candidate drug using the Open Targets Platform,
173 an integrated drug discovery database. Association scores are comprehensive
174 relatedness indicators based on literature information, genetic evidence, experimental
175 data, and other sources. This analysis revealed that all four candidate drugs showed
176 low association scores with COVID-19 (Table 1), suggesting that these drugs would not
177 be visible through existing knowledge bases. In particular, our top candidate, cilastatin,
178 had an association score of 0.01, which is markedly lower than scores (0.068–0.609) of
179 drugs that are known to have effects on COVID-19.

180 **2.2. Disproportionality analysis of candidate drugs in FAERS**

181 We further analyzed the four drugs identified by the GATE approach using real-world
 182 adverse event (AE) data from the FDA Adverse Event Reporting System (FAERS) to
 183 examine potential associations with COVID-19-related AE terms by disproportionality
 184 analysis (DPA). In this DPA, the reporting odds ratio (ROR) and its 95% confidence
 185 interval (CI) quantify whether COVID-19-related AE terms are reported
 186 disproportionately more or less often for a given drug relative to all other drugs in the
 187 database; an ROR < 1 indicates an inverse reporting association, and a 95% CI upper
 188 limit < 1 indicates that this association is statistically significant at the reporting level.
 189 Both cilastatin and megestrol showed significant inverse reporting associations for
 190 COVID-19-related AE terms, with RORs of 0.19 (95% CI: 0.09–0.43) and 0.63 (95% CI:
 191 0.42–0.94), respectively (Figure 3).
 192



193 **Figure 3.** DPA of repurposing candidate drugs and coronavirus infections
 194 (A) Reporting odds ratios (ROR) and 95% confidence intervals (CI) for candidate drugs
 195 through GATE-based analysis. “NA” indicates values that could not be calculated due
 196 to insufficient sample size. (B) Forest plot visualizing RORs for the same set of drugs.
 197 The horizontal bars represent 95% CIs. RORs for drotrecogin alfa and ethacrynic acid
 198 were not calculable and are indicated with asterisks (*).
 199

200 To examine whether the observed inverse associations were specific to each drug, we
201 conducted DPA for drugs in the same class as each candidate drug or for drugs that are
202 often co-administered with the candidate drug. Cilastatin is often co-administered with
203 imipenem, a β -lactamase inhibitor. DPA for β -lactamase inhibitors showed ROR values
204 around 1.0, and the remarkable inverse associations observed with cilastatin were not
205 observed (Figure S4). Megestrol is a progesterone receptor agonist. Progesterone
206 preparations classified as progesterone receptor agonists showed RORs around 0.5 with
207 95% CI upper limit less than 1, indicating inverse association trends similar to
208 megestrol (Figure S5).

209 Note that drotrecogin alfa and ethacrynic acid could not be evaluated using this DPA
210 due to insufficient co-occurrence reports with COVID-19-related AEs in the FAERS
211 database during the analysis period.

212 **2.3. Pathway-profile concordance with known COVID-19 candidates (LINCS 2020)**

213 To further explore these findings at the phenome level, we conducted correlation
214 analysis using pathway profiling with the Library of Integrated Network-Based
215 Cellular Signatures (LINCS) 2020 for cilastatin and megestrol, the two drugs that
216 showed statistically significant inverse associations in DPA. LINCS 2020 is the
217 successor database to the Connectivity Map (CMAP), an established platform for
218 comparing phenomic signatures of any given drug with those associated with specific
219 diseases or disease therapeutics [21]. For each drug, we explored compounds with
220 correlated pathway profiles from 17,170 compounds (241,597 profiles) using Fisher's
221 exact test.

222 The compounds showing the strongest correlations with cilastatin (Table 2) included
223 artesunate and sirolimus, which are drugs that have undergone clinical trials for
224 COVID-19. The analysis also revealed ipidacrine, ursolic acid, wortmannin, and
225 thapsigargin, which have been reported in the literature as COVID-19 therapeutic
226 candidates.

227

228 **Table 2.** Top-ranked compounds positively correlated with the pathway profile of cilastatin

PertID	Name	MOA	Cell	Conc	Time	Adj. P-value	COVID-19 relation
BRD-K79759031	artesanate		HT29	10uM	6h	4.4E-42	10 clinical studies
BRD-K66896231	ipidacrine	Acetylcholinesterase inhibitor	HT29	10uM	6h	1.0E-40	1 report
BRD-K60067222	BRD-K60067222		PC3	10uM	24h	8.7E-40	
BRD-K68185022	ursolic-acid	11-beta hydroxysteroid dehydrogenase inhibitor	PC3	70uM	24h	4.2E-39	4 reports
BRD-A19037878	BRD-A19037878		K562	2.5uM	24h	8.0E-39	
BRD-A75409952	wortmannin	PI3K inhibitor	PC3	10uM	6h	2.6E-38	2 reports
BRD-K81855038	roxatidine		HT29	10uM	6h	4.9E-38	
BRD-A79768653	sirolimus	MTOR inhibitor	PC3	3.33uM	24h	1.5E-37	9 clinical studies
BRD-K69023402	thapsigargin	ATPase inhibitor	JURKAT	0.37uM	24h	1.5E-37	3 reports
BRD-K56032964	AP-26113		HT29	0.37uM	24h	1.6E-37	

229 The "COVID-19 relation" column indicates whether each compound has been previously associated with COVID-19 through
230 publications or clinical trials. Details of the corresponding reports and studies are provided in Supporting Information 1 and 2.

231 Similarly, the compounds showing the strongest correlations with megestrol (Table 3)
232 included tianeptine, fedratinib (TG-101348), AZD-6482 and vorinostat. Tianeptine and
233 fedratinib (TG-101348) have undergone clinical trials for COVID-19. AZD-6482 and
234 vorinostat have been reported as COVID-19 therapeutic candidates.

235

236 **Table 3.** Top-ranked compounds positively correlated with the pathway profile of megestrol


PertID	Name	MOA	Cell	Conc	Time	Adj. P-value	COVID-19 relation
BRD-K37142460	MI-2		OCILY3	10uM	24h	1.3E-08	
BRD-A81370665	BI-D1870	Ribosomal protein inhibitor	HBL1	10uM	24h	1.3E-08	
BRD-K01436366	XMD-1150	Leucine rich repeat kinase inhibitor	BJAB	10uM	24h	3.4E-08	
BRD-K36363294	I-BET-151		HIMG002	2.5uM	24h	3.4E-08	
BRD-A53077924	tianeptine	Selective serotonin reuptake enhancer	HIMG001	15uM	24h	4.1E-08	1 clinical study
BRD-K12502280	TG-101348	FLT3 inhibitor	OCILY3	10uM	4h	2.5E-07	1 clinical study
BRD-K58772419	AZD-6482	PI3K inhibitor	MCF7	1.11uM	24h	3.0E-07	1 report
BRD-K12502280	TG-101348	FLT3 inhibitor	TMD8	2.5uM	4h	5.7E-07	1 clinical study
BRD-K81418486	vorinostat	HDAC inhibitor	JURKAT	10uM	24h	5.7E-07	3 reports
BRD-K54606188	BRD-K54606188		TMD8	0.66uM	4h	1.9E-06	

237 The "COVID-19 relation" column indicates whether each compound has been previously associated with COVID-19 through
238 publications or clinical trials. Details of the corresponding reports and studies are provided in Supporting Information 1 and 2.

239 **2.4. Pathway alterations following cilastatin treatment**

240 In the LINCS 2020 data, cilastatin treatment significantly decreased the expression of
241 42 pathways in the HT29 cell line (Table S2). Enrichment Map analysis revealed that
242 gene sets related to splicing, ribosome, mitochondria, telomere, and ubiquitin pathways
243 were altered (Figure 4).

244



245 **Figure 4.** Enrichment map of down-regulated pathways following cilastatin treatment
246 This network visualizes gene ontology (GO) biological process terms significantly down-
247 regulated by cilastatin treatment. Each node represents a gene set corresponding to a
248 GO term, with node size proportional to the number of genes in the set and node color
249 reflecting statistical significance (log₁₀-transformed p-value). Edges indicate gene
250 overlap between gene sets. Functionally related clusters are annotated.

251

252

253 **3. Discussion**

254 **3.1. Recovery of latent drug–disease associations by the GATE approach**

255 To assess whether these candidates had been overlooked in prior repurposing efforts,
256 we examined whether the four drugs had been reported as COVID-19 repurposing

257 candidates—and, where reported, investigated in depth—in existing studies. Cilastatin
258 was reported as one of 78 candidates for SARS-CoV-2 3CLpro inhibition through docking
259 simulation approaches, but detailed validation was not performed [22]. Megestrol was
260 proposed as one of 34 COVID-19 therapeutic candidates through network-based
261 approaches, though detailed investigation was not conducted [23]. Drotrecogin alfa had
262 no existing reports; to our knowledge, this is the first report of this drug as a COVID-19
263 candidate. Ethacrynic acid was reported as a promising candidate for SARS-CoV-2 main
264 protease (Mpro) inhibition through docking simulation approaches, with experimental
265 validation also performed [24].

266 Although some candidate drugs identified by the GATE approach have been previously
267 reported by other computational methods, those studies provided only preliminary
268 identification without comprehensive evaluation of therapeutic potential. Notably, three
269 of the four candidates (cilastatin, drotrecogin alfa, and ethacrynic acid) were not
270 identified by the conventional network-based approaches we surveyed. This suggests
271 that GATE may identify complex associations that remain undetected by conventional
272 network analysis, exemplified by drotrecogin alfa.

273 These results suggest that the GATE approach is complementary to existing methods
274 and has the potential to efficiently discover novel relationships between drug–disease
275 pairs. Furthermore, the identification of candidates such as ethacrynic acid, for which
276 experimental validation has been conducted, provides additional support for the
277 plausibility of the prioritized candidates. However, these remain computational
278 predictions that require experimental validation.

279 **3.2. Potential clinical relevance of candidate drugs from FAERS disproportionality** 280 **analysis**

281 For cilastatin, COVID-19-related AE terms were reported approximately 5-fold less
282 frequently among cilastatin reports than among non-cilastatin reports, and for
283 megestrol approximately 40% less frequently. Although these disproportionality signals

284 are hypothesis-generating and cannot by themselves establish a causal effect, the
285 inverse association for cilastatin was among the largest observed in this analysis,
286 suggesting that both drugs may have potential worth investigating as COVID-19
287 repurposing candidates.

288 The β -lactamase inhibitor comparison suggests that the inverse reporting association
289 observed for cilastatin is not explained by co-reported β -lactamase inhibitors and may
290 be relatively specific to cilastatin at the reporting level. For megestrol, the progesterone
291 receptor agonist comparison suggests that the observed inverse reporting association
292 reflects a pharmacological class effect mediated through progesterone receptors rather
293 than a signal specific to megestrol. However, more detailed analysis is needed to clearly
294 distinguish the influence of class effects.

295 The inverse reporting association observed for cilastatin is consistent with previously
296 proposed mechanisms by which this drug might reduce SARS-CoV-2 viral replication
297 and modulate the cytokine response observed in severe COVID-19 [25]. Based on these
298 findings, cilastatin warrants further investigation as a potential therapeutic candidate
299 for COVID-19.

300 Regarding megestrol, no literature has reported direct associations with COVID-19.
301 However, progesterone, a drug in the same class that showed a significant inverse
302 reporting association with COVID-19-related AE terms in our DPA, has anti-
303 inflammatory properties, suggesting potential as a COVID-19 treatment [26-28].
304 Clinical studies on COVID-19 treatment using progesterone have been conducted
305 (NCT04365127, NCT04865029) and in the NCT04865029 trial, progesterone
306 administration in moderate to severe COVID-19 patients resulted in significant
307 improvement in clinical status (a score of 1.5 points, corresponding to a 3-day reduction
308 in oxygen supplementation and 2.5-day reduction in hospitalization duration) [29].
309 Given the observed class effect of progesterone receptor agonists and the clinical

310 evidence supporting progesterone, megestrol merits evaluation to determine whether it
311 shows similar effects in confirmatory studies.

312 **3.3. Phenome-level plausibility from pathway profiling**


313 These phenome-level correlations suggest that both cilastatin and megestrol may have
314 potential as COVID-19 therapeutic candidates. The correlation with established
315 COVID-19 therapeutic candidates supported the plausibility of cilastatin as a candidate
316 worth further investigation.

317 **3.4. Hypothesized mechanisms of action of cilastatin**

318 To provide mechanistic insights that could guide future preclinical studies, we sought
319 to elucidate the underlying biological mechanisms based on pathway profiling data. We
320 argued that because pathway-level information represents a functional layer closer to
321 biological responses than individual genes, a detailed analysis of the LINCS 2020 data
322 can facilitate hypothesis generation regarding mechanisms of action.

323 Based on the pathway alterations following cilastatin treatment (Figure 4), we
324 constructed a mechanistic hypothesis for how cilastatin might intervene in the
325 pathophysiological processes of COVID-19 induced by SARS-CoV-2 infection (Figure 5).

326



327 **Figure 5.** Hypothesized mechanisms of action of cilastatin in COVID-19

328 This schematic illustrates potential therapeutic mechanisms by which cilastatin may

329 alleviate COVID-19 pathology, inferred from pathway-level gene expression changes.
330 Cilastatin-mediated inhibition of DPEP1 is proposed to influence multiple downstream
331 biological processes, including ribosome biogenesis, RNA splicing, and mitochondrial
332 metabolism. These changes converge on antiviral and anti-inflammatory outcomes such
333 as suppression of viral replication and inflammation. Pathways shaded in blue indicate
334 those with down-regulated gene expression following cilastatin treatment.

335

336 Cilastatin is clinically used as a dipeptidase-1 (DPEP1) inhibitor in combination with
337 imipenem to prevent renal degradation of the antibiotic by DPEP1. Beyond this
338 established role in antibiotic protection, our analysis of LINCS 2020 data indicated that
339 cilastatin modulates multiple cellular pathways related to splicing, ribosomal function,
340 and mitochondrial processes in cells. DPEP1, in addition to its primary function of
341 dipeptide hydrolysis in the kidney, also functions as a neutrophil adhesion receptor on
342 vascular endothelial cells and is known to promote neutrophil recruitment during
343 inflammation [30]. DPEP1 may restrict host ribosomes utilized by SARS-CoV-2 by
344 controlling the PI3K/Akt/mTOR pathway [31], which regulates cellular protein
345 synthesis and metabolism [32]. This may suppress viral replication through a
346 mechanism similar to that of mTOR inhibitors such as sirolimus (rapamycin) [33].

347 Of particular interest, it is known that the nonstructural protein 16 (NSP16) of SARS-
348 CoV-2 binds to U1/U2 snRNA and interferes with the host splicing mechanism [34].
349 Suppression of splicing-related pathways may function as a cellular stress response that
350 limits host gene expression and potentially suppresses viral replication by restricting
351 protein synthesis. Moreover, splicing abnormalities may activate antiviral responses via
352 RIG-I-like receptors [35].

353 Beyond the splicing mechanism, cilastatin also affects mitochondrial pathways.
354 Decreased expression of mitochondria-related genes suggests changes in cellular

355 metabolic activity and immune signaling. These changes are expected to achieve
356 balanced immune regulation by limiting energy available for viral replication while
357 maintaining type I interferon responses through mitochondrial antiviral signaling
358 protein (MAVS) [36].

359 **3.5. Study limitations**

360 While our multilayered computational approach successfully identified several
361 promising therapeutic candidates and generated mechanistic hypotheses, there are
362 several important limitations that must be addressed.

363 *Limitations of the GATE approach:* The predictive performance of our knowledge graph-
364 based method heavily depends on the comprehensiveness and quality of the base
365 knowledge graph being used. In biomedical knowledge graphs, data bias tends to occur
366 based on research progress and attention levels, with particularly insufficient
367 information regarding rare diseases and drugs in early development stages. This
368 deficiency may limit the discovery of novel drug-disease associations and reduce
369 predictability. Therefore, developing comprehensive knowledge graphs that
370 systematically incorporate data on rare diseases and emerging therapeutics will be
371 essential for improving comprehensiveness and reducing bias in future studies.

372 *Limitations of real-world data analysis:* Since FAERS is a spontaneous AE reporting
373 system, it inherently contains problems such as reporting bias, underreporting, and
374 selective reporting. A critical limitation emerged from our analysis of drotrecogin alfa
375 and ethacrynic acid, which revealed that the method cannot analyze drugs with
376 insufficient data in the database. This particularly affects newer drugs that lack
377 extensive real-world usage data, limiting our ability to detect potential associations.
378 Conversely, this suggests that classical, widely used drugs may have higher detection
379 sensitivity due to larger volumes of available data. Furthermore, inferring causal
380 relationships from observed associations remains challenging, and the influence of
381 confounding factors cannot be completely eliminated.

382 *Limitations of phenome analysis:* The LINCS database used is primarily based on drug
383 responses in cancer cell lines, which exhibit different characteristics from alveolar
384 epithelial cells, vascular endothelial cells, and immune cells, which are important in
385 COVID-19 pathogenesis. Additionally, drug responses under the inflammatory
386 environment in patients during SARS-CoV-2 infection may differ significantly from
387 normal cell culture conditions. Furthermore, it is difficult to capture host-virus
388 interactions and other complex biological processes that can only be detected in in vivo
389 conditions using cell line-based phenome data. While extensive animal data have been
390 accumulated and could potentially be curated and integrated into such databases,
391 comprehensive incorporation remains challenging due to species differences in drug
392 target molecules, including variations in binding site homology and pharmacological
393 responses between species.

394 *Need for experimental validation:* This study primarily focused on computational
395 predictions and analysis of existing databases to systematically identify therapeutic
396 candidates and elucidate potential mechanisms. While this approach provided valuable
397 insights into drug repurposing opportunities for COVID-19 treatment, future studies
398 should include direct validation using SARS-CoV-2-infected models and/or clinical
399 evaluation.

400

401

402 **4. Materials and Methods**

403 **4.1. Graph attention autoencoder (GATE)**

404 **4.1.1. Knowledge graph construction**

405 We constructed a heterogeneous knowledge graph comprising three node types —
406 disease, drug, and gene (target) — with data primarily obtained from Open Targets
407 (version 24.06) [20], which aggregates target–disease association data from multiple
408 public sources. Four edge types were defined: (i) "Molecular interactions" (target–target),

409 (ii) "Associations — direct (overall score)" (target–disease), (iii) "Drug — mechanism of
410 action" (drug–target), and (iv) "Drug — indications" (drug–disease). The resulting
411 knowledge graph comprised 11,882 disease, 4,314 drug, and 11,448 gene nodes
412 connected by 341,911 edges.

413 Initial node attribute vectors were generated by our proprietary LLM-based feature-
414 engineering pipeline, which produces an embedding vector reflecting the characteristics
415 of each node.

416 ***4.1.2. Model architecture and training***

417 Node embeddings were learned with a graph attention autoencoder (GATE)
418 implemented in Python using the Spektral library (v1.3.1) [37] with the TensorFlow
419 2.15.0/Keras backend. The encoder applied a single graph attention layer (GATConv)
420 with 8 attention heads (concatenated; 256 channels per head) and an exponential linear
421 unit (ELU) activation, with L2 regularization (2.5×10^{-4}) applied to the kernel, attention-
422 kernel, and bias weights. The decoder reconstructed the input node features through
423 two fully connected layers (256 units with ReLU activation, followed by a linear layer
424 matching the input feature dimension). The adjacency matrix was normalized using the
425 standard GATConv preprocessing. The model was trained as a feature-reconstruction
426 autoencoder by minimizing the mean squared error between the input and
427 reconstructed node features (plus the L2 penalty), using the Adam optimizer (learning
428 rate 1×10^{-3}) in full-graph mode for approximately 4×10^4 epochs, without dropout; a fixed
429 random seed was set.

430 ***4.1.3. Candidate identification***

431 From the trained model, an attention-based latent node representation was used to rank
432 all nodes by their cosine similarity to the COVID-19 node, with higher cosine similarity
433 indicating greater proximity in the latent space. Drug nodes that fell within the top 500
434 of all ranked nodes were retained as candidates; different salt or derivative forms of the
435 same compound (e.g., cilastatin and cilastatin sodium) were treated as a single drug

436 entity.

437 **4.2. Disproportionality analysis with spontaneous reporting systems**

438 We utilized a spontaneous adverse event (AE) reporting database constructed by U.S.
439 Food and Drug Administration (FDA). The FDA Adverse Event Reporting System
440 (FAERS) contains reports of spontaneous AEs reported by consumers, pharmaceutical
441 companies, and medical professionals worldwide. FAERS consists of seven datasets:
442 DEMO (patient demographic and administrative information), DRUG (drug
443 information), REAC (AEs), INDI (indications for the use of reported drugs), THER (start
444 and end dates for the reported drugs), OUTC (outcomes for the event) and RPSR
445 (sources for the event). The present study used FAERS data from January 2020 through
446 March 2022, representing 3,617,885 reports. This data was obtained through JAPIC-
447 AERS (Japan Pharmaceutical Information Center (JAPIC), Tokyo, Japan), a
448 commercially processed FAERS database.

449 AEs in the FAERS database are coded using Medical Dictionary for Regulatory
450 Activities (MedDRA) Preferred Terms (PTs). COVID-19-related AE terms were defined
451 as the preferred terms grouped under the MedDRA High Level Term (HLT)
452 "Coronavirus infections" (code 10084510); the complete list of these 17 preferred terms
453 is provided in Table S1, and this set is referred to as "coronavirus infections" in the
454 figures.

455 AE risk was evaluated by calculating the reporting odds ratio (ROR) and corresponding
456 95% confidence intervals (CI). The ROR and 95% CI were defined as follows: $ROR = (a/c) / (b/d)$;
457 $95\% CI = \exp [\ln(ROR) \pm 1.96\sqrt{(1/a + 1/b + 1/c + 1/d)}]$, where a, b, c and d refer to
458 the number of individuals in the FAERS database divided into: (a) those who received
459 the drug of interest (e.g., cilastatin) and had COVID-19-related AE terms reported; (b)
460 those who received the drug of interest without COVID-19-related AE terms; (c) those
461 who did not receive the drug of interest and had COVID-19-related AE terms reported;
462 and (d) those who did not receive the drug of interest and did not have COVID-19-related

463 AE terms. If the upper limit of the 95% CI was < 1 , a significant inverse association was
464 assumed between use of the drug of interest and the reporting of COVID-19-related AE
465 terms.

466 **4.3. Pathway profiling**

467 To explore potential therapeutic agents at the phenome level, we conducted analyses
468 using pathway alteration profiles following drug treatment. We utilized gene expression
469 profile data from the Expanded CMAP LINCS Resource 2020 [21], which contains
470 transcriptome data generated from cell lines treated with various compounds. To
471 mitigate noise inherent in high-throughput methodologies, we employed a quality
472 control process based on gene expression distribution across replicate samples,
473 retaining only high-confidence data; for each perturbation condition, approximately
474 three replicate samples were aggregated into a single expression profile. Plate-to-plate
475 normalization was performed to eliminate assay bias. Furthermore, we applied
476 distribution analysis to determine gene expression alteration levels, selecting only
477 profiles with gene alterations at a Q-value threshold of < 0.20 . Gene Set Enrichment
478 Analysis (GSEA) [38, 39] was conducted on the filtered profiles to identify significantly
479 altered pathways for each profile, using a false discovery rate (FDR) cutoff of < 0.20 ,
480 with up- and down-regulated pathways considered separately. For the GSEA analysis,
481 we utilized ontology gene sets (C5) [40] from the Molecular Signatures Database
482 (MSigDB) [41, 42]. Correlations between pathway profiles were evaluated using Fisher's
483 exact test to identify compounds with similar phenotypic effects. Statistical analysis
484 was implemented in C and Ruby, ensuring computational efficiency for large-scale
485 comparisons. Multiple testing correction was applied at the profile level using the
486 Benjamini-Hochberg method, with an FDR threshold of 0.20 to determine statistical
487 significance. These relatively lenient thresholds were chosen given the exploratory,
488 hypothesis-generating nature of the analysis. Pathway enrichment visualization was
489 performed using Enrichment Map (v3.4.0) [43], a plugin for Cytoscape software (Version

3.10.1) [44]. Significantly enriched pathways with $FDR < 0.20$ were included in the analysis. The pathway network was constructed with each node representing an enriched gene set, and edges connecting pathways with shared genes, where the edge weight corresponds to the overlap coefficient between the gene sets. To improve interpretability of the resulting network, we utilized the AutoAnnotate plugin (v1.5.1) [45] to cluster and label groups of functionally related pathways based on their shared gene content. Clusters were manually reviewed and annotated to ensure biological relevance.

5. Conclusions

In this study, we systematically evaluated the applicability of existing drugs for COVID-19 treatment using the GATE-based latent space analysis with real-world evidence and phenome-level validation. We identified two candidate drugs for further investigation: cilastatin and megestrol, both clinically approved medications with established safety profiles, which may be amenable to accelerated clinical evaluation.

Our multilayered validation approach achieved confident candidate drug selection by comprehensively assessing clinical and molecular biological plausibility. Integration of these heterogeneous data sources – static knowledge graphs, real-world clinical observations, and dynamic gene expression data – enabled multi-dimensional insights difficult to obtain with a single analytical method. Real-world data analysis indicated potential clinical relevance at the reporting level, while phenome data analysis supported molecular biological validity and mechanistic hypotheses for both drugs.

This integrated framework offers a practical approach to computational drug discovery that bridges the gap between in silico prediction and clinical translation. This multimodal methodology has applications for rapid candidate identification during pandemics and for diseases and conditions with limited therapeutic options. Future

517 developments in large language models (LLMs) and agentic AI may further strengthen
518 this multilayered framework, including by improving knowledge-graph
519 comprehensiveness.

520 Future research should focus on in vivo validation and clinical trials of these repurposed
521 drug candidates. It should also explore the applicability of this methodology to other
522 diseases and establish selection criteria for optimal computational methods based on
523 data scale and intended use.

524

525

526 **Supplementary Materials:** The following supporting information can be downloaded:
527 SupplementalTables.pdf, containing Table S1 (the preferred terms grouped under the
528 MedDRA HLT "Coronavirus infections") and Table S2 (significantly down-regulated
529 gene sets in cilastatin-treated HT29 cells (LINCS 2020)); SupplementalFigures.pdf,
530 containing Figure S1 (subnetwork connecting COVID-19 and megestrol), Figure S2
531 (subnetwork connecting COVID-19 and drotrecogin alfa), Figure S3 (subnetwork
532 connecting COVID-19 and ethacrynic acid), Figure S4 (DPA of β -lactamase inhibitors
533 and coronavirus infections), and Figure S5 (DPA of progestins and related drug classes
534 and coronavirus infections); SupportingInformation1_COVID-19_clinicaltrials.xlsx,
535 providing details of the clinical studies corresponding to the "COVID-19 relation"
536 column in Tables 2 and 3; and SupportingInformation2_COVID-19_reports.docx,
537 providing details of the reports corresponding to the "COVID-19 relation" column in
538 Tables 2 and 3.

539

540 **Author Contributions:** Conceptualization, Y.N., S.T., J.K.-S. and K.H.; methodology, S.T.,
541 K.H. and K.Y.; data curation, S.T. and K.Y.; formal analysis, Y.N. and K.H.; investigation,
542 Y.N. and K.H.; writing—original draft preparation, Y.N.; writing—review and editing,

543 S.T., J.K.-S. and K.H.; supervision, Y.N., S.T., J.K.-S. and K.H. All authors have read
544 and agreed to the published version of the manuscript.

545 **Funding:** This research and its publication were supported by GEXVal Inc. through its
546 internal research and development budget, and by the Business Restructuring Subsidy
547 Program of the Ministry of Economy, Trade and Industry (METI), Japan (grant number
548 R2148U00245-000). The funders provided financial support in the form of salaries and
549 research expenses but had no role in the study design; in the collection, analyses, or
550 interpretation of data; in the writing of the manuscript; or in the decision to publish the
551 results.

552

553 **Institutional Review Board Statement:** Not applicable. This study used only secondary,
554 de-identified data and did not involve human participants, animals, or human tissue;
555 institutional review board approval was therefore not required. All research was
556 conducted in accordance with ICMJE, MEXT, and JSPS ethical guidelines.

557

558 **Informed Consent Statement:** Not applicable. This study did not involve human
559 participants.

560

561 **Data Availability Statement:** Publicly available datasets were analyzed in this study.
562 These data can be found at: the Open Targets Platform
563 (<https://platform.opentargets.org/downloads>), the CMAP LINCS 2020 collection on the
564 CLUE platform (<https://clue.io/data/CMap2020#LINCS2020>), and the MSigDB C5
565 collection (<https://www.gsea-msigdb.org/gsea/msigdb/>). Spontaneous adverse event data
566 were obtained from JAPIC-AERS, a commercially licensed database derived from the
567 FDA Adverse Event Reporting System (FAERS); although the underlying FAERS data
568 are publicly available from the FDA, the JAPIC-AERS data used in this study are

569 subject to licensing and commercial restrictions and are not publicly available. Derived
570 data supporting the findings of this study are provided within the article and its
571 Supplementary Materials where possible. The node features used to initialize the model
572 were generated by a proprietary LLM-based feature-engineering pipeline; full public
573 release is not possible owing to commercial and intellectual-property considerations
574 associated with this proprietary embedding-generation methodology, although these
575 node features may be made available to editors and reviewers for verification during
576 peer review.

577

578 **Code Availability:** The code for this work was developed for and is proprietary to GEXVal
579 Inc. and may be made available to editors and reviewers for scrutiny during peer review;
580 full public release is not possible owing to commercial and intellectual-property
581 considerations.

582

583 **Acknowledgments:** The authors gratefully acknowledge the valuable discussions and
584 technical advice provided by K. Matsubara, T. Morimoto, N. Hirashima, Dr. K. Shimada,
585 and T. Naito. We would like to thank K. Hashizume for collaboration on this work. We
586 also thank Dr. J. Urano for insightful comments and critical suggestions on the
587 manuscript. During the preparation of this manuscript, the authors used ChatGPT
588 (OpenAI) and Claude (Anthropic) for English-language editing and to support literature
589 searches. The authors have reviewed and edited the output and take full responsibility
590 for the content of this publication.

591

592 **Conflicts of Interest:** J.K.-S. is the founder and representative director of GEXVal Inc.
593 Y.N. is an employee of GEXVal Inc. S.T., K.Y. and K.H. are external consultants
594 providing scientific and technical advice under service agreements with GEXVal Inc. for

595 this academic study. The algorithms and methodologies reported in this study will be
596 utilized by GEXVal Inc. for internal research and development activities and external
597 collaborations. All authors have reviewed and approved this statement.

598

599

600 **References**

- 601 1. Serafin, M.B., et al., *Drug repositioning is an alternative for the treatment of coronavirus COVID-*
602 *19*. Int J Antimicrob Agents, 2020. **55**(6): p. 105969.
- 603 2. Thomas, D., et al. *Clinical Development Success Rates and Contributing Factors 2011–2020*
604 *(Biotechnology Innovation Organization, 2021)*. 2021; Available from: [https://go.bio.org/rs/490-](https://go.bio.org/rs/490-EHZ-999/images/ClinicalDevelopmentSuccessRates2011_2020.pdf)
605 [EHZ-999/images/ClinicalDevelopmentSuccessRates2011_2020.pdf](https://go.bio.org/rs/490-EHZ-999/images/ClinicalDevelopmentSuccessRates2011_2020.pdf).
- 606 3. Simoens, S. and I. Huys, *R&D Costs of New Medicines: A Landscape Analysis*. Front Med
607 (Lausanne), 2021. **8**: p. 760762.
- 608 4. Sertkaya, A., et al., *Costs of Drug Development and Research and Development Intensity in the*
609 *US, 2000-2018*. JAMA Network Open, 2024. **7**(6): p. e2415445.
- 610 5. Wong, C.H., K.W. Siah, and A.W. Lo, *Estimation of clinical trial success rates and related*
611 *parameters*. Biostatistics, 2018. **20**(2): p. 273-286.
- 612 6. Pushpakom, S., et al., *Drug repurposing: progress, challenges and recommendations*. Nat Rev
613 Drug Discov, 2019. **18**(1): p. 41-58.
- 614 7. Nosengo, N., *Can you teach old drugs new tricks?* Nature, 2016. **534**(7607): p. 314-316.
- 615 8. Saber-Ayad, M., et al., *Current Status of Baricitinib as a Repurposed Therapy for COVID-19*.
616 Pharmaceuticals (Basel), 2021. **14**(7).
- 617 9. Morselli Gysi, D., et al., *Network medicine framework for identifying drug-repurposing*
618 *opportunities for COVID-19*. Proc Natl Acad Sci U S A, 2021. **118**(19).
- 619 10. Cai, H., V.W. Zheng, and K.C.-C. Chang, *A Comprehensive Survey of Graph Embedding: Problems,*
620 *Techniques, and Applications*. IEEE Transactions on Knowledge and Data Engineering, 2018.
621 **30**(9): p. 1616-1637.
- 622 11. Cao, S., W. Lu, and Q. Xu, *Deep Neural Networks for Learning Graph Representations*.
623 Proceedings of the AAAI Conference on Artificial Intelligence, 2016. **30**(1).
- 624 12. Salehi, A. and H. Davulcu. *Graph Attention Auto-Encoders*. IEEE.
- 625 13. Vaswani, A.S., Noam; Parmar, Niki; Uszkoreit, Jakob; Jones, Llion; Gomez, Aidan N; Kaiser,
626 Łukasz; Polosukhin, Illia, *Attention Is All You Need*. Advances in Neural Information Processing
627 Systems (Curran Associates, Inc.), 2017. **30**.
- 628 14. Li, W., et al., *Drug repurposing based on the DTD-GNN graph neural network: revealing the*
629 *relationships among drugs, targets and diseases*. BMC Genomics, 2024. **25**(1).
- 630 15. Muhammed, Y., *Molecular targets for COVID-19 drug development: Enlightening Nigerians*
631 *about the pandemic and future treatment*. Biosafety and Health, 2020. **2**(4): p. 210-216.
- 632 16. Kindrachuk, J., et al., *Antiviral potential of ERK/MAPK and PI3K/AKT/mTOR signaling*
633 *modulation for Middle East respiratory syndrome coronavirus infection as identified by temporal*
634 *kinome analysis*. Antimicrob Agents Chemother, 2015. **59**(2): p. 1088-99.
- 635 17. Gassen, N.C., et al., *SKP2 attenuates autophagy through Beclin1-ubiquitination and its*
636 *inhibition reduces MERS-Coronavirus infection*. Nat Commun, 2019. **10**(1): p. 5770.
- 637 18. Sureda, A., et al., *Endoplasmic reticulum as a potential therapeutic target for covid-19 infection*
638 *management?* Eur J Pharmacol, 2020. **882**: p. 173288.
- 639 19. Han, N., et al., *Identification of SARS-CoV-2-induced pathways reveals drug repurposing*
640 *strategies*. Sci Adv, 2021. **7**(27).
- 641 20. *Open Targets Platform Datasets*. Available from: <https://platform.opentargets.org/downloads>.
- 642 21. *CMAP LINCS 2020*. Available from: <https://clue.io/data/CMap2020#LINCS2020>.
- 643 22. Wu, C., et al., *Analysis of therapeutic targets for SARS-CoV-2 and discovery of potential drugs by*
644 *computational methods*. Acta Pharmaceutica Sinica B, 2020. **10**(5): p. 766-788.
- 645 23. Zhou, Y., et al., *A network medicine approach to investigation and population-based validation of*
646 *disease manifestations and drug repurposing for COVID-19*. PLOS Biology, 2020. **18**(11): p.
647 e3000970.
- 648 24. Isgrò, C., A.M. Sardanelli, and L.L. Palese, *Systematic Search for SARS-CoV-2 Main Protease*

- Inhibitors for Drug Repurposing: Ethacrynic Acid as a Potential Drug.* Viruses, 2021. **13**(1): p. 106.
25. González-Nicolás, M., et al., *Cilastatin: a potential treatment strategy against COVID-19 that may decrease viral replication and protect from the cytokine storm.* Clin Kidney J, 2020. **13**(5): p. 903-905.
 26. Shilpa, *COVID-19 and Progesterone: Part 1. SARS-CoV-2, Progesterone and its potential clinical use.* Endocrine and Metabolic Science, 2021. **5**: p. 100109.
 27. Su, S., et al., *Modulation of innate immune response to viruses including SARS-CoV-2 by progesterone.* Signal Transduction and Targeted Therapy, 2022. **7**(1).
 28. Al-Kuraishy, H.M., et al., *New insights on the potential effect of progesterone in Covid-19: Anti-inflammatory and immunosuppressive effects.* Immunity, Inflammation and Disease, 2023. **11**(11).
 29. Ghandehari, S., et al., *Progesterone in Addition to Standard of Care vs Standard of Care Alone in the Treatment of Men Hospitalized With Moderate to Severe COVID-19: A Randomized, Controlled Pilot Trial.* Chest, 2021. **160**(1): p. 74-84.
 30. Choudhury, S.R., et al., *Dipeptidase-1 Is an Adhesion Receptor for Neutrophil Recruitment in Lungs and Liver.* Cell, 2019. **178**(5): p. 1205-1221.e17.
 31. Cui, X., et al., *DPEP1 is a direct target of miR-193a-5p and promotes hepatoblastoma progression by PI3K/Akt/mTOR pathway.* Cell Death Dis, 2019. **10**(10): p. 701.
 32. Appelberg, S., et al., *Dysregulation in Akt/mTOR/HIF-1 signaling identified by proteo-transcriptomics of SARS-CoV-2 infected cells.* Emerg Microbes Infect, 2020. **9**(1): p. 1748-1760.
 33. Khan, N., *mTOR: A possible therapeutic target against SARS-CoV-2 infection.* Arch Stem Cell Ther, 2021. **2**(1): p. 5-7.
 34. Banerjee, A.K., et al., *SARS-CoV-2 Disrupts Splicing, Translation, and Protein Trafficking to Suppress Host Defenses.* Cell, 2020. **183**(5): p. 1325-1339.e21.
 35. Chang, A.Y., et al., *Modulation of SF3B1 in the pre-mRNA spliceosome induces a RIG-I-dependent type I IFN response.* J Biol Chem, 2021. **297**(5): p. 101277.
 36. Hanada, Y., et al., *MAVS is energized by Mff which senses mitochondrial metabolism via AMPK for acute antiviral immunity.* Nature Communications, 2020. **11**(1): p. 5711.
 37. Grattarola, D. and C. Alippi *Graph Neural Networks in TensorFlow and Keras with Spektral.* 2020. arXiv:2006.12138 DOI: 10.48550/arXiv.2006.12138.
 38. Subramanian, A., et al., *Gene set enrichment analysis: A knowledge-based approach for interpreting genome-wide expression profiles.* Proceedings of the National Academy of Sciences, 2005. **102**(43): p. 15545-15550.
 39. Mootha, V.K., et al., *PGC-1 α -responsive genes involved in oxidative phosphorylation are coordinately downregulated in human diabetes.* Nature Genetics, 2003. **34**(3): p. 267-273.
 40. MSigDB: c5. Available from: <https://www.gsea-msigdb.org/gsea/msigdb/human/genesets.jsp?collection=C5>.
 41. MSigDB. Available from: <https://www.gsea-msigdb.org/gsea/msigdb/>.
 42. Liberzon, A., et al., *Molecular signatures database (MSigDB) 3.0.* Bioinformatics, 2011. **27**(12): p. 1739-1740.
 43. Merico, D., et al., *Enrichment Map: A Network-Based Method for Gene-Set Enrichment Visualization and Interpretation.* PLOS ONE, 2010. **5**(11): p. e13984.
 44. Shannon, P., et al., *Cytoscape: a software environment for integrated models of biomolecular interaction networks.* Genome Res, 2003. **13**(11): p. 2498-504.
 45. Kucera, M., et al., *AutoAnnotate: A Cytoscape app for summarizing networks with semantic annotations.* F1000Res, 2016. **5**: p. 1717.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementalFigures.pdf](#)
- [SupplementalTables.pdf](#)
- [SupportingInformation1COVID19clinicaltrials.xlsx](#)
- [SupportingInformation2COVID19reports.pdf](#)