Research Square

Preprints are preliminary reports that have not undergone peer review.
They should not be considered conclusive, used to inform clinical practice,
or referenced by the media as validated information.

Quantum Bio-Inorganic Chemistry of Lithium: Nuclear Spin Effects, Radical Pairs, and the Thermodynamic Regulation of Neural Dynamics

Artem Borozentsev

a.borozentsev05@outlook.fr

University of Toulouse-Jean Jaurès

Article

Keywords: Magnetic Isotope Effect, Radical Pair Mechanism, Lithium, GSK-3β, Alzheimer's, Epilepsy, Predictive Coding, Quantum Biology

Posted Date: March 3rd, 2026

DOI: https://doi.org/10.21203/rs.3.rs-8302041/v1

License: This work is licensed under a Creative Commons Attribution 4.0 International License.
Read Full License

Additional Declarations: No competing interests reported.

Quantum Bio-Inorganic Chemistry of Lithium: Nuclear Spin Effects, Radical Pairs, and the Thermodynamic Regulation of Neural Dynamics

Artem Borozentsev*

Independent Researcher

December 8, 2025

Abstract

Lithium remains the "dark matter" of psychopharmacology. Despite being the gold standard treatment for Bipolar Disorder and a promising neuroprotective agent for Alzheimer's disease, its molecular mechanism of action presents an enduring challenge for modern medicine. The prevailing hypothesis of "ionic mimicry"—which suggests that lithium competes with magnesium due to ionic radius similarity—has faced difficulties in quantitatively explaining therapeutic efficacy at sub-toxic concentrations. Furthermore, this classical view does not fully account for the "Isotope Anomaly," wherein Lithium-6 and Lithium-7 exhibit distinct effects on animal behavior and mitochondrial function despite being chemically identical. In this paper, I propose a comprehensive **Quantum-Thermodynamic Model** of lithium action. The simulations presented here suggest that lithium may act as a quantum modulator of the Radical Pair Mechanism (RPM), potentially regulating mitochondrial **Reactive Oxygen Species (ROS)** and magnesium-dependent enzymatic sites. By exploiting the **Magnetic Isotope Effect (MIE)** via Hyperfine Interactions (> 5 mT), lithium is hypothesized to stabilize the brain's metabolic coherence. This paper provides a mathematical derivation linking spin dynamics to reaction yields via the Haberkorn equation, analyzes the clinical spectrum through the lens of phase transitions, and integrates these findings with the Free Energy Principle. Finally, a definitive "Zero-Field" experiment with a ^{24}Mg negative control is proposed to empirically test this paradigm.

Keywords: Magnetic Isotope Effect, Radical Pair Mechanism, Lithium, GSK-3β, Alzheimer's, Epilepsy, Predictive Coding, Quantum Biology.

1 Introduction: The Crisis of Explanation

The pharmacodynamics of Lithium (Li^+) present a profound paradox in molecular biology. As a monoatomic cation lacking complex conformational structure, its ability to stabilize the dynamic systems of human cognition and mood is remarkable. While it remains a critical medication for reducing suicide rates in bipolar patients, the physical basis of its action is not yet fully understood.

The standard pharmacological model relies on the concept of competitive inhibition. Given that the ionic radius of Li^+ (0.76 Å) is nearly identical to that of Magnesium Mg^{2+} (0.72 Å), it is generally assumed that lithium displaces magnesium in critical Mg-dependent enzymes, such as Glycogen Synthase Kinase-3β (GSK-3β) and Inositol Monophosphatase (IMPase) [1]. However, this model encounters quantitative discrepancies. The inhibition constant K_i for lithium typically

*Correspondence: a.borozentsev05@outlook.fr

falls in the range of $2 - 5$ mM, whereas clinical efficacy is maintained at $0.6 - 1.0$ mM. At these concentrations, classical competition kinetics suggest that lithium should be outcompeted by intracellular magnesium.

A significant anomaly exists in the form of Isotope Selectivity. Historical studies [3] and more recent work [4] have reported that ^6Li ($I = 1$) can be more potent in altering behavior in rodent models than its heavier counterpart, ^7Li ($I = 3/2$). Furthermore, recent biochemical assays indicate potential mitochondrial discrimination between these isotopes [7]. If biological activity were governed solely by electrostatic forces, such isotopic differentiation would be unexpected. These observations suggest the potential involvement of non-classical, quantum mechanical degrees of freedom dependent on nuclear properties.

2 Theoretical Framework: Spin Dynamics

We postulate that the mechanism of lithium action may be governed by the **Radical Pair Mechanism (RPM)**. While this mechanism is established in the context of avian magnetoreception [5], its application to enzymology offers a potential resolution to the sensitivity paradox observed in lithium therapy.

2.1 Target 1: Mitochondrial ROS Regulation

We identify the regulation of mitochondrial **Reactive Oxygen Species (ROS)** as a physically viable target. The recombination of radical pairs, such as the Flavin-Superoxide pair ($FADH^\bullet ... O_2^{\bullet -}$), is spin-selective. The Singlet state typically leads to hydrogen peroxide (H_2O_2) formation, a signaling molecule, whereas the Triplet state may allow the escape of free superoxide. Lithium, by introducing a nuclear spin, could modulate the Singlet/Triplet mixing rate via Hyperfine Interaction [6], potentially linking nuclear spin dynamics to cellular health.

2.2 Target 2: The ATP-Magnesium Complex

A secondary target involves the ATP-Magnesium complex in kinase reactions. Evidence of magnetic isotope effects in kinase catalysis [2] suggests the existence of a transient **Single Electron Transfer (SET)** step:

$$ATP + Mg^{2+} \xrightarrow{SET} [ATP \dots Mg^+]^\bullet \rightarrow \text{Products} \tag{1}$$

If lithium replaces magnesium, its nuclear spin would modulate the lifetime of this intermediate, potentially acting as a "kinetic valve" for the overall reaction rate.

2.3 Quantum Arithmetic and Coherence Times

A common objection to quantum biology is thermal decoherence. However, reaction coherence differs from computational coherence. The spin state needs only to survive long enough to influence a chemical bond choice. The interaction strength is determined by the **Hyperfine Interaction (HFI)**:

$$E_{HFI} \approx g\mu_B B_{loc} \tag{2}$$

For a lithium nucleus, the local magnetic field B_{loc} generated by the nucleus is estimated in the range of **1–5 mT**. The timescale of spin mixing ($\tau_{mix} \sim \hbar / E_{HFI}$) is in the nanosecond range, while the lifetime of radical pairs in enzyme pockets often extends to microseconds. Thus, the quantum operation may occur faster than thermal destruction.

2.4 Kinetic Evolution: The Haberkorn Equation

To link quantum states to macroscopic yields, we employ the Haberkorn master equation for the density matrix ρ:

$$\frac{d\rho}{dt} = -\frac{i}{\hbar}[\hat{H}, \rho]_- - \frac{k_S}{2}\{P_S, \rho\}_+ - \frac{k_T}{2}\{P_T, \rho\}_+ \tag{3}$$

The Hamiltonian $\hat{H}$ contains the nuclear spin term $\hat{I}$:

$$\hat{H} = \hat{H}_{Zeeman} + \sum a_k \hat{S} \cdot \hat{I}_k \tag{4}$$

Since ^6Li and ^7Li possess different gyromagnetic ratios and spin quantum numbers, they generate different mixing frequencies ω_{ST}. Consequently, the simulation predicts different product yields for the two isotopes.




Figure 1: **Simulation of Enzymatic Amplification (Haberkorn Kinetics).** Under modeled conditions of extended coherence ($\tau > 10\mu s$), typical of the viscous enzyme active site, the simulation predicts an amplification of the quantum difference between isotopes. ^6Li (Spin $I = 1$, Blue) demonstrates a sustained ability to preserve the Singlet state compared to ^7Li (Spin $I = 3/2$, Red), while the spin-zero control ^{24}Mg remains inert. This supports the "Quantum Valve" hypothesis.

This implies that lithium could act as a **"Quantum Probability Valve,"** modulating the time the system spends in a reactive state.

3 The Clinical Spectrum

If lithium modulates metabolic kinetics via spin dynamics, its effects should be observable across pathologies involving energy regulation and neural rhythmicity.

3.1 Bipolar Disorder and Homeostasis

In the manic phase, neural circuits exhibit hyper-synchronous firing. By introducing nuclear spin interactions, lithium is hypothesized to induce a **Quantum Zeno Effect**. It effectively measures the state of the reaction frequently, forcing the system into a non-reactive subspace. This prevents the system from accelerating into a runaway metabolic state, potentially acting as a thermodynamic damper.




Figure 2: **Time-Domain Quantum Coherence (Oscillatory Dynamics).** The temporal evolution of the Singlet state probability at a fixed magnetic field ($B = 0.5$). While ^{24}Mg (dotted black) maintains a static state, lithium isotopes induce coherent oscillations between Singlet and Triplet states. Crucially, ^{6}Li (blue) and ^{7}Li (red) exhibit distinct phase and frequency characteristics, theoretically allowing ^{6}Li to act as a unique "frequency modulator" for neural synchronization.

In depression, lithium's inhibition of GSK-3β promotes neurotrophic factors, facilitating synaptic plasticity [15]. The drug thus acts not as a sedative or a stimulant, but as a stabilizer of the phase transition.

3.2 Neuroprotection in Alzheimer's Disease

Clinical trials indicate that microdoses of lithium can arrest cognitive decline [10]. A purely competitive inhibitor would require toxic doses to fully block GSK-3β. However, a spin-dependent mechanism can alter the branching ratio of the enzyme at trace concentrations. By stabilizing the Singlet state, lithium may prevent the hyper-phosphorylation of tau.

3.3 Epilepsy and Phase Transitions

Lithium exhibits a biphasic profile in epilepsy [11], characteristic of a **Control Parameter** near a phase transition. Lithium tunes the system towards criticality. Low doses introduce necessary

spin entropy, breaking rigid seizure patterns. High doses, however, may saturate the system.

3.4 Cotard's Syndrome and Interoception

From the perspective of the **Free Energy Principle** [17], Cotard's syndrome represents a failure of interoceptive inference. Lithium augmentation has been reported as effective [12]. Our model suggests a mechanistic explanation: by restoring mitochondrial respiration via the spin-dependent RPM, lithium re-energizes the Salience Network, allowing the brain to update its model of the "living self."

4 Discussion

4.1 Addressing Alternative Explanations

While mass effects (Kinetic Isotope Effect) exist, they are independent of magnetic fields. Spin effects, by definition, vanish in zero field. The magnitude of the isotope effect in biological models points toward a mechanism beyond simple diffusion [6].

4.2 Cognitive Implications

From a cognitive neuroscience perspective, the stabilization of mitochondrial kinetics translates into the regularization of neural oscillations. By modulating ATP availability, lithium may tune the **"precision weighting"** of prediction errors. Lithium restores the signal-to-noise ratio at the quantum level, manifesting macroscopically as regained cognitive control.

4.3 Strategic Roadmap

If this model holds, it implies the possibility of **Isotope-Enriched Therapeutics** (using ^{6}Li to reduce toxic load) and **Magnetic-Pharmacological Resonance** (using external fields to modulate drug activity). This points toward a new era of Quantum Pharmacogenomics.

5 Proposed Experiment: The Decisive Test

To definitively distinguish between mass and spin effects, I propose a comparative assay: Run an in vitro kinetic assay of purified GSK-3β or mitochondrial ROS generation using three groups: Enzyme + $^{6}LiCl$, Enzyme + $^{7}LiCl$, and a **Negative Control Group** with Enzyme + ^{24}Mg (Spin $I = 0$). The variable is the magnetic field. The assay should be performed in Earth's Magnetic Field and in a **Hypomagnetic Chamber** (Zero-Field).




Figure 3: **Computational Verification of the Magnetic Isotope Effect.** A first-principles simulation of the Radical Pair Mechanism using exact diagonalization. **(Main Plot):** The negative control ^{24}Mg (dotted line) shows zero sensitivity to the magnetic field, validating the model. ^{6}Li and ^{7}Li show distinct magnetic field dependencies. **(Inset):** The differential plot (Δ) reveals a theoretical "Quantum Advantage" window where ^{6}Li efficiency exceeds ^{7}Li by approximately 8-10%.

Prediction: The difference between ^{6}Li and ^{7}Li should vanish or invert in Zero-Field if the mechanism is spin-dependent. Conversely, the kinetics of ^{24}Mg should remain unchanged. This result would rigorously test the quantum model against classical mass-based explanations.

6 Conclusion

The results suggest a shift in focus from finding a specific lithium receptor to understanding its interaction with the **quantum fabric** of biological matter. By modulating nuclear spin dynamics, lithium appears to adjust the thermodynamic parameters of the brain. This model offers a unified explanation for disconnected clinical data and points toward the potential of **Quantum Pharmacology**.

Data Availability

The datasets generated during and/or analysed during the current study, including the software code used for the mathematical experiment (exact diagonalization simulations) and the resulting data, are available in the **Zenodo** repository at `https://doi.org/10.5281/zenodo.17834884`. This code is provided to enhance transparency and ensure full reproducibility of the quantum mechanical simulations presented in Figures 1–3.

Funding

The author declares that no funds, grants, or other support were received during the preparation of this manuscript.

Competing Interests

The author declares no competing financial or non-financial interests.

Author Contributions

A.B. conceived the study, developed the Quantum-Thermodynamic theoretical framework, wrote the Python simulation code, performed the data analysis, and wrote the manuscript.

Acknowledgements

I am grateful to the Google Gemini project for assistance with translation and stylistic refinement. I confirm that the intellectual content, theoretical derivation, experimental design, and scientific conclusions are my own original work.

References

[1] Quiroz, J. A., et al. (2004). Molecular therapeutics for bipolar disorder. *Archives of General Psychiatry*, 61(9), 922-931.

[2] Buchachenko, A. L., et al. (2005). Magnetic isotope effect of magnesium in phosphokinase catalysis. *PNAS*, 102(31), 10793-10796.

[3] Sechzer, J. A., et al. (1986). Lithium isotopes: Differential effects on behavior. *Biological Psychiatry*, 21(13), 1258-1266.

[4] Ettenberg, A., et al. (2020). Differential effects of lithium isotopes in a ketamine-induced hyperactivity model of mania. *Pharmacol. Biochem. Behav.*, 190, 172875.

[5] Hore, P. J., & Mouritsen, H. (2016). The radical-pair mechanism of magnetoreception. *Annual Review of Biophysics*, 45, 299-344.

[6] Zadeh-Haghighi, H., & Simon, C. (2021). Entangled radicals may explain lithium effects on hyperactivity. *Scientific Reports*, 11, 1542.

[7] Deline, M. L., et al. (2023). Lithium isotopes differentially modify mitochondrial amorphous calcium phosphate cluster size distribution. *Frontiers in Cell and Developmental Biology*, 11, 1200119.

[8] Poet, M., et al. (2023). Biological fractionation of lithium isotopes by cellular Na+/H+ exchangers unravels fundamental transport mechanisms. *iScience*, 26(11), 108160.

[9] Forlenza, O. V., et al. (2011). Disease-modifying properties of long-term lithium treatment. *Br. J. Psychiatry*, 198, 351-356.

[10] Nunes, M. A., et al. (2013). Microdose lithium treatment stabilized cognitive impairment in patients with Alzheimer's disease. *Current Alzheimer Research*, 10(1), 104-107.

[11] Shorvon, S. D. (2009). Drug treatment of epilepsy in the century of the ILAE. *Epilepsia*, 50, 69-130.

[12] Grover, S., et al. (2014). Cotard's syndrome: Two case reports and a brief review of literature. *Journal of Neurosciences in Rural Practice*, 5, S59.

[13] Jakobsson, E., et al. (2017). Isotope-specific lithium bioactivity in mitochondria. *RSC Advances*, 7, 123-130.

[14] Wada, A. (2009). Lithium and neuropsychiatric therapeutics: neuroplasticity via GSK-3beta. *Journal of Pharmacological Sciences*, 110, 14-28.

[15] Manji, H. K., et al. (2000). The underlying neurobiology of bipolar disorder. *World Psychiatry*, 2, 136-146.

[16] Fisher, M. P. A. (2015). Quantum cognition: The possibility of processing with nuclear spins in the brain. *Annals of Physics*, 362, 593-602.

[17] Friston, K. (2010). The free-energy principle: a unified brain theory? *Nature Reviews Neuroscience*, 11(2), 127-138.

[18] Klein, P. S., & Melton, D. A. (1996). A molecular mechanism for the effect of lithium on development. *PNAS*, 93(16), 8455-8459.

[19] Hashimoto, R., et al. (2002). Lithium protects against oxidative stress-induced cell death in mouse hippocampal neurons. *Life Sciences*, 72(21), 2431-2446.