



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 4, 2025 – 02:03 pm GMT

PDB ID : 9TFG / pdb_00009tfg
Title : KIT kinase domain in complex with a 6-,7-dimethoxyquinoline-based compound (19)
Deposited on : 2025-11-27
Resolution : 1.70 Å(reported)

This wwPDB validation report is for manuscript review

This is a Full wwPDB X-ray Structure Validation Report.

This report is produced by the wwPDB biocuration pipeline after annotation of the structure.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	5161 (1.70-1.70)
Clashscore	180529	5671 (1.70-1.70)
Ramachandran outliers	177936	5594 (1.70-1.70)
Sidechain outliers	177891	5594 (1.70-1.70)
RSRZ outliers	164620	5159 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	327	<div> <div>7%</div> <div>93%</div> <div>5%</div> </div>
1	B	327	<div> <div>6%</div> <div>86%</div> <div>5%</div> <div>9%</div> </div>

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
 Validation Pipeline (wwPDB-VP) : 2.46

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 5662 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mast/stem cell growth factor receptor Kit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	319	Total	C	N	O	S	0	14	0
			2597	1674	419	484	20			
1	B	298	Total	C	N	O	S	0	15	0
			2400	1550	396	436	18			

There are 70 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	548	GLY	-	expression tag	UNP P10721
A	549	SER	-	expression tag	UNP P10721
A	550	MET	-	expression tag	UNP P10721
A	563	SER	ILE	engineered mutation	UNP P10721
A	569	SER	VAL	engineered mutation	UNP P10721
A	609	GLN	TYR	engineered mutation	UNP P10721
A	631	SER	LEU	engineered mutation	UNP P10721
A	651	GLU	MET	engineered mutation	UNP P10721
A	662	HIS	ILE	engineered mutation	UNP P10721
A	688	GLU	-	linker	UNP P10721
A	689	PHE	-	linker	UNP P10721
A	690	VAL	-	linker	UNP P10721
A	691	PRO	-	linker	UNP P10721
A	692	TYR	-	linker	UNP P10721
A	693	LYS	-	linker	UNP P10721
A	754	VAL	-	linker	UNP P10721
A	755	ALA	-	linker	UNP P10721
A	756	PRO	-	linker	UNP P10721
A	757	GLU	-	linker	UNP P10721
A	758	ASP	-	linker	UNP P10721
A	759	LEU	-	linker	UNP P10721
A	760	TYR	-	linker	UNP P10721
A	761	LYS	-	linker	UNP P10721
A	762	ASP	-	linker	UNP P10721
A	763	PHE	-	linker	UNP P10721

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	764	LEU	-	linker	UNP P10721
A	765	THR	-	linker	UNP P10721
A	768	HIS	ASP	engineered mutation	UNP P10721
A	804	ASN	ARG	engineered mutation	UNP P10721
A	825	ASP	VAL	engineered mutation	UNP P10721
A	844	SER	CYS	engineered mutation	UNP P10721
A	890	SER	LEU	engineered mutation	UNP P10721
A	894	TYR	HIS	engineered mutation	UNP P10721
A	912	ASP	LEU	engineered mutation	UNP P10721
A	923	ASP	LEU	engineered mutation	UNP P10721
B	548	GLY	-	expression tag	UNP P10721
B	549	SER	-	expression tag	UNP P10721
B	550	MET	-	expression tag	UNP P10721
B	563	SER	ILE	engineered mutation	UNP P10721
B	569	SER	VAL	engineered mutation	UNP P10721
B	609	GLN	TYR	engineered mutation	UNP P10721
B	631	SER	LEU	engineered mutation	UNP P10721
B	651	GLU	MET	engineered mutation	UNP P10721
B	662	HIS	ILE	engineered mutation	UNP P10721
B	688	GLU	-	linker	UNP P10721
B	689	PHE	-	linker	UNP P10721
B	690	VAL	-	linker	UNP P10721
B	691	PRO	-	linker	UNP P10721
B	692	TYR	-	linker	UNP P10721
B	693	LYS	-	linker	UNP P10721
B	754	VAL	-	linker	UNP P10721
B	755	ALA	-	linker	UNP P10721
B	756	PRO	-	linker	UNP P10721
B	757	GLU	-	linker	UNP P10721
B	758	ASP	-	linker	UNP P10721
B	759	LEU	-	linker	UNP P10721
B	760	TYR	-	linker	UNP P10721
B	761	LYS	-	linker	UNP P10721
B	762	ASP	-	linker	UNP P10721
B	763	PHE	-	linker	UNP P10721
B	764	LEU	-	linker	UNP P10721
B	765	THR	-	linker	UNP P10721
B	768	HIS	ASP	engineered mutation	UNP P10721
B	804	ASN	ARG	engineered mutation	UNP P10721
B	825	ASP	VAL	engineered mutation	UNP P10721
B	844	SER	CYS	engineered mutation	UNP P10721
B	890	SER	LEU	engineered mutation	UNP P10721

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	894	TYR	HIS	engineered mutation	UNP P10721
B	912	ASP	LEU	engineered mutation	UNP P10721
B	923	ASP	LEU	engineered mutation	UNP P10721

- Molecule 2 is 6,7-dimethoxy-4-[4-[5-(phenylmethyl)pyrimidin-2-yl]piperazin-1-yl]quinoline (CCD ID: A1JVT) (formula: C₂₆H₂₇N₅O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			33	26	5	2		
2	B	1	Total	C	N	O	0	0
			33	26	5	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	337	Total	O	0	0
			337	337		
3	B	262	Total	O	0	0
			262	262		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Mast/stem cell growth factor receptor Kit



- Molecule 1: Mast/stem cell growth factor receptor Kit



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	58.65Å 58.97Å 192.45Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.11 – 1.70 48.11 – 1.70	Depositor EDS
% Data completeness (in resolution range)	99.8 (48.11-1.70) 99.8 (48.11-1.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.38 (at 1.70Å)	Xtriage
Refinement program	PHENIX 1.21.2_5419	Depositor
R, R_{free}	0.174 , 0.214 0.175 , 0.215	Depositor DCC
R_{free} test set	3715 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	28.6	Xtriage
Anisotropy	0.436	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 42.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.019 for k,h,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5662	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: A1JVT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.27	0/2700	0.45	0/3653
1	B	0.26	0/2505	0.46	0/3395
All	All	0.26	0/5205	0.46	0/7048

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2597	0	2545	11	0
1	B	2400	0	2353	14	0
2	A	33	0	0	0	0
2	B	33	0	0	0	0
3	A	337	0	0	1	0
3	B	262	0	0	2	0
All	All	5662	0	4898	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:878:LYS:HG2	1:B:882:MET:HE2	1.70	0.73
1:A:774:TYR:HE1	1:A:925:GLU:HG3	1.63	0.64
1:B:873:MET:HE3	1:B:882:MET:HE3	1.81	0.62
1:B:873:MET:HE3	1:B:882:MET:CE	2.29	0.61
1:A:684:ARG:NH2	3:A:1106:HOH:O	2.37	0.57
1:A:817:ILE:HD12	1:A:821:SER:HA	1.86	0.57
1:A:559:VAL:HA	1:A:568:TYR:CD1	2.42	0.55
1:A:774:TYR:CE1	1:A:925:GLU:HG3	2.42	0.53
1:A:584:PHE:CD1	1:A:585:PRO:HD2	2.47	0.50
1:A:638[A]:MET:HE1	1:A:661:THR:HG21	1.93	0.49
1:B:638[A]:MET:HE1	1:B:661:THR:HG21	1.94	0.49
1:B:873:MET:HE2	1:B:879:PHE:HD1	1.78	0.49
1:B:570:TYR:CE2	1:B:819:ASN:HA	2.48	0.48
1:A:587:ASN:ND2	1:A:587:ASN:H	2.12	0.47
1:B:787:ASN:HD22	1:B:818:LYS:HA	1.80	0.47
1:B:631:SER:OG	1:B:634[B]:ARG:NH2	2.49	0.45
1:B:770[B]:LEU:HD23	3:B:1318:HOH:O	2.16	0.45
1:A:864[B]:SER:OG	1:A:892:PRO:HB3	2.18	0.43
1:B:586:ARG:HG2	3:B:1221:HOH:O	2.19	0.42
1:A:559:VAL:HA	1:A:568:TYR:HD1	1.81	0.42
1:B:690:VAL:O	1:B:765:THR:HA	2.19	0.42
1:B:878:LYS:CG	1:B:882:MET:HE2	2.46	0.41
1:B:770[B]:LEU:HG	1:B:928:ILE:HG12	2.03	0.41
1:A:913:LYS:HB3	1:A:913:LYS:HE2	1.88	0.41
1:B:873:MET:HE3	1:B:882:MET:HE1	2.01	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	327/327 (100%)	320 (98%)	7 (2%)	0	100	100
1	B	309/327 (94%)	303 (98%)	6 (2%)	0	100	100
All	All	636/654 (97%)	623 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	286/285 (100%)	286 (100%)	0	100	100
1	B	259/285 (91%)	259 (100%)	0	100	100
All	All	545/570 (96%)	545 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	567	ASN
1	A	587	ASN
1	A	680	ASN
1	B	787	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	A1JVT	B	1001	-	37,37,37	1.20	4 (10%)	50,51,51	1.52	9 (18%)
2	A1JVT	A	1001	-	37,37,37	1.19	5 (13%)	50,51,51	1.38	7 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A1JVT	B	1001	-	-	0/16/26/26	0/5/5/5
2	A1JVT	A	1001	-	-	0/16/26/26	0/5/5/5

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1001	A1JVT	C21-N20	3.39	1.42	1.35
2	A	1001	A1JVT	C21-N22	2.95	1.39	1.34
2	B	1001	A1JVT	C13-C12	2.76	1.44	1.38
2	A	1001	A1JVT	C13-C12	2.64	1.43	1.38
2	A	1001	A1JVT	C21-N20	2.62	1.40	1.35
2	B	1001	A1JVT	C21-N33	2.59	1.39	1.34
2	B	1001	A1JVT	C21-N22	2.27	1.38	1.34
2	A	1001	A1JVT	C18-N15	2.26	1.50	1.46
2	A	1001	A1JVT	C17-N20	2.01	1.49	1.46

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1001	A1JVT	O02-C03-C04	-4.89	118.92	125.24
2	A	1001	A1JVT	O02-C03-C04	-4.61	119.28	125.24
2	B	1001	A1JVT	O02-C03-C08	4.50	121.68	115.41
2	A	1001	A1JVT	O02-C03-C08	4.26	121.34	115.41
2	B	1001	A1JVT	C25-C24-C32	3.03	125.49	120.83
2	B	1001	A1JVT	N33-C21-N22	-2.79	122.36	127.06
2	B	1001	A1JVT	N33-C21-N20	2.38	119.47	116.90
2	A	1001	A1JVT	C04-C05-C14	2.29	125.53	122.81
2	A	1001	A1JVT	N33-C21-N22	-2.28	123.21	127.06
2	A	1001	A1JVT	N22-C21-N20	2.21	119.30	116.90
2	B	1001	A1JVT	C16-C17-N20	2.21	114.99	110.70
2	B	1001	A1JVT	C17-C16-N15	2.17	114.92	110.70
2	B	1001	A1JVT	C04-C05-C14	2.17	125.39	122.81
2	A	1001	A1JVT	C04-C05-C06	-2.14	116.31	118.94
2	B	1001	A1JVT	C32-N33-C21	2.07	117.47	115.64
2	A	1001	A1JVT	C25-C24-C23	2.03	123.95	120.83


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	319/327 (97%)	0.09	24 (7%) 22 22	13, 30, 63, 80	14 (4%)
1	B	298/327 (91%)	0.10	19 (6%) 27 27	15, 32, 62, 79	15 (5%)
All	All	617/654 (94%)	0.09	43 (6%) 24 24	13, 31, 63, 80	29 (4%)

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	692	TYR	4.4
1	B	614	SER	4.3
1	B	615	ASP	4.3
1	A	548	GLY	3.8
1	A	557	TRP	3.8
1	B	612	ILE	3.6
1	A	560	VAL	3.5
1	B	613	LYS	3.4
1	A	570[A]	TYR	3.3
1	A	760	TYR	3.3
1	A	693	LYS	3.1
1	A	819	ASN	3.0
1	A	562	GLU	3.0
1	B	566	ASN	3.0
1	A	821	SER	2.9
1	B	600	PHE	2.9
1	A	565	GLY	2.8
1	A	759	LEU	2.8
1	B	616	ALA	2.8
1	B	821	SER	2.7
1	A	559	VAL	2.7
1	B	828	ASN	2.7
1	B	819	ASN	2.7
1	B	762	ASP	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	632	THR	2.6
1	A	828	ASN	2.5
1	A	817	ILE	2.4
1	A	757	GLU	2.4
1	A	931	SER	2.4
1	B	688	GLU	2.4
1	A	555	VAL	2.4
1	A	872	GLY	2.4
1	B	763	PHE	2.3
1	B	693	LYS	2.2
1	A	612	ILE	2.2
1	A	630	HIS	2.2
1	B	646	TYR	2.2
1	A	822	ASN	2.1
1	B	931	SER	2.1
1	B	827	GLY	2.1
1	B	687	ASP	2.0
1	B	820	ASP	2.0
1	A	549	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	A1JVT	B	1001	33/33	0.94	0.08	24,29,37,39	0
2	A1JVT	A	1001	33/33	0.96	0.06	22,26,31,33	0


The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around A1JVT B 1001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.