

		Total survivors (n=237,075)
Age (years)		72.3(18.7)
	<15	5862 (2.47%)
	15 - 44	13,180 (5.56%)
	45 - 64	37,783 (15.9%)
	65 - 74	42,645 (18%)
	75 - 84	75,103 (31.7%)
	>84	62,502 (26.4%)
Gender		
	Male	132,514 (70.7%)
Charlson Index		2.36 (2.12)
	=0	45,401 (19.2%)
	1 - 2	98,915 (41.7%)
	3 - 4	62,386 (26.3%)
	≥ 5	30,373 (12.8%)
Comorbidities		
	Chronic kidney disease	69,081 (29.1%)
	Chronic obstructive pulmonary disease	60,455 (25.5%)
	Active neoplasms	59,370 (25%)
	Uncomplicated diabetes mellitus	54,338 (22.9%)
	Congestive heart failure	50,152 (21.2%)
	Liver disease	28,612 (12.1%)
	Dementia	18,644 (7.86%)
	Cerebrovascular disease	17,423 (7.3%)
	Complicated diabetes mellitus	12,631 (5.3%)
	Liver Cirrhosis	8,710 (3.7%)
	Disseminated neoplasms	9,852 (4.16%)
	Human Immunodeficiency virus	2,096 (0.8%)
Sepsis Source		
	Genitourinary	98,641 (41.6%)
	Respiratory	75,831 (32%)
	Unknown	26,950 (11.4%)
	Abdomin-biliary	23,837 (10.1%)
	Related to intravascular devices	17,862 (7.53%)

	Tissues and soft tissues	10,901 (4.6%)
	Others	8,112 (3.42%)
	Endocarditis	2,687 (1.13%)
	Central Nervous System	1,839 (0.78%)
Hospital stay		
	<4 days	37,225 (15.7%)
	4 - 7 days	55,853 (23.6%)
	8 - 14 days	67,673 (28.5%)
	15 - 21 days	30,515 (12.9%)
	22 - 28 days	15,482 (6.53%)
	>28 days	30,327 (12.8%)

Table 1. Baseline characteristics of surviving patients at hospital discharge.