

Supplementary Information-1

Adipose-derived stem cell exosomes promote critical-sized bone defect repair by enhancing the homing of bone marrow mesenchymal stem cells

Jingjie Yang¹, Jing Jing¹, Xuesha Tong¹, Songyang Ma¹, Ye Qiu¹, Yang Liu^{1*}, Yunji Wang^{1*}.

Table S1. Sequences of qRT-PCR primers.

Gene	Forward	Reverse
GAPDH	GGCACAGTCAAGGCTGAGAATG	ATGGTGGTGAAGACGCCAGTA
Cxcr4	GCGTTTGGTGCTCCGGTA	GAAGCAGGGTTCCTTGTTGGA
Ccr7	GGAGACAGAGACAAGAGACACAA	TTCTTGAAGCACACCGACTCATA
MMP2	CGCTGCCTTTAACTGGAGTAAGA	CCAGGTTATCAGGGATGGCATTC
MMP9	CTACTCTGCCTGCACCACTAAAG	CTCGAGTAGGACAGAAGCCATAC
Rac1	AATGCGTTCCCTGGAGAGTACAT	CAAATGATGCAGGACTCACAAGG
Alp	CCAAAGGCTTCTTCTTGCTAGTG	TGATCAGCAGTAACCACAGTCAA
Col1a1	GTGAGACGTGGAAACCTGATGTA	GACTTCTGCGTCTGGTGATACAT
OCN	GAGGGCAGTAAGGTGGTGAATAG	GGGTCGAGTCCTGGAGAGTAG
Runx2	GGCCACTTACCACAGAGCTATTA	GTGTCTGCCTGGGATCTGTAATC
OPN	CGCATTACAGCAAACACTCAGAT	CCGTTTCTTCAGAGGACACAGAA

Supplementary Information-2

Fig.1G

