

The Association Between Short-Term Nitrogen Dioxide (NO₂) Exposure and Ischemic Stroke or Transient Ischemic Attacks (TIA): A National Study in Israel

Placidus S. Fernando

Icahn School of Medicine at Mount Sinai

Helena Krasnov

Icahn School of Medicine at Mount Sinai

Itai Kloog

Icahn School of Medicine at Mount Sinai

Mike Z. He

Icahn School of Medicine at Mount Sinai

Gal Ifergane

Soroka University Medical Center

Inbar Zucker

Soroka University Medical Center

Victor Novack

Soroka University Medical Center

Carmit Libruder

Soroka University Medical Center

Maayan Yitshak-Sade

`maayan.yitshak-sade@mssm.edu`

Icahn School of Medicine at Mount Sinai

Article

Keywords: Air pollution, Nitrogen dioxide, Ischemic stroke, Transient ischemic attack

Posted Date: February 25th, 2026

DOI: <https://doi.org/10.21203/rs.3.rs-8875648/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.
[Read Full License](#)

Additional Declarations: No competing interests reported.

Version of Record: A version of this preprint was published at Scientific Reports on October 15th, 2026.
See the published version at <https://doi.org/10.1038/s41598-026-75950-x>.

Abstract

Background: Air pollution is an environmental risk factor for stroke, but most studies have focused on fine particulate matter (PM_{2.5}), and fewer have investigated the independent association with exposure to nitrogen dioxide (NO₂).

Methods: We conducted a nationwide case-crossover study including all ischemic stroke or transient ischemic attack (TIA) reported in the Israeli National Stroke Registry from 2014 to 2018. We obtained daily NO₂ levels from high-resolution geospatial models and linked to residential addresses. We used distributed lag nonlinear models to assess the cumulative association with NO₂ exposure over a lag of 0–5 days, adjusting for PM_{2.5} and temperature. We further evaluated the modification of the associations by age, sex, urbanicity, and comorbidities (ischemic heart disease, hypertension, diabetes).

Results: We included 85,162 cases, about 54% were males, and the mean age was 71.8. Cumulative NO₂ exposure was associated with increased odds of ischemic stroke or TIA (odd ratio – OR and 95% confidence interval –1.064 [1.041–1.088]), with the strongest effects 3–5 days before the event. The associations were more pronounced in urban areas (1.070[1.038;1.103]) compared to suburban areas (1.054[1.021;1.088]). Although the overall association was similar across ages and sexes, the lag specific associations on lag days 3-5 were more pronounced in younger adults and women.

Conclusion: Short-term exposure to NO₂ is associated with an increased risk of ischemic stroke and TIA, with differential associations across urbanicity, sex and age groups. These results underscore the urgent need for public health initiatives to reduce NO₂ emissions.

Synopsis

Short-term NO₂ exposure increases ischemic stroke and TIA risk, informing air-quality policy to curb traffic emissions and protect vulnerable populations.

Introduction

Stroke is one of the leading causes of death and disability worldwide¹. Each year, there are over 12 million new stroke cases worldwide, a number that has seen over a 50% increase over the last two decades². Therefore, research in recent decades has focused on identifying modifiable risk factors that can be targeted to reduce the global stroke burden³. There is substantial evidence indicating that exposure to air pollution increases the risk of cardiovascular diseases (CVD) beyond traditional CVD risk factors⁴. Given that the vast majority of the global population currently lives in regions where the air pollution levels exceed the World Health Organization's Global Air Quality Guidelines, air pollution represents a critical target for stroke prevention^{5,6}. Most existing research has focused on particulate matter smaller than 2.5 microns (PM_{2.5}), indicating an increased risk of stroke associated with both long

(1.01 per 10 $\mu\text{g}/\text{m}^3$ CI 1.011-1.012) and short-term exposures (1.06 per 10 $\mu\text{g}/\text{m}^3$ 95% CI 1.018–1.105) ^{6,7}. Although nitrogen dioxide (NO_2) has been studied, the evidence is less extensive than for $\text{PM}_{2.5}$, and important research gaps remain ^{8,9}. These gaps underscore the need to investigate the health effects independently associated with NO_2 exposure ⁶.

NO_2 is an anthropogenic air pollutant originating primarily from traffic and industrial sources through the combustion of fossil fuels ¹⁰. Studies suggest that NO_2 may increase stroke risk by triggering oxidative stress, systemic inflammation, and endothelial dysfunction, leading to blood clots and restricted blood flow to the brain ¹¹. A large meta-analysis in 2023 found that NO_2 exposure is associated with stroke hospitalizations and mortality (1.014 per 10 ppb, CI 1.009-1.019) ⁶. However, the analysis concluded that it had insufficient information to establish a causal relationship between exposure to NO_2 and stroke ⁶.

In this comprehensive national study, we leverage the Israeli National Stroke Registry (INSR) data to investigate the association between short-term NO_2 exposure and ischemic stroke or transient ischemic attacks (TIA). We further investigate differential susceptibility among population groups by urbanicity, demographic, and clinical characteristics. Israel's unique topography, which includes mountains, coastal lowlands, and deserts, offers a diverse environment to investigate NO_2 exposure and cardiovascular health effects in varying climates ¹². These findings may inform policy and interventions aimed at reducing NO_2 exposure levels and stroke burden in the population.

Methods

Population and outcome

We used data from the INSR, which includes all permanent residents in Israel admitted to general hospitals with a primary diagnosis of stroke or TIA, reported using the International Classification of Diseases, ninth edition (ICD-9). We included all ischemic stroke (ICD-9:433, 434.x1, 436, or 997.02) and TIA (ICD-9: 435) cases of individuals 18 years and older across Israel between the years 2014 - 2018. The INSR data are retrieved from electronic medical records and include diagnoses and comorbidities coded using ICD-9, encompassing hospitalization diagnoses, chronic conditions, and discharge diagnoses. Sociodemographic information (age, sex, ethnicity, and geocoded address) is retrieved from the Ministry of Interior database and linked to the INSR. Ethnicity was grouped into Jewish versus non-Jewish due to the small percentage of the non-Jewish population (Arab, Druze, or other ethnic groups).

This study was approved by the institutional review boards at Soroka University Medical Center in Beer Sheva, Israel (study 01718-19), and the Icahn School of Medicine at Mount Sinai, New York, USA (study 21-00076). A waiver of informed consent was granted due to the minimal risk nature of the study and the use of de-identified registry data with no direct participant contact. All methods were performed in accordance with the relevant institutional and national guidelines and regulations.

Exposure

We obtained daily 1-km² predictions of NO₂ from a hybrid spatiotemporal model we built over Israel from 2004 to 2022¹³. The model, which is based on methodology we used previously for an NO₂ model in Mexico City, incorporates a combination of satellite, land use, and meteorological data and employs machine learning-based methods (random forest and gradient boosting) for prediction¹⁴. All models were rigorously evaluated using 10-fold cross-validation and demonstrated strong performance¹⁴⁻¹⁷. The exposures were then linked to each individual based on the proximity of the geocoded residential address to the corresponding 1-km² grid cells of the satellite-based model predictions. We excluded 1,289 individuals without a valid residential address (1.6%).

Covariates

Models were adjusted for temperature averaged over lag days 0-4. In a sensitivity analysis, we further adjusted the model for fine particulate matter (PM_{2.5}) exposure. We obtained daily mean PM_{2.5} and temperature estimates from satellite-based spatiotemporal models that apply a hybrid approach and estimate daily mean temperature and PM_{2.5} exposures at a fine resolution^{18,19}. Briefly, these models use both extreme gradient boosting and inverse-distance weighting techniques that incorporate aerosol optical depth, meteorology, and land-use variables to generate daily predictions at 1-km resolutions. The models were calibrated daily by applying a mixed-modeling approach using the MODIS-based multiangle implementation of atmospheric correction (MAIAC) aerosol optical depth (AOD) and various spatial and temporal predictors. This resulted in excellent performance with out-of-sample cross-validated R² values of 0.82–0.92 for the PM_{2.5} model and 0.96–0.98 for the temperature model.

Statistical analysis

We used distributed lag nonlinear models (DLNMs) in a case crossover design to investigate the associations between NO₂ exposures and ischemic stroke or TIA. This approach accounts for confounders that do not vary within a short period by design and allows the flexible estimation of lag-specific and cumulative associations over time²⁰. We defined the admission day as the case day, and matched control days by the day of the week within the same month and year. We defined the exposure window as exposure on lag days 0-5, based on the smallest Bayesian Information Criteria (BIC). We used a linear function for the exposure response association and natural splines with two degrees of freedom (df) to estimate the lag structures, based on the smallest BIC. Then, we used generalized nonlinear models (GNM) with conditional stratification. We adjusted the models for the temperature mean for lags 0-4. We selected the exposure window that yielded the strongest association with the outcome to ensure there is no residual confounding by temperature exposure.

We conducted an interaction analysis to evaluate whether the cumulative association between NO₂ exposure and ischemic stroke/TIA differed across urbanicity (cities with over 100,000 residents versus smaller cities, towns and villages), comorbidities (diabetes, hypertension, and ischemic heart disease -

IHD), sex, and age. We classified individuals as younger than 55 years versus 55 years and older, a cutoff commonly used to define stroke in young adults ²¹. We repeated the models adding an interaction term between the NO₂ cross-basis and each subgroup indicator separately. Because the exposure-response is represented by multiple spline and lag parameters, the interaction produces a set of coefficients rather than a single term. We therefore used the likelihood ratio test (LRT) to assess whether all interaction parameters significantly differ from the null, comparing models with and without the interaction term. A statistically significant LRT p-value reflects evidence of differences in the *shape* or *timing* of the lagged response across groups, which may be present even if the estimated cumulative risks over the full lag period appear similar. For interpretability, we present group-specific cumulative OR and 95% confidence intervals (CI) and provide lag-specific effects among subgroups of statistically significant modifiers.

Sensitivity analyses

We conducted several sensitivity analyses to ensure the robustness of our findings. First, we repeated the main model using alternative definitions of cumulative NO₂ exposure, summing exposure over 3-, 4-, 5-, and 6-day lag windows. Second, we repeated the main models only with ischemic stroke, excluding TIA, to ensure the associations with NO₂ did not differ between the groups. Third, since the baseline stroke risk may differ between first occurrence and repeated stroke events, we repeated the models excluding recurrent events to ensure the association did not change. Finally, we further adjusted the models to PM_{2.5} exposure on the day of the event to ensure that our results are not confounded by PM_{2.5} exposure.

Results

We included 85,162 cases, of whom 69.9% had ischemic stroke and 30.1% had TIA. About 54% were males, 80.9% were Jewish, 50.28% resided in large cities, and the mean age was 71.8 years. Cardiometabolic diseases were highly prevalent in this population, with 45.8% presenting diabetes, about 15% CHF, and 77.9% hypertension. Finally, about 3% of the cases had a history of myocardial infarction, and 13.64% had recurring stroke events (**Table 1**).

Table 1. Population characteristics (n=85,162)

Variable	Mean (SD)/ N (%)
Age	71.7 (13.4)
Sex	
<i>Male</i>	46,014 (54.03)
<i>Female</i>	39,148 (45.97)
Ethnicity	
<i>Jewish</i>	68,837 (80.83)
<i>Other</i>	16,325 (19.17)
Residence type	
<i>Urban</i>	42,795 (50.28)
<i>Suburban and rural</i>	42,311 (49.71)
Stroke type	
<i>Ischemic</i>	59,606 (69.9)
<i>TIA</i>	25,556 (30.1)
Diabetes	39,034 (45.83)
Hypertension	66,376 (77.94)
IHD	28,251 (33.17)
Died in the hospital	59,825 (70.24)
Stroke events	
<i>Single</i>	73,466 (86.26)
<i>Recurring</i>	11,696 (13.74)

The largest number of cases was observed in the summer season (33.42%). Overall, the mean NO₂ exposure level during the observation period was 12.34 ppb, with higher levels in large cities (13.62 ppb) compared to suburban and rural areas (11.02 ppb). The mean PM_{2.5} exposure level was 19.04 µg/m³, with similar levels between large cities (19.28 µg/m³) and suburban and rural areas (18.79 µg/m³). The mean temperature during the observation period was 21.40 °C (**Table 2**). The correlation between the air pollutants and temperature was low, with the highest correlation between NO₂ and temperature (r=-0.2), and NO₂ and PM_{2.5} (r=0.13). A very low correlation was observed between temperature and PM_{2.5} (r=0.05).

Table 2. Environmental characteristics

	All	Large cities	Suburban and rural
Temperature			
Mean (SD)	21.40 (5.55)	21.40 (5.53)	21.40 (5.56)
Median (IQR)	21.81 (9.79)	21.79 (9.74)	21.82 (9.82)
NO ₂			
Mean (SD)	12.32 (8.56)	13.62 (9.09)	11.02 (7.78)
Median (IQR)	10.01 (10.25)	11.40 (11.43)	8.87 (8.75)
PM _{2.5}			
Mean (SD)	19.04 (9.92)	19.28 (7.02)	18.79 (9.48)
Median (IQR)	17.25 (6.59)	17.40 (7.02)	17.13 (6.17)

Large cities were defined as those with populations exceeding 100,000 residents. Suburban and rural areas include smaller cities, towns, and villages

The cumulative OR for ischemic stroke or TIA associated with an 8 ppb (standard deviation (SD)) increase in NO₂ exposure, summed over lag days 0–5, was 1.064 (95% confidence interval [CI]: 1.041, 1.088) (**Figure 1.A**). The effects were more pronounced in lag days 3-5. For example, the OR observed on lag 1 was 0.990 (0.980; 1.001) and the OR observed on lag 5 was 1.044 (1.024; 1.065) (**Figure 1.B**). The cumulative ORs remained consistent across the alternative exposure windows, with estimates ranging from 1.050 to 1.061 when using 3- to 6-day cumulative lags (**Supplementary Table 1**).

The associations between NO₂ and ischemic stroke/TIA were statistically significantly modified by urbanicity (LRT p value 0.029), sex (LRT p value 0.008), and age (LRT p value 0.037). Overall, the cumulative associations were very similar among younger (1.068, 95% CI 1.036; 1.101) and older adults (1.060, 95% CI 1.027; 1.094), and among men (1.069, 95% CI 1.037; 1.101) and women (1.061, 95% CI 1.027; 1.096). However, as shown in **Figure 2**, the lag specific associations on lag days 3-5 were more pronounced in younger adults and women. For example, on lag day 5, the OR for NO₂ exposure was 1.114 (95% CI: 1.047, 1.184) in younger adults compared with 1.037 (95% CI: 1.016, 1.059) in older adults, and 1.074 (95% CI: 1.044, 1.105) in women compared with 1.019 (95% CI: 0.993, 1.047) in men. Finally, the cumulative OR was larger among people in urban (1.070, 95% CI 1.038; 1.103) versus rural and suburban areas (1.054, 95% CI 1.021; 1.088) with different lag structures between the groups (**Figure 2**).

Table 3. The modification of the associations between NO₂ and ischemic stroke/TIA by patient characteristics.

Subgroup	OR (95% CI)	LRT P value
Residence type		
<i>Rural/suburban</i>	1.054 (1.021; 1.088)	
<i>Urban</i>	1.070 (1.038; 1.103)	0.029
Diabetes		
<i>No</i>	1.047 (1.016; 1.078)	
<i>Yes</i>	1.087 (1.052; 1.123)	0.179
Hypertension		
<i>No</i>	1.076 (1.027; 1.127)	
<i>Yes</i>	1.062 (1.036; 1.088)	0.865
IHD		
<i>No</i>	1.064 (1.036; 1.093)	
<i>Yes</i>	1.067 (1.027; 1.109)	0.89
Sex		
<i>Men</i>	1.069 (1.037; 1.101)	
<i>Women</i>	1.061 (1.027; 1.096)	0.008
Age		
<i>55 and older</i>	1.060 (1.027; 1.094)	
<i>Younger than 55</i>	1.068 (1.036; 1.101)	0.037

We used distributed lag nonlinear models (DLNMs) in a case crossover design to investigate the associations between cumulative NO₂ exposures (lags 0-5) and ischemic stroke or TIA, adjusted for temperature. We repeated the models, adding an interaction term between the NO₂ cross-basis and each subgroup indicator separately. We used the likelihood ratio test (LRT) to assess whether all interaction parameters significantly differ from the null, comparing models with and without the interaction term. A statistically significant LRT p-value reflects evidence of differences in the *shape* or *timing* of the lagged response across groups.

Odds ratios are presented per incremental increase in NO₂ concentration and were estimated using distributed lag nonlinear models adjusted for PM_{2.5} and temperature. Error bars represent 95% confidence intervals.

As a sensitivity analysis, we repeated the main model excluding repeated events. The cumulative OR of ischemic stroke or TIA, associated with an 8-ppb increase in NO₂ exposure, summed over lag days 0-5,

remained very similar (1.061, 95% CI 1.036, 1.086). Additionally, the results remained similar after adjusting for PM_{2.5} exposure on the day of the event (1.065, 95% CI 1.042; 1.089), and no association was observed with PM_{2.5} (OR 0.997, 95%CI 0.993; 1.001 per 10 mg/m³ increase). Finally, we repeated the model only among ischemic stroke cases, excluding TIA, and the results remained similar (1.042, 95% CI 1.015, 1.070).

Discussion

In this nationwide study encompassing diverse populations and climate regions, we observed increased odds of ischemic stroke or TIA associated with short-term NO₂ exposure. The associations were strongest at lags 3-5 days and were marginally stronger among residents of large cities, women, and adults <55 years, while remaining broadly similar across cardiometabolic subgroups.

Previous studies have demonstrated associations between short-term NO₂ exposure and an increased risk of ischemic stroke, supporting our findings^{9,22}. A 2023 meta-analysis, for example, reported a 1.4% increase in stroke mortality and hospitalizations per 10 ppb rise in NO₂⁶. A recent multi-country analysis of 7.4 million deaths reported stronger short-term associations among older adults²³. That study focuses on fatal outcomes, while our study examines incident ischemic stroke and TIA; differences in (mortality vs incidence), age cut-points, exposure profiles, and healthcare systems may explain why age-related susceptibility is different between studies. Compared with prior studies, the effects observed in our analysis are similar in direction and magnitude and may appear larger in part because of differences in exposure assessment, spatial resolution, or population susceptibility.

Most existing research on air pollution and stroke has focused on PM_{2.5}, which can originate from various sources, including natural atmospheric reactions in the atmosphere²⁴. In a recent study conducted in Israel using the same data source of stroke cases between 2014-2018, researchers found elevated ischemic stroke risks associated with PM_{2.5} exposure lower than the national threshold²⁵. Our findings indicate that the NO₂ and Stroke association was independent of PM_{2.5} and temperature, suggesting that NO₂ may contribute to distinct health risks beyond PM_{2.5}. Nevertheless, NO₂ can be a marker for a broader mixture of traffic emissions, and residual confounding by unmeasured pollutants cannot be completely excluded.

A key finding of our study is the differential susceptibility observed across population groups. We found evidence of effect modification by urbanicity, sex, and age, with somewhat stronger cumulative associations among residents of large cities, women, and younger adults. These patterns are consistent with existing epidemiological studies showing that individuals in certain sociodemographic groups may be more susceptible to the adverse effects of air pollution^{23,26,27}. Urbanicity may reflect differences in exposure intensity, co-exposures (heat, noise), as well as socioeconomic and environmental factors that may amplify NO₂ effects^{28,29}.

The sex- and age-specific patterns were notable, with stronger and delayed associations between NO₂ exposure and ischemic stroke/TIA observed among women and younger adults. Lag-specific analyses indicated that the risk increased among women, particularly at lags of 3–5 days, compared with men. Although men often have higher age-specific stroke incidence, women experience a greater lifetime stroke burden and may be more vulnerable to vascular risk factors³⁰⁻³². In addition, women are exposed to sex hormone–related risk factors for stroke, including age at menarche, menopause, pregnancy-related complications, and hormonal therapies³¹. The stronger associations we observed among younger adults were consistent with a previous study from Israel reporting disproportionately stronger short-term associations between ambient air pollution and ischemic stroke among younger adults, despite their lower baseline stroke risk³³. In that study, the authors suggested that air pollution–related vascular injury may exert greater relative effects in younger individuals with fewer competing risk factors, potentially mediated through endothelial dysfunction, inflammatory responses, and prothrombotic pathways³³. This proposed mechanism aligns with our observed delayed effects at lags of 3–5 days and supports heightened susceptibility among women and younger adults.

We also observed slightly higher odds among individuals with diabetes compared to those without, although the test for interaction was not statistically significant. This is consistent with previous studies suggesting that underlying cardiometabolic dysfunction, with diabetes, may intensify the effects of NO₂ exposure, adding to greater cardiovascular risk from air pollution^{34,35}.

In addition to the cumulative association, we observed that the risk of ischemic stroke and TIA peaked 3–5 days after NO₂ exposure, with particularly pronounced delayed effects among women and younger adults. This pattern is consistent with studies that suggest exposure to air pollution can trigger pulmonary and systemic inflammation, pro-coagulant changes, and vascular dysfunction, which typically evolve over hours to several days after exposure³⁶. Several mechanisms could explain these patterns. While our data cannot directly account for all the pathways, the observed heterogeneity by sex and age suggests that short-term NO₂ effects disproportionately affect specific demographic groups. These highlight the importance of considering sex and age groups in air pollution and stroke research.

Strengths of our study include high spatiotemporally resolved exposure assessment, population-based setup, rich outcome, and covariate data. The case-crossover design inherently accounts for confounders that do not vary within a short period. However, some limitations remain. Although numerous confounders were accounted for by design, residual confounding from simultaneous environmental exposures is still possible. Considering this, we adjusted the models for PM_{2.5} and temperature. We observed no significant change in the results, reaffirming that NO₂ is independently associated with stroke. Another key limitation is that NO₂ exposure levels were only linked to residential addresses, not accounting for daily mobility. Since the NO₂ exposure levels may differ at their workplaces or other locations, this could lead to exposure misclassification⁴⁰. Nevertheless, this would be considered a non-

differential misclassification, resulting in bias towards the null, likely underestimating the true association between NO₂ exposure and stroke⁴¹.

In conclusion, we found an increased risk of ischemic stroke and TIA associated with higher NO₂ exposure. The peak effects were observed 3-5 days after exposure, and there was evidence of stronger effects in urban residents, women, and younger adults. The findings support NO₂ as an acute, modifiable risk factor for ischemic stroke/TIA, suggesting the need to develop public health interventions to reduce NO₂ exposure at the population level.

Abbreviations

NO ₂	Nitrogen Dioxide
TIA	Transient Ischemic Attack
PM _{2.5}	Particulate Matter less than 2.5 microns
DLNM	Distributed Lag Nonlinear Model
IHD	Ischemic Heart Disease
CVD	Cardiovascular Disease
INSR	Israeli National Stroke Registry
AOD	Aerosol Optical Depth
MAIAC	Multiangle Implementation of Atmospheric Correction
ICD-9	International Classification of Diseases, Ninth Revision
OR	Odds Ratio
CI	Confidence Interval
SD	Standard Deviation
IQR	Interquartile Range

Declarations

This study was approved by the institutional review boards at Soroka University Medical Center in Beer Sheva, Israel (study 01718-19), and the Icahn School of Medicine at Mount Sinai, New York, USA (study 21-00076). A waiver of informed consent was granted due to the minimal risk nature of the study and the use of de-identified registry data with no direct participant contact. All methods were performed in accordance with the relevant institutional and national guidelines and regulations.

Funding

This work was supported by the United States–Israel Binational Science Foundation (BSF), grant number 2021197

Clinical trial number: Not applicable

References

1. Pacheco-Barrios, K. *et al.* Burden of Stroke and Population-Attributable Fractions of Risk Factors in Latin America and the Caribbean. *J Am Heart Assoc* **11**, e027044, doi:10.1161/JAHA.122.027044 (2022).
2. Saposnik, G. *et al.* The World Stroke Academy: A World Stroke Organization global pathway to improve knowledge in stroke care. *Int J Stroke* **17**, 829-834, doi:10.1177/17474930221085895 (2022).
3. de Bont, J. *et al.* Mixtures of long-term exposure to ambient air pollution, built environment and temperature and stroke incidence across Europe. *Environ Int* **179**, 108136, doi:10.1016/j.envint.2023.108136 (2023).
4. Lee, K. K., Miller, M. R. & Shah, A. S. V. Air Pollution and Stroke. *J Stroke* **20**, 2-11, doi:10.5853/jos.2017.02894 (2018).
5. Organization, W. H. *Ambient (outdoor) air pollution*, <[https://www.who.int/news-room/fact-sheets/detail/ambient-\(outdoor\)-air-quality-and-health](https://www.who.int/news-room/fact-sheets/detail/ambient-(outdoor)-air-quality-and-health)> (
6. Kulick, E. R., Kaufman, J. D. & Sack, C. Ambient Air Pollution and Stroke: An Updated Review. *Stroke* **54**, 882-893, doi:10.1161/strokeaha.122.035498 (2023).
7. Shah, A. S. *et al.* Short term exposure to air pollution and stroke: systematic review and meta-analysis. *Bmj* **350**, h1295, doi:10.1136/bmj.h1295 (2015).
8. Collaborators, G. B. D. R. F. Global, regional, and national comparative risk assessment of 79 behavioural, environmental and occupational, and metabolic risks or clusters of risks, 1990-2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* **388**, 1659-1724, doi:10.1016/S0140-6736(16)31679-8 (2016).
9. Chen, Z. *et al.* Short-Term Effects of Low-Level Ambient Air NO₂ on the Risk of Incident Stroke in Enshi City, China. *Int J Environ Res Public Health* **19**, doi:10.3390/ijerph19116683 (2022).
10. Agency, U. S. E. P. *Nitrogen Dioxide (NO₂) Pollution*, <<https://www.epa.gov/no2-pollution/basic-information-about-no2>> (2024).
11. Block, M. L. & Calderon-Garciduenas, L. Air pollution: mechanisms of neuroinflammation and CNS disease. *Trends Neurosci* **32**, 506-516, doi:10.1016/j.tins.2009.05.009 (2009).
12. *Geography of Israel*, <<https://bje.org.au/knowledge-centre/israel/geography-of-israel/>> (2021).
13. Mike Zhongyu He, M. Y. S., Itai Kloog. A hybrid spatiotemporal model for estimating daily NO₂ concentrations across Israel. *ISEE Conference Abstracts* **Volume 2023**, doi:<https://doi.org/10.1289/isee.2023.MP-011> (2023).

14. He, M. Z. *et al.* Predicting fine-scale daily NO₂ over Mexico City using an ensemble modeling approach. *Atmos Pollut Res* **14**, doi:10.1016/j.apr.2023.101763 (2023).
15. Gutierrez-Avila, I. *et al.* Prediction of daily mean and one-hour maximum PM_{2.5} concentrations and applications in Central Mexico using satellite-based machine-learning models. *J Expo Sci Environ Epidemiol* **32**, 917-925, doi:10.1038/s41370-022-00471-4 (2022).
16. Gutierrez-Avila, I. *et al.* A spatiotemporal reconstruction of daily ambient temperature using satellite data in the Megalopolis of Central Mexico from 2003 to 2019. *Int J Climatol* **41**, 4095-4111, doi:10.1002/joc.7060 (2021).
17. Just, A. C. *et al.* Using High-Resolution Satellite Aerosol Optical Depth To Estimate Daily PM_{2.5} Geographical Distribution in Mexico City. *Environ Sci Technol* **49**, 8576-8584, doi:10.1021/acs.est.5b00859 (2015).
18. Kloog, I. *et al.* Estimating daily PM_{2.5} and PM₁₀ across the complex geo-climate region of Israel using MAIAC satellite-based AOD data. *Atmos Environ (1994)* **122**, 409-416, doi:10.1016/j.atmosenv.2015.10.004 (2015).
19. Rosenfeld, A. *et al.* Estimating daily minimum, maximum, and mean near surface air temperature using hybrid satellite models across Israel. *Environ Res* **159**, 297-312, doi:10.1016/j.envres.2017.08.017 (2017).
20. Gasparrini, A. Modeling exposure-lag-response associations with distributed lag non-linear models. *Stat Med* **33**, 881-899, doi:10.1002/sim.5963 (2014).
21. Yitshak Sade, M., Novack, V., Ifergane, G., Horev, A. & Kloog, I. Air Pollution and Ischemic Stroke Among Young Adults. *Stroke; a journal of cerebral circulation* **46**, 3348-3353, doi:10.1161/strokeaha.115.010992 (2015).
22. Ho, A. F. W. *et al.* Association of air pollution with acute ischemic stroke risk in Singapore: a time-stratified case-crossover study. *Int J Stroke* **17**, 983-989, doi:10.1177/17474930211066745 (2022).
23. Marb, A. *et al.* Short-term exposure to ambient nitrogen dioxide and fine particulate matter and cause-specific mortality: A causal modeling approach in four regions. *Environ Pollut* **372**, 126059, doi:10.1016/j.envpol.2025.126059 (2025).
24. Tessum, M. W. *et al.* Sources of ambient PM_{2.5} exposure in 96 global cities. *Atmos Environ (1994)* **286**, 119234, doi:10.1016/j.atmosenv.2022.119234 (2022).
25. Gaines, B. *et al.* Particulate Air Pollution Exposure and Stroke among Adults in Israel. *Int J Environ Res Public Health* **20**, doi:10.3390/ijerph20021482 (2023).
26. Zhang, J. *et al.* Effects of air pollution on cardiovascular health in patients with type 2 diabetes mellitus: Evidence from a large tertiary hospital in Shandong Province, China. *Front Public Health* **10**, 1050676, doi:10.3389/fpubh.2022.1050676 (2022).
27. Rajagopalan, S., Al-Kindi, S. G. & Brook, R. D. Air Pollution and Cardiovascular Disease: JACC State-of-the-Art Review. *J Am Coll Cardiol* **72**, 2054-2070, doi:10.1016/j.jacc.2018.07.099 (2018).
28. Zhang, G. *et al.* Rural-urban differences in associations between air pollution and cardiovascular hospital admissions in Guangxi, southwest China. *Environ Sci Pollut Res Int* **29**, 40711-40723,

- doi:10.1007/s11356-021-18196-6 (2022).
29. Anenberg, S. C. et al. Long-term trends in urban NO₂ concentrations and associated paediatric asthma incidence: estimates from global datasets. *Lancet Planet Health* **6**, e49-e58, doi:10.1016/S2542-5196(21)00255-2 (2022).
 30. Hanna, M., Wabnitz, A. & Grewal, P. Sex and stroke risk factors: A review of differences and impact. *J Stroke Cerebrovasc Dis* **33**, 107624, doi:10.1016/j.jstrokecerebrovasdis.2024.107624 (2024).
 31. Leppert, M. H. et al. Systematic Review of Sex Differences in Ischemic Strokes Among Young Adults: Are Young Women Disproportionately at Risk? *Stroke* **53**, 319-327, doi:10.1161/STROKEAHA.121.037117 (2022).
 32. Ospel, J., Singh, N., Ganesh, A. & Goyal, M. Sex and Gender Differences in Stroke and Their Practical Implications in Acute Care. *J Stroke* **25**, 16-25, doi:10.5853/jos.2022.04077 (2023).
 33. Yitshak Sade, M., Novack, V., Ifergane, G., Horev, A. & Kloog, I. Air Pollution and Ischemic Stroke Among Young Adults. *Stroke* **46**, 3348-3353, doi:10.1161/STROKEAHA.115.010992 (2015).
 34. Niemann, B. et al. Oxidative Stress and Cardiovascular Risk: Obesity, Diabetes, Smoking, and Pollution: Part 3 of a 3-Part Series. *J Am Coll Cardiol* **70**, 230-251, doi:10.1016/j.jacc.2017.05.043 (2017).
 35. Bonanni, L. J., Wittkopp, S., Long, C., Aleman, J. O. & Newman, J. D. A review of air pollution as a driver of cardiovascular disease risk across the diabetes spectrum. *Front Endocrinol (Lausanne)* **15**, 1321323, doi:10.3389/fendo.2024.1321323 (2024).
 36. Maheswaran, R. Air pollution and stroke - an overview of the evidence base. *Spat Spatiotemporal Epidemiol* **18**, 74-81, doi:10.1016/j.sste.2016.04.004 (2016).

Figures




Figure 1

The association between NO_2 exposure in lag days 0-5 and ischemic stroke or TIA.

Panel A shows the cumulative NO_2 -stroke/TIA association (solid line) with 95% CIs (dashed). Panel B shows lag-specific odds ratios (lags 0-5 days) with 95% CIs. Estimates were derived from distributed lag nonlinear models adjusted for $\text{PM}_{2.5}$ and temperature; the horizontal line indicates $\text{OR} = 1.0$.




Figure 2

Lag-specific associations between short-term NO₂ exposure and ischemic stroke or TIA, stratified by urbanicity, sex, and age.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [TableS1.docx](#)