



# wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 4, 2025 – 04:54 PM JST

PDB ID : 9WD3 / pdb\_00009wd3  
Title : Crystal structure of HPK1 kinase in complex with a potent and selective inhibitor with in vivo efficacy  
Deposited on : 2025-08-18  
Resolution : 3.22 Å(reported)

**This wwPDB validation report is for manuscript review**

This is a wwPDB X-ray Structure Validation Summary Report.

This report is produced by the wwPDB biocuration pipeline after annotation of the structure.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0rc1
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	2.0rc1
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.006 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.22 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	1638 (3.24-3.20)
Clashscore	180529	1778 (3.24-3.20)
Ramachandran outliers	177936	1751 (3.24-3.20)
Sidechain outliers	177891	1750 (3.24-3.20)
RSRZ outliers	164620	1639 (3.24-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	299	
1	B	299	
1	C	299	
1	D	299	

*Continued on next page...*

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
 Validation Pipeline (wwPDB-VP) : 2.45.1

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E	299	
1	F	299	
1	G	299	
1	H	299	

For Manuscript Review

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 18304 atoms, of which 290 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mitogen-activated protein kinase kinase kinase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	288	2249	1448	387	403	11	0	0	0
1	B	272	2145	1380	366	388	11	0	0	0
1	C	281	2197	1417	380	390	10	0	0	0
1	D	288	2260	1455	389	405	11	0	0	0
1	E	284	2223	1431	384	397	11	0	0	0
1	F	274	2156	1391	368	386	11	0	0	0
1	G	275	2156	1388	370	387	11	0	0	0
1	H	287	2248	1448	387	402	11	0	0	0

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP Q92918
A	1	PRO	-	expression tag	UNP Q92918
A	171	ALA	SER	engineered mutation	UNP Q92918
B	0	GLY	-	expression tag	UNP Q92918
B	1	PRO	-	expression tag	UNP Q92918
B	171	ALA	SER	engineered mutation	UNP Q92918
C	0	GLY	-	expression tag	UNP Q92918
C	1	PRO	-	expression tag	UNP Q92918
C	171	ALA	SER	engineered mutation	UNP Q92918
D	0	GLY	-	expression tag	UNP Q92918
D	1	PRO	-	expression tag	UNP Q92918
D	171	ALA	SER	engineered mutation	UNP Q92918
E	0	GLY	-	expression tag	UNP Q92918

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	1	PRO	-	expression tag	UNP Q92918
E	171	ALA	SER	engineered mutation	UNP Q92918
F	0	GLY	-	expression tag	UNP Q92918
F	1	PRO	-	expression tag	UNP Q92918
F	171	ALA	SER	engineered mutation	UNP Q92918
G	0	GLY	-	expression tag	UNP Q92918
G	1	PRO	-	expression tag	UNP Q92918
G	171	ALA	SER	engineered mutation	UNP Q92918
H	0	GLY	-	expression tag	UNP Q92918
H	1	PRO	-	expression tag	UNP Q92918
H	171	ALA	SER	engineered mutation	UNP Q92918

- Molecule 2 is 5-(8-methoxyimidazo[1,2-a]pyridin-3-yl)-7,7-dimethyl-9-morpholin-4-yl-2,12-dihydrobenzo[b][1,9]phenanthroline-1-one (CCD ID: A1EVW) (formula: C<sub>30</sub>H<sub>29</sub>N<sub>5</sub>O<sub>3</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	H	N			O
2	A	1	Total	C	H	N	O	0	0
			67	30	29	5	3		
2	B	1	Total	C	H	N	O	0	0
			67	30	29	5	3		
2	C	1	Total	C	H	N	O	0	0
			67	30	29	5	3		
2	C	1	Total	C	H	N	O	0	0
			67	30	29	5	3		
2	D	1	Total	C	H	N	O	0	0
			67	30	29	5	3		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
2	D	1	Total 67	C 30	H 29	N 5	O 3	0	0
2	E	1	Total 67	C 30	H 29	N 5	O 3	0	0
2	F	1	Total 67	C 30	H 29	N 5	O 3	0	0
2	G	1	Total 67	C 30	H 29	N 5	O 3	0	0
2	H	1	Total 67	C 30	H 29	N 5	O 3	0	0

For Manuscript Review


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Mitogen-activated protein kinase kinase kinase kinase 1




- Molecule 1: Mitogen-activated protein kinase kinase kinase kinase 1




- Molecule 1: Mitogen-activated protein kinase kinase kinase kinase 1









● Molecule 1: Mitogen-activated protein kinase kinase kinase kinase 1



● Molecule 1: Mitogen-activated protein kinase kinase kinase kinase 1



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	116.62Å 99.60Å 117.00Å 90.00° 90.44° 90.00°	Depositor
Resolution (Å)	22.78 – 3.22 22.78 – 3.22	Depositor EDS
% Data completeness (in resolution range)	92.6 (22.78-3.22) 92.5 (22.78-3.22)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.78 (at 3.23Å)	Xtriage
Refinement program	PHENIX 1.17.1_3660, PHENIX 1.17.1_3660	Depositor
R, $R_{free}$	0.252 , 0.308 0.250 , 0.302	Depositor DCC
$R_{free}$ test set	2096 reflections (4.72%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	76.9	Xtriage
Anisotropy	0.085	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 90.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.000 for l,k,-h 0.000 for h,-k,-l 0.000 for l,-k,h	Xtriage
$F_o, F_c$ correlation	0.88	EDS
Total number of atoms	18304	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	81.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 23.27 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.8229e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: A1EVW

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.29	0/2298	0.59	0/3112
1	B	0.35	0/2191	0.61	0/2964
1	C	0.23	0/2243	0.47	0/3035
1	D	0.47	0/2308	0.68	0/3122
1	E	0.31	0/2269	0.56	0/3070
1	F	0.32	0/2202	0.64	0/2977
1	G	0.34	0/2201	0.61	0/2975
1	H	0.27	0/2296	0.52	0/3106
All	All	0.33	0/18008	0.59	0/24361

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2249	0	2285	74	0
1	B	2145	0	2177	86	0
1	C	2197	0	2244	82	0
1	D	2260	0	2304	123	0
1	E	2223	0	2271	111	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	2156	0	2191	125	0
1	G	2156	0	2188	86	0
1	H	2248	0	2290	74	0
2	A	38	29	0	14	0
2	B	38	29	0	10	0
2	C	76	58	0	21	0
2	D	76	58	0	19	0
2	E	38	29	0	5	0
2	F	38	29	0	9	0
2	G	38	29	0	4	0
2	H	38	29	0	9	0
All	All	18014	290	17950	706	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 706 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:106:THR:CG2	1:B:291:LEU:HD23	1.64	1.26
1:H:185:ALA:HB1	1:H:188:LEU:CG	1.73	1.17
1:E:161:GLN:HG3	1:E:166:LEU:HD13	1.22	1.15
1:E:106:THR:HG22	1:E:291:LEU:HD21	1.18	1.12
1:F:106:THR:HG22	1:F:291:LEU:HD21	1.29	1.10

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	286/299 (96%)	250 (87%)	29 (10%)	7 (2%)	<b>5</b> <b>27</b>

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	266/299 (89%)	249 (94%)	15 (6%)	2 (1%)	16	50
1	C	277/299 (93%)	253 (91%)	22 (8%)	2 (1%)	19	53
1	D	284/299 (95%)	255 (90%)	25 (9%)	4 (1%)	9	39
1	E	280/299 (94%)	250 (89%)	25 (9%)	5 (2%)	7	34
1	F	268/299 (90%)	247 (92%)	15 (6%)	6 (2%)	5	29
1	G	269/299 (90%)	252 (94%)	13 (5%)	4 (2%)	8	37
1	H	283/299 (95%)	257 (91%)	25 (9%)	1 (0%)	30	63
All	All	2213/2392 (92%)	2013 (91%)	169 (8%)	31 (1%)	9	39

5 of 31 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	53	ASP
1	A	55	ASP
1	A	56	VAL
1	A	186	VAL
1	D	54	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	239/252 (95%)	236 (99%)	3 (1%)	65	81
1	B	233/252 (92%)	227 (97%)	6 (3%)	41	68
1	C	233/252 (92%)	232 (100%)	1 (0%)	89	93
1	D	242/252 (96%)	237 (98%)	5 (2%)	48	72
1	E	237/252 (94%)	237 (100%)	0	100	100
1	F	232/252 (92%)	231 (100%)	1 (0%)	89	93
1	G	231/252 (92%)	231 (100%)	0	100	100
1	H	240/252 (95%)	239 (100%)	1 (0%)	89	93
All	All	1887/2016 (94%)	1870 (99%)	17 (1%)	75	87

5 of 17 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	186	VAL
1	H	186	VAL
1	B	63	ILE
1	B	64	LEU
1	C	170	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	212	GLN
1	G	161	GLN
1	H	248	HIS
1	H	212	GLN
1	F	233	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	A1EVW	B	301	-	41,44,44	4.62	20 (48%)	53,67,67	2.66	20 (37%)
2	A1EVW	F	301	-	41,44,44	4.62	20 (48%)	53,67,67	2.72	20 (37%)
2	A1EVW	A	301	-	41,44,44	4.63	20 (48%)	53,67,67	2.59	23 (43%)
2	A1EVW	G	301	-	41,44,44	4.37	19 (46%)	53,67,67	2.45	24 (45%)
2	A1EVW	C	301	-	41,44,44	4.61	20 (48%)	53,67,67	2.60	24 (45%)
2	A1EVW	E	301	-	41,44,44	4.34	18 (43%)	53,67,67	2.56	20 (37%)
2	A1EVW	D	301	-	41,44,44	4.46	20 (48%)	53,67,67	2.65	24 (45%)
2	A1EVW	H	301	-	41,44,44	4.64	20 (48%)	53,67,67	2.77	24 (45%)
2	A1EVW	C	302	-	41,44,44	4.57	22 (53%)	53,67,67	2.67	22 (41%)
2	A1EVW	D	302	-	41,44,44	4.75	22 (53%)	53,67,67	2.36	19 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A1EVW	B	301	-	-	5/10/34/34	0/7/7/7
2	A1EVW	F	301	-	-	6/10/34/34	0/7/7/7
2	A1EVW	A	301	-	-	3/10/34/34	0/7/7/7
2	A1EVW	G	301	-	-	5/10/34/34	0/7/7/7
2	A1EVW	C	301	-	-	3/10/34/34	0/7/7/7
2	A1EVW	E	301	-	-	3/10/34/34	0/7/7/7
2	A1EVW	D	301	-	-	7/10/34/34	0/7/7/7
2	A1EVW	H	301	-	-	7/10/34/34	0/7/7/7
2	A1EVW	C	302	-	-	2/10/34/34	0/7/7/7
2	A1EVW	D	302	-	-	4/10/34/34	0/7/7/7

The worst 5 of 201 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	301	A1EVW	C21-C20	-12.67	1.20	1.39
2	B	301	A1EVW	C21-C20	-12.20	1.21	1.39
2	H	301	A1EVW	C24-N04	12.13	1.61	1.36
2	C	302	A1EVW	C21-C20	-12.13	1.21	1.39
2	A	301	A1EVW	C21-C20	-12.08	1.21	1.39

The worst 5 of 220 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	301	A1EVW	C22-C21-C20	7.38	135.75	119.62
2	F	301	A1EVW	C22-C21-C20	7.38	135.75	119.62
2	B	301	A1EVW	C18-C17-C16	-7.30	96.21	109.65
2	C	302	A1EVW	C22-C21-C20	7.19	135.33	119.62
2	A	301	A1EVW	C22-C21-C20	7.12	135.18	119.62

There are no chirality outliers.

5 of 45 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	A	301	A1EVW	C21-C22-C23-N05
2	B	301	A1EVW	C21-C22-C23-N05
2	C	301	A1EVW	C21-C22-C23-N05
2	C	302	A1EVW	C25-C29-O03-C30
2	D	301	A1EVW	C25-C29-O03-C30

There are no ring outliers.



10 monomers are involved in 91 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	301	A1EVW	10	0
2	F	301	A1EVW	9	0
2	A	301	A1EVW	14	0
2	G	301	A1EVW	4	0
2	C	301	A1EVW	11	0
2	E	301	A1EVW	5	0
2	D	301	A1EVW	7	0
2	H	301	A1EVW	9	0
2	C	302	A1EVW	10	0
2	D	302	A1EVW	12	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





For Manuscript Review




For Manuscript



For Manuscript Review



For Manuscript



## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	288/299 (96%)	1.01	33 (11%) 11 8	82, 127, 159, 182	0
1	B	272/299 (90%)	0.02	5 (1%) 67 52	33, 65, 103, 122	0
1	C	281/299 (93%)	0.51	14 (4%) 35 24	57, 90, 125, 145	0
1	D	288/299 (96%)	0.18	6 (2%) 63 48	38, 70, 103, 122	0
1	E	284/299 (94%)	0.32	19 (6%) 25 18	42, 75, 130, 163	0
1	F	274/299 (91%)	-0.01	5 (1%) 67 52	33, 55, 94, 108	0
1	G	275/299 (91%)	-0.08	6 (2%) 62 46	30, 54, 93, 121	0
1	H	287/299 (95%)	0.46	22 (7%) 21 15	44, 88, 134, 168	0
All	All	2249/2392 (94%)	0.31	110 (4%) 36 25	30, 76, 137, 182	0

The worst 5 of 110 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	258	SER	5.2
1	A	220	PRO	4.8
1	H	53	ASP	4.4
1	A	159	SER	4.3
1	C	186	VAL	3.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [\(i\)](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	A1EVW	C	302	38/38	0.74	0.20	129,156,188,200	0
2	A1EVW	D	302	38/38	0.75	0.21	129,177,323,337	0
2	A1EVW	A	301	38/38	0.76	0.20	81,106,133,141	0
2	A1EVW	E	301	38/38	0.77	0.17	58,77,103,126	0
2	A1EVW	H	301	38/38	0.81	0.17	64,90,109,129	0
2	A1EVW	D	301	38/38	0.82	0.15	51,80,105,115	0
2	A1EVW	G	301	38/38	0.84	0.16	40,72,94,119	0
2	A1EVW	C	301	38/38	0.84	0.14	55,86,106,117	0
2	A1EVW	B	301	38/38	0.85	0.15	47,75,97,117	0
2	A1EVW	F	301	38/38	0.85	0.15	36,68,97,117	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




### Electron density around A1EVW C 302:

2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
 mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
 and green (positive)






**Electron density around A1EVW D 302:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
 and green (positive)






**Electron density around A1EVW A 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
 and green (positive)






**Electron density around A1EVW E 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around A1EVW H 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around A1EVW D 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
 and green (positive)






**Electron density around A1EVW G 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
 and green (positive)






**Electron density around A1EVW C 301:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around A1EVW B 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.

For Manuscript Review