

1 **Supplementary information**

2

3 **Multi-Omics Mapping of Human Kidney Reveals Complement-Mediated Cellular**
4 **Dynamics During Progression of Focal Segmental Glomerulosclerosis**

5

6

7 Sahomi Hayashi, MD, PhD^{1,2}, Miyoshi Takeuchi, MD^{1,2}, Toshiaki Nakano, MD, PhD^{1*#}, Daiki
8 Setoyama, PhD^{2*#}, Sasha A. Singh, PhD³, Abhijeet R. Sonawane, PhD³, Takaki Iwamoto, MD¹, Hiroshi
9 Kishimoto, MD, PhD¹, Akihiro Tsuchimoto, MD, PhD¹, Shunsuke Yamada, MD, PhD¹, Dongchon
10 Kang, MD, PhD^{2,5}, Tetsuro Ago, MD, PhD^{1,6}, Takanari Kitazono, MD, PhD¹, Masanori Aikawa, MD,
11 PhD^{3,4}, Yuya Kunisaki, MD, PhD^{2,5#}

12

13 **Methods**

14 **LC-MS/MS data acquisition**

15 Full MS scans were acquired at a resolution of 70,000 (automatic gain control (AGC) target of $3e6$,
16 max injection time of 100 ms, m/z range 350-1500). MS/MS scans were conducted at a resolution of
17 17,500 (AGC target of $1e5$, max injection time of 50 ms, m/z range 200-2000), selecting the top 10
18 precursor ions with an isolation window of 2.0 m/z. Dynamic exclusion was set to auto, with a
19 minimum AGC target of $1.00e3$ and an intensity threshold of $2.0e4$. Precursor ions with charge states
20 of 1, 5, more than 5, or unassigned were excluded. Isotope exclusion was enabled. The spectra were
21 imported into Proteome Discoverer 2.5 (PD 2.5; Thermo Fisher Scientific) as follows: the four SDB-
22 SCX fractions per kidney biopsy sample were imported as 'fractions' thereby combining them into a
23 single query file; for a total of 73 queries that were analyzed against the Uniprot fasta database
24 (downloaded March 16, 2023; n = 42,326 entries) using the Sequest HT algorithm. Trypsin was set
25 as the digestion enzyme, allowing up to 2 missed cleavages and a minimum peptide length of 6
26 amino acids. Oxidation (+15.995 Da) of methionine; acetylation (+42.011 Da) of the N-terminus; the
27 loss of N-terminal methionine (-131.040 Da); and combining loss of methionine and addition of
28 acetylation at N-terminal methionine (-89.030 Da), were set as dynamic modifications. Spectral mass
29 tolerances were 10 ppm for the precursor and 20 mmu for the fragments.

30

31

32

33

34

35

36

37 **Supplementary table S1. The patients' background at kidney biopsy in the proteomics cohort.**

Total n = 73	FSGS n = 23	Donor n = 14	MCD n = 15	IgAN n = 21	P value
Sex (%)	a)		d)	e)	0.028
Female	8 (34.8)	10 (76.9)	4 (26.7)	7 (33.3)	
Male	15 (65.2)	3 (23.1)	11 (73.3)	14 (66.7)	
Age	63.00 [44.50, 71.00]	66.00 [45.00, 67.00]	70.00 [52.50, 72.50]	64.00 [39.00, 69.00]	0.549
BMI (kg/m ²)	24.60 [21.53, 27.17]	22.54 [21.25, 27.12]	25.09 [22.84, 28.69]	23.36 [21.70, 27.18]	0.64
Mean BP (mmHg)	100.00 [93.33, 105.17]	90.33 [81.50, 95.67]	90.00 [83.83, 99.17]	98.67 ^{e)} [94.00, 107.00]	0.012
TP (g/dl)	6.50 ^{b)} [5.50, 7.25]	7.00 [6.75, 7.15]	4.70 ^{d)} [4.30, 5.00]	6.90 ^{f)} [6.50, 7.20]	<0.001
Alb (g/dl)	3.80 ^{a)b)} [2.90, 4.15]	4.30 [4.10, 4.55]	1.70 ^{d)} [1.45, 2.05]	4.10 ^{f)} [3.50, 4.10]	<0.001
T.chol (g/dl)	231.00 ^{b)} [204.25, 280.50]	195.00 [164.50, 233.50]	356.00 ^{d)} [296.00, 469.00]	190.00 ^{f)} [171.50, 229.00]	<0.001
LDL (mg/dl)	146.70 ^{b)} [92.40, 183.05]	95.60 [90.00, 133.10]	214.00 ^{d)} [184.80, 363.20]	110.70 ^{f)} [80.15, 126.05]	<0.001
CRP (mg/dl)	0.05 [0.02, 0.22]	0.08 [0.03, 0.18]	0.08 [0.04, 0.10]	0.07 [0.02, 0.10]	0.947
Cr (mg/dl)	1.16 ^{a)} [1.05, 1.42]	0.62 [0.58, 0.67]	1.02 ^{d)} [0.84, 1.73]	1.08 ^{e)} [0.88, 1.52]	<0.001
eGFR (ml/min/1.73m ²)	44.94 ^{a)} [37.35, 55.19]	75.66 [70.69, 85.99]	53.98 ^{d)} [31.37, 80.86]	50.25 ^{e)} [40.70, 65.92]	0.001
UP/Ucr (g/gCr)	2.50 ^{a)b)} [1.48, 7.92]	0.06 [0.06, 0.07]	10.15 ^{d)} [8.15, 14.28]	0.98 ^{e) f)} [0.50, 2.72]	<0.001

38 Data are expressed as Median [interquartile range] or numbers (%).

39 Statistic test: Kruskal-Wallis test for numeric data, and Fisher's exact test for categorical variables.

40

41 P values indicate the results of the Kruskal-Wallis test or Fisher's exact test performed on the four

42 groups. P values less than 0.05 were considered statistically significant. If there are any significant

43 differences, multiple tests were performed by Bonferroni method. When any significant differences
44 are detected, following a), b), c), d), e) and f) are shown.

45 a) FSGS – Kidney Donor, b) FSGS – MCD, c) FSGS – IgAN, d) Kidney Donor – MCD, e) Kidney
46 Donor – IgAN, f) MCD – IgAN

47 FSGS: focal segmental glomerulosclerosis, MCD: minimal change disease, IgAN: IgA nephropathy,

48 BMI: body mass index, Mean BP: mean blood pressure, TP: serum total protein, Alb: serum albumin,

49 T.chol: serum total cholesterol, LDL: serum low density lipoprotein cholesterol, CRP: serum c reactive

50 protein, Cr: serum creatinine, eGFR: estimated glomerular filtration rate, UP/Ucr: urinary protein/
51 urinary creatinine ratio

52

53

54

55

56

57

58

59

60

61

62

63

64

65

66

67 **Supplementary table S2. The patients' background at kidney biopsy in the spatial**
68 **transcriptomics cohort.**

Total n = 21	FSGS n = 15	Donor n = 6	P value
Sex (%)			0.028
Female	6 (40.0)	4 (66.7)	
Male	9 (60.0)	2 (33.3)	
Age	60.00 [52.00, 74.00]	44.50 [38.00, 62.25]	0.129
BMI (kg/m ²)	23.60 [22.89, 27.72]	21.86 [19.92, 25.43]	0.096
Mean BP (mmHg)	103.33 [89.42, 108.83]	94.33 [89.08, 97.33]	0.386
TP (g/dl)	5.40 [4.85, 6.80]	7.30 [7.12, 7.47]	0.001
Alb (g/dl)	2.50 [2.05, 3.75]	4.40 [4.18, 4.40]	0.002
T.chol (g/dl)	283.00 [236.25, 325.00]	169.00 [166.50, 204.50]	0.015
LDL (mg/dl)	180.65 [142.50, 206.00]	93.50 [81.50, 119.00]	0.017
CRP (mg/dl)	0.05 [0.01, 0.18]	0.03 [0.02, 0.07]	0.557
Cr (mg/dl)	1.05 [0.80, 1.54]	0.64 [0.63, 0.84]	0.087
eGFR (ml/min/1.73m ²)	48.72 [31.22, 74.41]	74.50 [70.20, 80.83]	0.073
UP/Ucr (g/gCr)	6.41 [3.08, 10.86]	0.06 [0.05, 0.08]	<0.001

69 Data are expressed as Median [interquartile range] or numbers (%).

70 Statistic test: Mann-Whitney U test for numeric data, and Fisher's exact test for categorical variables

71 P-values less than 0.05 were considered statistically significant.

72

73 FSGS: focal segmental glomerulosclerosis, BMI: body mass index, Mean BP: mean blood pressure,
74 TP: serum total protein, Alb: serum albumin, T.chol: serum total cholesterol, LDL: serum low density
75 lipoprotein cholesterol, CRP: serum c reactive protein, Cr: serum creatinine, eGFR: estimated
76 glomerular filtration rate, UP/Ucr: urinary protein/ urinary creatinine ratio

77

78

79

80

81

82

83

84

85

86

87

88

89

90

91

92

93

94

95

96

97

98 **Supplementary table S3. Classification, treatment, and electron microscopic findings of FSGS**
 99 **groups used in the proteomics and spatial transcriptomics cohorts.**

	Proteomics cohort		Spatial cohort	
	n = 23		n = 15	
	Number of patients	%	Number of patients	%
Primary FSGS	9	39.1	10	66.7
Secondary FSGS	8	34.8	0	0.0
Undetermined cause	6	26.1	5	33.3
Steroid therapy	11	47.8	9	60.0
Immunosuppressants therapy	4	17.4	2	13.3
Foot process effacement				
Diffuse	5	21.7	8	53.3
Focal	4	17.4	3	20.0
Negative	0	0.0	1	6.7
Not available	14	60.9	3	20.0

100 Data are expressed as Number of patients or percentages (%).

101 FSGS: focal segmental glomerulosclerosis

102

103

104

105

106

107

108

109

110

111

112 **Supplementary table S4. Evaluation of pathological differences in the kidney biopsy samples**
 113 **among the four groups in proteomics cohort.**

Total n = 73	FSGS n = 23	Donor n = 14	MCD n = 15	IgAN n = 21	P value
Global sclerosis (%)	31.25 ^{a) b)} [13.10, 49.15]	6.70 [0.00, 10.44]	11.11 [6.38, 23.08]	17.39 ^{e)} [11.11, 26.32]	0.001
Segmental sclerosis (%)	9.09 ^{a) b) c)} [4.88, 17.42]	0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.00 [0.00, 4.35]	<0.001
Interstitial cell infiltration (%)	^{a)}			^{f) e)}	0.047
None	13 (56.5)	14 (100.0)	12 (80.0)	8 (38.1)	
Mild	7 (30.4)	0 (0.0)	3 (20.0)	10 (47.6)	
Moderate	2 (8.7)	0 (0.0)	0 (0.0)	2 (9.5)	
Severe	1 (4.3)	0 (0.0)	0 (0.0)	1 (4.8)	
Interstitial fibrosis (%)	^{a) b)}			^{e) f)}	<0.001
None	5 (21.7)	14 (100.0)	11 (73.3)	4 (19.0)	
Mild	13 (56.5)	0 (0.0)	3 (20.0)	10 (47.6)	
Moderate	4 (17.4)	0 (0.0)	1 (6.7)	7 (33.3)	
Severe	1 (4.3)	0 (0.0)	0 (0.0)	0 (0.0)	
IF C3	^{c)}			^{e) f)}	<0.001
(-)	22 (95.7)	12 (85.7)	15 (100.0)	5 (23.8)	
(+)	0 (0.0)	2 (14.3)	0 (0.0)	13 (61.9)	
(++)	1 (4.3)	0 (0.0)	0 (0.0)	3 (14.3)	
IF Fib	^{c)}			^{e) f)}	<0.001
(-)	21 (91.3)	14 (100.0)	15 (100.0)	10 (47.6)	
(+)	2 (8.7)	0 (0.0)	0 (0.0)	9 (42.9)	
(++)	0 (0.0)	0 (0.0)	0 (0.0)	2 (9.5)	
IF IgA	^{c)}			^{e) f)}	<0.001
(-)	20 (87.0)	10 (71.4)	14 (93.3)	0 (0.0)	
(+)	3 (13.0)	4 (28.6)	1 (6.7)	8 (38.1)	
(++)	0 (0.0)	0 (0.0)	0 (0.0)	9 (42.9)	
(+++)	0 (0.0)	0 (0.0)	0 (0.0)	4 (19.0)	
IF IgG					0.422
(-)	22 (95.7)	14 (100)	15 100.0()	19 (90.5)	

(+)	1 (4.3)	0 (0.0)	0 (0.0)	2 (9.5)	
IF IgM					0.311
(-)	18 (78.3)	14 (100.0)	13 (86.7)	17 (81.0)	
(+)	5 (21.7)	0 (0.0)	2 (13.3)	4 (19.0)	
IF kappa chain	^{c)}			^{e) f)}	<0.001
(-)	21 (91.3)	14 (100.0)	15 (100.0)	3 (14.3)	
(+)	2 (8.7)	0 (0.0)	0 (0.0)	15 (71.4)	
(++)	0 (0.0)	0 (0.0)	0 (0.0)	3 (14.3)	
IF lamda chain	^{c)}			^{e) f)}	<0.001
(-)	21 (91.3)	12 (85.7)	15 (100.0)	3 (14.3)	
(+)	2 (8.7)	2 (14.3)	0 (0.0)	14 (66.7)	
(++)	0 (0.0)	0 (0.0)	0 (0.0)	4 (19.0)	

114 Data are expressed as Median [interquartile range] or numbers (%).

115 Statistic test: Kruskal-Wallis test for numeric data, and Fisher's exact test for categorical variables.

116

117 P-values indicate the results of the Kruskal-Wallis test or Fisher's exact test performed on the four
 118 groups. P-values less than 0.05 were considered statistically significant. If there are any significant
 119 differences, multiple tests were performed by Bonferroni method. When any significant differences
 120 are detected, following a), b), c), d), e) and f) are shown.

121 a) FSGS - Kidney Donor, b) FSGS – MCD, c) FSGS – IgAN, d) Kidney Donor – MCD, e) Kidney
 122 Donor – IgAN, f) MCD – IgAN

123 Interstitial cell infiltration, interstitial fibrosis are defined as follows: None: <10% of cortical area,

124 Mild: 10–24%, Moderate: 25–49%, Severe: >50%.

125 FSGS: focal segmental glomerulosclerosis, MCD: minimal change disease, IgAN: IgA nephropathy

126

127

128

129 **Supplementary table S5. Evaluation of pathological differences in the kidney biopsy samples**
 130 **among the two groups in spatial transcriptomics cohort.**

Total n = 21	FSGS n = 15	Donor n = 6	P value
Global sclerosis (%)	15.79 [7.99, 45.69]	2.38 [0.00, 20.93]	0.086
Segmental sclerosis (%)	9.09 [3.24, 11.81]	0.00 [0.00, 0.00]	0.004
Interstitial cell infiltration (%)			0.156
None	8 (53.3)	6 (100.0)	
Mild	6 (40.0)	0 (0.0)	
Moderate	1 (6.7)	0 (0.0)	
Interstitial fibrosis (%)			0.29
None	5 (33.3)	5 (83.3)	
Mild	6 (40.0)	1 (16.7)	
Moderate	3 (20.0)	0 (0.0)	
Severe	1 (6.7)	0 (0.0)	
IF C3	°)		1
(-)	14 (93.3)	6 (100.0)	
(+)	1 (6.7)	0 (0.0)	
IF Fib			N/A
(-)	15 (100.0)	6 (100.0)	
IF IgA			0.167
(-)	12 (80.0)	5 (83.3)	0.167
(±)	3 (20.0)	0 (0.0)	
(+)	0 (0.0)	1 (16.7)	
IF IgG			N/A
(-)	15 (100.0)	6 (100.0)	
IF IgM			0.43
(-)	13 (86.7)	4 (66.7)	
(±)	0 (0.0)	1 (16.7)	
(+)	2 (13.3)	1 (16.7)	
IF kappa chain			N/A
(-)	15 (100.0)	6 (100.0)	

IF lamda chain

N/A

(-)

15 (100.0)

6 (100.0)

131 Data are expressed as Median [interquartile range] or numbers (%).

132 Statistic test: Kruskal-Wallis test for numeric data, and Fisher's exact test for categorical variables.

133 P-values indicate the results of the Kruskal-Wallis test or Fisher's exact test performed on the four

134 groups. P-values less than 0.05 were considered statistically significant. N/A indicates not applicable.

135 If there are any significant differences, multiple tests were performed by Bonferroni method. When

136 any significant differences are detected, following a), b), c), d), e) and f) are shown.

137 a) FSGS - Kidney Donor, b) FSGS – MCD, c) FSGS – IgAN, d) Kidney Donor – MCD, e) Kidney

138 Donor – IgAN, f) MCD – IgAN

139 Interstitial cell infiltration, interstitial fibrosis are defined as follows: None: <10% of cortical area,

140 Mild: 10–24%, Moderate: 25–49%, Severe: >50%.

141 FSGS: focal segmental glomerulosclerosis, MCD: minimal change disease, IgAN: IgA nephropathy