

**MONTREAL CLASSIFICATION (n=641)**

<b>Montreal A1 (%)</b>	85 (13.2%)
<b>Montreal A2 (%)</b>	469 (73.2%)
<b>Montreal A3 (%)</b>	87 (13.6%)
<b>Montreal L1 (%)</b>	254 (39.6%)
<b>Montreal L2 (%)</b>	136 (21.2%)
<b>Montreal L3 (%)</b>	303 (47.3%)
<b>Montreal L4 (%)</b>	46 (7.2%)
<b>Montreal B1 (%)</b>	11 (1.7%)
<b>Montreal B2 (%)</b>	528 (82.4%)
<b>Montreal B3 (%)</b>	414 (64.6%)
<b>Montreal p (%)</b>	74 (11.5%)

**POSTOPERATIVE COMPLICATIONS AS PER CLAVIEN-DINDO CLASSIFICATION (n=641)**

<b>0+1 (%)</b>	507 (79.1%)
<b>2 (%)</b>	89 (13.9%)
<b>3a (%)</b>	8 (1.2%)
<b>3b (%)</b>	29 (4.5%)
<b>4a (%)</b>	6 (0.9%)
<b>5 (%)</b>	2 (0.3%)

**Supplementary Table 1.** Montreal Classification and postoperative complications according to the Clavien-Dindo-Classification. Mean±SD