

Supplementary 1: Subgroup analysis of patients who had <3 months of anticoagulation

	Overall (n=369)	DOACs (n=170)	VKA (n=43)	LMWH (n=156)
Age, mean (SD)	62.78(16.42)	68.83(15.985)	58.72(18.44)	62.76(16.25)
Age groups(years)				
18-40	45	17	8	20
41-64	129	59	19	51
>65	195	94	16	85
Gender				
Male	178	81	20	77
Female	191	89	23	79
Ethnicity				
Malay	103	49	13	41
Chinese	159	68	16	75
Indian	83	44	9	30
Others	24	9	5	10
Anatomical site				
DVT	121	62	18	41
PE	211	94	21	96
DVT and PE	28	11	4	13
Splanchnic vein thrombosis	5	1	0	4
Renal vein thrombosis	1	1	0	0
Subclavian vein thrombosis	1	0	0	1
Multiple site thrombosis ≥2(other than PE and DVT)	2	1	0	1
Number of comorbidities				
0	121	60	14	47
≤2	198	89	22	87
≤3	50	21	7	22
Cause of VTE				
Unprovoked	29	19	4	6
Provoked	340	151	39	150
Malignancy	122	55	6	61
Recurrent VTE	9	6	0	3
Major Bleed	47	19	8	20
CRNMB	14	10	0	4
Minor bleed	6	5	0	1
Death	294	118	30	146
Cause of death				
VTE	42	9	1	32
Bleed secondary to anticoagulant	5	1	3	1
others	247	108	26	113
Time from diagnosis to death in weeks Mean(SD)	16.31(37.36)	23.47(46.97)	20.43(46.17)	9.63(22.75)