

Online Resource 5

		Overall n = 81	No HoLEP n = 54	HoLEP n = 27 n (%)	p
Pad use (/day) at 3 months postoperatively	0	23 (28.4)	16 (29.6)	7 (25.9)	0.32
	1	39 (48.1)	27 (50.0)	12 (44.4)	
	2	8 (9.9)	6 (11.1)	2 (7.4)	
	3 ≥	11 (13.5)	5 (9.3)	6 (22.2)	
Pad use (/day) at 6 months postoperatively	0	44 (54.3)	30 (55.6)	14 (51.9)	0.12
	1	28 (34.6)	21 (38.9)	7 (25.9)	
	2	5 (6.2)	2 (3.7)	3 (11.1)	
	3	4 (4.9)	1 (1.9)	3 (11.1)	
Pad use (/day) at 1 year postoperatively	0	57 (70.4)	39 (72.2)	18 (66.7)	0.032
	1	20 (24.7)	15 (27.8)	5 (18.5)	
	2	2 (2.5)	0 (0.0)	2 (7.4)	
	3	2 (2.5)	0 (0.0)	2 (7.4)	

Pad usage at 3, 6, and 12 months after RARP, stratified by prior HoLEP status

HoLEP: Holmium laser enucleation of the prostate