Research Square

Preprints are preliminary reports that have not undergone peer review.
They should not be considered conclusive, used to inform clinical practice,
or referenced by the media as validated information.

Navigating Complexity: An Analysis of Consumer Decision-Making in Modern Healthcare

Rupam Kumar Saha

rupamkumarsaha@gmail.com

NSHM Knowledge Campus, Durgapur, West Bengal, India. https://orcid.org/0009-0007-1727-6504

Sachin Sidhra

Lovely Professional University

Systematic Review

Keywords: Consumer Behaviour, Healthcare Decision-Making, Tertiary Care, Digital Health, Patient Empowerment

Posted Date: March 8th, 2026

DOI: https://doi.org/10.21203/rs.3.rs-9021460/v1

License: This work is licensed under a Creative Commons Attribution 4.0 International License.
Read Full License

Additional Declarations: The authors declare no competing interests.

Abstract

Background: The transition of healthcare from a paternalistic model to a consumer-driven marketplace has fundamentally altered how individuals make medical decisions. Cost considerations, quality perceptions, accessibility, health literacy, and cultural factors intersect to shape contemporary healthcare consumer behaviour, yet the mechanisms underlying these decisions remain inadequately understood, particularly in emerging economy contexts.

Objectives: This study investigates how patients in urban India navigate decisions when selecting medical services, with three specific objectives: (1) to identify the primary determinants of healthcare consumer decisions; (2) to examine how digital technologies—including online information, reviews, and telehealth—influence behaviour; and (3) to analyse barriers to informed decision-making and develop stakeholder recommendations.

Methods: A sequential mixed-methods design employing triangulation was utilised. Phase 1 comprised a systematic literature review (SLR) following PRISMA guidelines, synthesising 87 peer-reviewed studies from 2000–2024. Phase 2 involved multiple case studies of four tertiary care hospitals in eastern India, incorporating document analysis and observational data. Phase 3 encompassed semi-structured interviews with 42 participants: 24 patients (12 who had completed treatment decisions, 8 currently deciding, and 4 who declined recommended treatment), 8 healthcare providers, 6 hospital administrators, and 4 health insurance representatives. Thematic analysis was conducted using NVivo 14, with triangulation across methods, investigators, and data sources to enhance validity.

Findings: Consumer decision-making in healthcare operates at the intersection of rational information-seeking, emotional heuristics, and digital empowerment. Six overarching themes emerged: (1) the digitally augmented information seeker—consumers actively research online but struggle with source credibility; (2) trust as the foundational currency—provider endorsement remains the single most powerful influence; (3) price sensitivity amidst opacity—cost considerations are rising but transparency deficits persist; (4) emotional heuristics under uncertainty—anxiety and fear trigger decision shortcuts; (5) the convenience imperative—digital access strongly predicts provider choice; and (6) systemic barriers to informed choice—complexity, health literacy gaps, and administrative friction. Triangulation revealed convergence across patient, provider, and insurer perspectives on these themes.

Conclusions: While consumers are increasingly empowered through digital access, meaningful decision-making remains constrained by complexity, trust deficits, and transparency gaps. Healthcare stakeholders must embrace consumer-centric strategies that simplify choice, enhance cost transparency, leverage technology judiciously, and build authentic trust. This study contributes a comprehensive framework for understanding healthcare consumer behaviour in urban Indian contexts, with implications for theory, practice, and policy.

1. Introduction

1.1 Background: The Shifting Paradigm of Healthcare Consumption

Healthcare globally is transitioning from a paternalistic model—wherein patients passively received physician-directed care—to a consumer-driven marketplace characterised by activated individuals making consequential choices. This transformation is propelled by multiple forces: rising out-of-pocket expenditures, the proliferation of high-deductible health plans (where applicable), unprecedented access to online health information (OHI), and the diffusion of digital health tools that both empower and, paradoxically, overwhelm consumers .

In the Indian context, this shift assumes particular significance. India's healthcare ecosystem comprises a complex mix of public and private providers, with tertiary care increasingly concentrated in corporate hospital chains. The country's urban middle class—estimated at over 400 million—increasingly approaches healthcare with consumerist expectations shaped by experiences in retail, banking, and hospitality. Simultaneously, out-of-pocket expenditure accounts for approximately 62% of total health spending, rendering cost considerations salient. Yet, price transparency remains elusive, quality information is asymmetrically distributed, and health literacy varies widely across socioeconomic strata.

1.2 Problem Statement

Understanding how consumers navigate healthcare choices—selecting providers, treatments, insurance products, and digital tools—is crucial for improving access, satisfaction, clinical outcomes, and system efficiency. However, traditional marketing and consumer behaviour models, developed primarily for low-involvement product categories, often fail to capture healthcare's unique characteristics: the emotional weight of illness, the life-altering consequences of decisions, the profound information asymmetry between provider and patient, and the non-standardised nature of medical services.

Furthermore, extant research predominantly originates from Western developed markets, with limited empirical investigation of healthcare consumer behaviour in rapidly transforming economies like India. The interplay between traditional cultural norms (deference to physicians, family involvement in decisions) and emerging digital empowerment remains under theorised.

1.3 Research Objectives

This study addresses these gaps through the following objectives:

Primary Objective: To identify and analyse the principal determinants of healthcare consumer decisions in urban Indian tertiary care settings.

Secondary Objectives:

To examine how digital technologies—online information, reviews, telehealth, and health applications—influence consumer behaviour and decision processes.

To analyse the barriers that impede healthcare consumers from making informed, value-congruent decisions.

To develop evidence-based recommendations for healthcare stakeholders (providers, insurers, policymakers, technology developers) seeking to better meet consumer needs.

1.4 Scope and Delimitations

This study focuses on elective and planned care decisions in tertiary care hospitals within urban India. It encompasses decisions regarding provider selection, treatment choices (where alternatives exist), and engagement with digital health tools. Emergency care decisions, characterised by extreme time constraints and minimal deliberation, are excluded. While the empirical context is urban India, the conceptual framework and findings hold relevance for other emerging economies undergoing similar healthcare transitions.

1.5 Article Structure

Section 2 presents a systematic literature review, synthesising theoretical frameworks and empirical findings. Section 3 details the mixed-methods triangulation methodology. Section 4 reports findings organised by research objective, with integration across data sources. Section 5 discusses theoretical and practical implications. Section 6 outlines limitations and future research directions. Section 7 concludes.

2. Systematic Literature Review

2.1 Review Methodology

A systematic literature review was conducted following PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) guidelines. The review aimed to synthesise existing knowledge on healthcare consumer decision-making and identify gaps warranting empirical investigation.

Search Strategy: Five electronic databases were searched: PubMed, Scopus, Web of Science, EBSCOhost, and Google Scholar. The search covered January 2000 to May 2024, capturing the emergence and evolution of consumer-directed healthcare. Search strings combined terms related to ("healthcare consumer" OR "patient decision-making" OR "health consumer behaviour") AND ("determinants" OR "factors" OR "influences") AND ("digital health" OR "online information" OR "telehealth") AND ("barriers" OR "obstacles").

Inclusion Criteria: Peer-reviewed empirical studies (quantitative, qualitative, or mixed-methods) and systematic reviews; English language; focused on adult consumers making healthcare decisions; addressing determinants, digital influences, or decision barriers.

Exclusion Criteria: Editorials, commentaries, conference abstracts; studies focused solely on clinical decision-making by physicians; paediatric populations; non-English publications.

Quality Assessment: Methodological quality was appraised using the Mixed Methods Appraisal Tool (MMAT) for empirical studies and AMSTAR-2 for systematic reviews .

Synthesis Approach: Thematic synthesis was employed to identify patterns across studies, with findings organised around theoretical frameworks, influencing factors, digital transformations, and persistent barriers.

2.2 Search Results

The initial search yielded 1,847 records. After removing 412 duplicates, 1,435 titles and abstracts were screened, resulting in 312 full-text articles assessed for eligibility. Ultimately, 87 studies satisfied inclusion criteria and were incorporated into the synthesis. This comprised 54 quantitative studies (including 12 discrete choice experiments), 23 qualitative investigations, 7 mixed-methods studies, and 3 systematic reviews .

2.3 Defining the Healthcare Consumer

A foundational tension in the literature concerns the conceptualisation of the individual seeking medical care. Traditional discourse employs the term "patient," connoting a passive recipient of care, an ill person deserving of compassion but not necessarily an active agent in economic transactions. Contemporary scholarship increasingly deploys "consumer" or "healthcare consumer," reflecting the reality that individuals often bear significant financial responsibility, exercise choice among alternatives, and approach healthcare with expectations shaped by experiences in other service industries .

However, scholars caution against uncritical application of the consumer label. Hibbard and Greene distinguish between "patients" and "consumers" while emphasising that "patient activation"—the knowledge, skills, and confidence to manage one's health—exists on a continuum rather than as a binary state. The Health Belief Model (HBM) and Theory of Planned Behaviour (TPB), discussed below, provide frameworks for understanding when and how individuals act as engaged consumers versus distressed patients.

2.4 Theoretical Frameworks

Three theoretical frameworks dominate the literature on healthcare consumer behaviour, each offering complementary insights.

2.4.1 Health Belief Model (HBM)

Originally developed by Rosenstock in the 1950s to explain preventive health behaviour, the HBM posits that health-related action depends on six key constructs:

Perceived Susceptibility: Belief about the likelihood of experiencing a health threat.

Perceived Severity: Belief about the seriousness of the threat and its potential consequences.

Perceived Benefits: Belief in the efficacy of recommended action to reduce risk or severity.

Perceived Barriers: Belief about the tangible and psychological costs of recommended action.

Cues to Action: Triggers that prompt decision-making (symptoms, media reports, physician recommendations).

Self-Efficacy: Confidence in one's ability to successfully perform the action.

In the context of healthcare consumer decisions, the HBM illuminates why individuals with similar clinical profiles may make radically different choices. For instance, a patient perceiving high susceptibility to complications and strong benefits from surgery may proceed despite significant out-of-pocket costs, while another with identical financial circumstances but lower perceived severity may decline.

2.4.2 Theory of Planned Behaviour (TPB)

Ajzen's TPB extends the HBM by explicitly incorporating social and control factors. The theory posits that behavioural intention—the immediate antecedent of action—is determined by:

Attitude toward the behaviour: Positive or negative evaluation of performing the behaviour.

Subjective norm: Perceived social pressure to perform or not perform the behaviour.

Perceived behavioural control: Perceived ease or difficulty of performing the behaviour, reflecting past experience and anticipated obstacles.

In healthcare decisions, TPB captures the influence of family members (particularly salient in collectivist cultures like India), peer recommendations, and the perceived feasibility of navigating complex healthcare systems .

2.4.3 Consumer Decision-Making Models

The Engel-Kollat-Blackwell (EKB) model of consumer behaviour, adapted for healthcare contexts, conceptualises decision-making as a five-stage process: (1) need recognition, (2) information search, (3) evaluation of alternatives, (4) purchase decision, and (5) post-purchase evaluation. Recent empirical work confirms that healthcare consumers traverse these stages, though with modifications necessitated by the high-involvement, high-risk nature of medical decisions .

A significant refinement comes from research demonstrating that "seeking value" and "assessing value" constitute distinct sub-stages within the EKB framework. In a qualitative study of 54 insured adults, researchers found that consumers actively sought value through price comparisons, negotiations with providers, and responses to financial incentives—behaviours that map imperfectly onto traditional healthcare utilisation models .

2.5 Key Influencing Factors

Synthesis of the literature reveals five categories of factors shaping healthcare consumer decisions.

2.5.1 Trust and Reputation

Trust emerges consistently as the paramount factor in healthcare decisions. Unlike other service contexts where trust may be secondary to price or convenience, healthcare decisions hinge on confidence in the provider's competence, benevolence, and integrity. Trust is built through multiple channels: clinical reputation (often transmitted through professional networks), credentials and affiliations, personal recommendations from trusted social contacts, and increasingly, online ratings and reviews .

Provider endorsement functions as a particularly potent trust signal. Research on remote hypertension program enrollment found that "the decision to enroll hinges on provider endorsement"—patients consistently looked to their physicians for guidance on whether and how to engage with digital health innovations. This finding aligns with broader evidence that physician recommendations serve as heuristic substitutes for complex information processing when patients face uncertainty.

2.5.2 Accessibility and Convenience

Location proximity, appointment availability, and wait times significantly influence provider selection, particularly for non-emergency care. The convenience imperative extends to digital access: consumers increasingly expect online scheduling, telehealth options, electronic access to medical records, and patient portals for communication. Research demonstrates "widespread comfort with hybrid care delivery" among patients, with telehealth particularly valued for routine follow-ups and mental health consultations .

2.5.3 Cost and Financial Transparency

The rise of high-deductible health plans and out-of-pocket payment models has heightened price sensitivity among healthcare consumers. Studies document extensive price-seeking behaviours: 72% of participants sought cost information before receiving care, 56% compared provider prices, and 83% engaged in some form of negotiation regarding healthcare purchasing .

However, pervasive lack of price transparency remains a fundamental barrier to consumerism. Unlike other markets where prices are readily available pre-purchase, healthcare consumers often encounter prices only after services are rendered, rendering comparison shopping impossible. Even when prices are available, the complexity of insurance coverage—deductibles, co-payments, co-insurance, out-of-network penalties—obscures true out-of-pocket costs.

2.5.4 Quality and Outcomes Information

Consumers increasingly seek quality information to guide decisions, relying on online reviews, hospital rating systems (such as CMS Star Ratings in the US), physician scorecards, and word-of-mouth.

However, significant challenges attend the use of such information: questions about representativeness and validity, potential manipulation, and difficulty interpreting clinical quality metrics .

A systematic review of online health information (OHI) seeking found that the most frequent reasons for seeking OHI were to obtain diagnostic and treatment information, and to seek well-being and emotional support. Education level and socioeconomic status significantly influenced OHI-seeking behaviour, raising equity concerns .

2.5.5 Digital Engagement

Digital technologies are fundamentally reshaping healthcare consumer behaviour across multiple dimensions. Telehealth adoption, accelerated by the COVID-19 pandemic, has shifted expectations toward convenient, remote access for appropriate conditions. Patient portals and health applications enable consumers to access records, communicate with providers, and monitor health metrics. Wearable devices generate data that consumers increasingly bring to clinical encounters.

Research on continued telemedicine usage behaviour in emerging economies identifies multiple determinants: perceived usefulness, ease of use, social influence, facilitating conditions, and trust in technology. Importantly, technology adoption is not uniform across populations; age, digital literacy, and socioeconomic factors create differential patterns of engagement .

2.5.6 Emotional State

The emotional context of illness fundamentally shapes decision processing. Anxiety, fear, and urgency—hallmarks of healthcare decisions—impair systematic information processing and increase reliance on heuristics or decision shortcuts. Consumers facing high emotional arousal may choose the closest provider, follow a single recommendation without seeking alternatives, or defer decisions entirely to family members or physicians.

Research on sentiment in healthcare purchasing found that positive sentiment correlated with identifying options and evaluating decisions, while negative sentiment correlated with seeking value and making the purchase decision. This suggests that the emotionally charged phases of decision-making—confronting costs and committing to a course of action—may be particularly vulnerable to distress.

2.5.7 Socioeconomic and Cultural Factors

Income, education, health literacy, cultural beliefs, and social support networks profoundly shape healthcare decisions. Health literacy—the capacity to obtain, process, and understand basic health information needed to make appropriate health decisions—emerges as a critical determinant. Significant disparities in health literacy limit the ability of many consumers to make fully informed choices, exacerbating inequities in access and outcomes.

Cultural factors operate at multiple levels: family involvement in decisions (particularly pronounced in collectivist cultures), beliefs about illness causation and appropriate treatment, trust in biomedical

versus traditional systems, and communication preferences with providers .

2.6 Barriers to Informed Consumer Decisions

The literature identifies persistent barriers that impede truly informed consumer choice.

Complexity and Uncertainty: Medical jargon, intricate insurance terms, and inherently unpredictable outcomes create confusion and decision paralysis. Consumers frequently encounter information they cannot interpret, choices they cannot compare, and consequences they cannot anticipate.

Information Asymmetry: Despite increased access to online information, providers retain fundamental advantages in medical knowledge and experience. This asymmetry creates vulnerability and dependence, limiting the extent to which consumers can function as autonomous decision-makers.

Emotional Overload: As noted above, the affective dimensions of illness impair rational information processing, leading to decisions that may not align with patients' values or preferences when considered calmly .

Trust Deficits: Mistrust of insurer motives, pharmaceutical pricing, and even provider recommendations (fuelled by perceived conflicts of interest or negative experiences) hinders engagement and may lead to suboptimal decisions .

Transparency Gaps: Cost opacity remains the single most cited barrier to consumer shopping behaviour. Without price information available before purchase, consumers cannot compare alternatives on cost grounds.

2.7 Gaps in Existing Literature

The systematic review reveals several gaps warranting empirical investigation:

Geographic Concentration: The vast majority of research originates from North America and Western Europe, with limited empirical work in emerging economies like India, where healthcare systems, cultural norms, and consumer expectations differ substantially.

Methodological Monoculture: Quantitative surveys and discrete choice experiments predominate, with fewer qualitative studies capturing the lived experience of healthcare decision-making. Mixed-methods triangulation studies are particularly scarce.

Stakeholder Integration: Studies typically examine either consumer perspectives or provider perspectives, rarely integrating multiple stakeholder views within a single investigation.

Dynamic Processes: Most research captures decisions at a single time point, obscuring the iterative, evolving nature of healthcare choices as information accumulates and circumstances change.

Digital Transformation: The rapid evolution of digital health means that findings may become outdated quickly; ongoing research is needed to track changing patterns of technology adoption and use .

This study addresses these gaps through its focus on urban India, qualitative depth, multi-stakeholder design, and triangulation methodology.

3. Methodology

3.1 Research Philosophy and Design

This study adopted a pragmatist philosophical stance, prioritising research questions over paradigmatic commitments and employing methods best suited to addressing the objectives. A sequential mixed-methods design was employed, incorporating three complementary phases: systematic literature review, multiple case studies, and semi-structured interviews. The design was underpinned by triangulation principles—methodological, data source, and investigator triangulation—to enhance validity and comprehensiveness .

3.2 Phase 1: Systematic Literature Review

The systematic literature review followed the methodology detailed in Section 2.1, with findings used to develop the interview protocol and case study framework.

3.3 Phase 2: Multiple Case Studies

Case Selection: Four tertiary care hospitals in eastern India were purposively selected to maximise variation on key dimensions: ownership (two private corporate hospitals, one trust-run hospital, one public teaching hospital), size (ranging from 250 to 850 beds), and specialty focus (two general tertiary, one cardiac specialty, one oncology specialty). This diversity enabled examination of how organisational characteristics shape consumer decision contexts.

Data Collection: For each case, data were collected from three sources:

Documentary evidence: Websites, patient information materials, price lists (where available), quality reports, and social media presence.

Observational data: Researchers observed outpatient departments, admission counters, and patient information centres for 20 hours per hospital, documenting information availability, patient interactions, and decision environments.

Key informant discussions: Informal discussions with front-line staff (receptionists, billing clerks, nurses) to understand typical patient queries and decision processes.

Analysis: Case study data were analysed using within-case and cross-case synthesis techniques, identifying patterns in how hospitals presented information and facilitated (or impeded) consumer

decision-making.

3.4 Phase 3: Semi-Structured Interviews

Participant Sampling: Purposive sampling was employed to recruit 42 participants across four stakeholder groups:

Stakeholder Group	Sub-category	N
Patients (Total 24)	Completed treatment decisions (post-consultation)	12
	Currently making decisions (in process)	8
	Declined recommended treatment	4
Healthcare Providers (Total 8)	Specialist physicians	5
	General physicians	3
Hospital Administrators (Total 6)	Medical superintendents	2
	Patient relations managers	2
	Finance/billing managers	2
Health Insurance Representatives (Total 4)	Corporate agents	2
	Third-party administrators	2
TOTAL		42

Patient inclusion criteria: age ≥ 25 years, made a decision regarding tertiary care in the preceding 12 months, urban residence (Kolkata, Durgapur, or adjacent cities), able to provide informed consent. Patients under active treatment for life-threatening conditions were excluded to avoid undue burden.

Interview Protocol: A semi-structured interview guide was developed based on literature review findings and theoretical frameworks (HBM, TPB, EKB model). The guide explored:

Decision context and triggers (need recognition)

Information sources consulted and their perceived credibility

Evaluation processes (how alternatives were compared)

Role of family, friends, and social networks

Digital technology use (online research, reviews, telehealth)

Cost considerations and price transparency experiences

Barriers encountered and how they were navigated

Post-decision evaluation and reflections

Patient interviews averaged 55 minutes (range: 35–90 minutes). Provider/administrator/insurer interviews averaged 45 minutes (range: 30–70 minutes).

Procedure: Interviews were conducted between January and April 2024 at locations convenient for participants (hospital private spaces, participants' homes, or researcher's university office). All interviews were audio-recorded with permission, transcribed verbatim, and translated where necessary (some participants code-switched between English, Hindi, and Bengali). Field notes documented non-verbal cues and contextual observations.

Ethical Considerations: Ethical approval was obtained from the Institutional Ethics Committee of NSHM Knowledge Campus (IEC/NSHM/2023/47). All participants provided written informed consent. Confidentiality was assured through anonymisation (participants identified by codes: P[patient number], DR[provider number], AD[administrator number], IN[insurer number]). Participants received no financial compensation but were offered summary findings.

3.5 Data Analysis

Thematic Analysis: Interview transcripts were analysed using NVivo 14 following Braun and Clarke's six-phase thematic analysis framework:

Data familiarisation (repeated reading, initial note-taking)

Initial coding (systematic coding of entire dataset)

Theme generation (collating codes into potential themes)

Theme review (checking themes against coded extracts and dataset)

Theme definition (refining specifics of each theme)

Report writing (selecting vivid exemplars, relating to research questions)

Coding was initially inductive, allowing themes to emerge from participant accounts, followed by deductive coding aligned with theoretical frameworks and literature review findings.

Triangulation Procedures: Three forms of triangulation were employed :

Methodological triangulation: Comparing findings from literature review, case studies, and interviews to identify convergence and divergence.

Data source triangulation: Comparing perspectives across stakeholder groups (patients, providers, administrators, insurers) to develop multi-faceted understanding.

Investigator triangulation: Two researchers independently coded a subset of transcripts (20%) and compared coding decisions, discussing discrepancies until consensus reached. Inter-coder reliability (Cohen's κ = 0.82) indicated substantial agreement.

Rigor and Trustworthiness: Lincoln and Guba's criteria for qualitative rigor guided the study:

Credibility: Prolonged engagement (researchers spent extended time in hospital settings), member checking (summaries shared with 10 participants for verification), and peer debriefing.

Transferability: Thick description of context and participants enables readers to assess applicability to other settings.

Dependability: Audit trail documenting all methodological decisions.

Confirmability: Reflexive journaling to bracket researcher assumptions and biases.

3.6 Integration and Synthesis

Findings from all three phases were integrated through narrative synthesis, with interview themes illustrated by case study observations and situated within the broader literature. Tables and matrices facilitated cross-source comparison, with particular attention to areas of convergence (strengthening confidence) and divergence (generating insights about context-dependence).

4. Findings

The analysis yielded six overarching themes characterising healthcare consumer decision-making in urban India. Each theme is presented with supporting evidence from interviews, case study observations, and literature, followed by triangulation insights.

4.1 Theme 1: The Digitally Augmented Information Seeker

Description: Contemporary healthcare consumers actively research symptoms, conditions, providers, and treatment options online before and after clinical encounters. However, this digital engagement is best characterised as augmentation rather than replacement of traditional information sources.

Interview Evidence: All 24 patient participants reported using the internet for health-related searches, though with varying frequency and sophistication. The modal pattern involved Google searches for symptoms (often at night or when symptoms first appeared), followed by attempts to understand potential conditions before consulting a physician.

"When I first felt the chest pain, I didn't go to the doctor immediately. First, I Googled. Actually, I Googled at 2 AM because I couldn't sleep. It said it could be acidity, could be heart. That's when I got scared and decided I must see a cardiologist." (P7, 52-year-old male, post-angioplasty)

Younger participants (under 45) demonstrated more extensive online engagement, including reading hospital websites, physician profiles, and patient reviews. Several described "validating" physician recommendations through subsequent online research:

"The doctor said I needed surgery. I came home, I didn't sleep. I watched YouTube videos of the procedure, read articles, found forums where patients discussed their experiences. I even found a medical paper about success rates. Only then I felt ready to say yes." (P15, 38-year-old female, post-cholecystectomy)

However, participants also expressed difficulty assessing source credibility:

"There is too much information. One website says this, another says that. How do I know which doctor to believe? In the end, I just went with the one my cousin recommended." (P3, 45-year-old male, declined recommended knee replacement)

Provider Perspectives: Physicians acknowledged patients' online research, with ambivalent responses:

"Sometimes patients come with printouts from the internet. They want to discuss. If the information is good, it saves time. But often it's wrong, or it's for Western populations, not relevant here. Then I have to first un-teach before I can teach." (DR2, senior cardiologist)

Case Study Observations: All four hospitals maintained websites with physician directories and service listings. However, only the two corporate hospitals provided detailed physician profiles (training, experience, special interests). None provided price information online; all required in-person or telephone inquiries for cost estimates. Two hospitals had active social media presences (Facebook, Instagram) featuring patient testimonials and doctor interviews.

Triangulation: Literature strongly supports this theme: systematic reviews identify OHI-seeking as ubiquitous, with diagnostic/treatment information and emotional support as primary motivations. The credibility challenges voiced by participants mirror findings that consumers struggle to evaluate online health information quality. The provider ambivalence documented in interviews aligns with research showing that OHI influences patient-provider relationships and communication patterns .

4.2 Theme 2: Trust as the Foundational Currency

Description: Despite digital information proliferation, trust remains the fundamental determinant of healthcare decisions. Trust operates through multiple channels: provider reputation, personal recommendations, institutional affiliations, and relational continuity.

Interview Evidence: When asked what most influenced their final decision, 19 of 24 patients spontaneously mentioned trust-related factors. Provider recommendations carried exceptional weight:

"My family physician said, 'Go to Dr. X at Hospital Y. I have known him for twenty years. He is good.' That was enough. I didn't look anywhere else." (P11, 60-year-old female, completed cataract surgery)

Personal recommendations from family and friends functioned as trust shortcuts:

"My sister-in-law had the same surgery there last year. She said the doctor explains everything, the nurses are kind, the room was clean. I didn't need to know more." (P19, 48-year-old female, post-hysterectomy)

Institutional reputation (the hospital's "brand") also influenced trust, particularly among higher-income participants:

"In Kolkata, there is a hierarchy. [Hospital A] and [Hospital B] are top-tier. If you go there, you assume quality. It's like buying from Tata versus an unknown brand." (P22, 55-year-old male, completed cardiac procedure)

The Trust Paradox: Several participants described situations where trust was breached, with lasting consequences:

"The doctor was very famous. But he spent only five minutes with me. He didn't look at me when speaking. He dictated the prescription while typing. I felt like a number. I didn't go back." (P6, 42-year-old female, discontinued care)

Provider Perspectives: Physicians recognised their trust-signalling role:

"Patients don't know medicine. They can't judge my clinical skills directly. So they judge everything else: Do I listen? Do I explain? Is my clinic clean? Are my staff courteous? These become proxies for clinical trust." (DR5, general physician)

Administrator Perspectives: Hospital administrators described deliberate trust-building strategies:

"We invest heavily in patient experience training. Receptionists, nurses, billing staff—everyone understands they represent the hospital's trustworthiness. One rude interaction can undo years of reputation building." (AD3, patient relations manager)

Triangulation: This theme strongly converges with literature identifying trust as paramount. The finding that provider endorsement drives enrollment decisions directly aligns with research on remote health programs. The "trust proxies" identified by physicians (listening, explanation, courtesy) echo SERVQUAL dimensions of service quality. The breach narratives parallel research on trust violations and patient disengagement.

4.3 Theme 3: Price Sensitivity amidst Opacity

Description: Cost considerations significantly influence healthcare decisions, yet consumers face pervasive opacity that prevents meaningful price comparison. The result is a paradox: price matters greatly, but price information is rarely available when needed.

Interview Evidence: Twenty of 24 patients identified cost as a "major" or "significant" factor in their decisions. However, the timing of cost consideration varied:

"First I decided the doctor. Then I worried about the cost. But by then, I was committed. If it was too expensive, what could I do? I would borrow, sell something, manage somehow." (P12, 50-year-old male, completed bypass surgery)

This sequence—choose provider first, worry about cost later—was modal, reflecting the primacy of clinical trust over price considerations. However, a minority (particularly younger, employed participants) attempted price comparison:

"I called three hospitals. [Hospital C] quoted ₹80,000, [Hospital D] said ₹1.2 lakhs, [Hospital E] said ₹95,000. But when I asked what's included, nobody could give a clear breakdown. Package means different things to different hospitals." (P16, 41-year-old female, comparing maternity packages)

Negotiation Behaviours: Consistent with literature, negotiation emerged as a common value-seeking strategy:

"I told the billing manager, 'I have cash, I can pay today, but this is my budget.' They reduced the package by 15%. I didn't know you could negotiate in a hospital, but you can." (P9, 53-year-old male, completed orthopaedic surgery)

Insurer Perspectives: Insurance representatives highlighted how coverage complexity compounds opacity:

"Patients think 'I have insurance, so cost doesn't matter.' Then they discover: this doctor is out of network, this procedure has a sub-limit, this room rent exceeds coverage. The shock is tremendous. They feel cheated." (IN2, TPA representative)

Case Study Observations: None of the four hospitals displayed price information publicly. Price inquiries were handled at admission counters or billing departments, typically requiring personal interaction. Two hospitals offered "package rates" for common procedures (cataract, normal delivery, knee replacement), but package components varied, preventing apples-to-apples comparison.

Triangulation: Findings closely mirror US-based research documenting widespread price-seeking and negotiation behaviours (83% engaged in some negotiation). The sequencing pattern (choose first, cost later) aligns with literature suggesting clinical trust dominates initial decisions, with cost considerations entering during implementation rather than evaluation phases. The opacity finding is extensively documented as a systemic barrier .

4.4 Theme 4: Emotional Heuristics under Uncertainty

Description: Healthcare decisions are made under conditions of emotional arousal—anxiety, fear, hope, despair—that fundamentally alter information processing. Rather than systematic rational calculation,

consumers employ heuristics (mental shortcuts) that simplify choice under uncertainty.

Interview Evidence: Emotional narratives permeated patient accounts, particularly surrounding diagnosis and treatment decisions:

"When he said 'cancer,' I didn't hear anything after that. My wife was crying. My daughter was holding my hand. We just agreed to whatever he suggested. How could we think clearly at that moment?" (P20, 62-year-old male, post-oncology treatment)

Common Heuristics Identified:

Authority heuristic: Deferring to physician recommendations without question.

Social proof heuristic: Choosing providers or treatments that others have chosen.

Familiarity heuristic: Staying with known providers rather than exploring alternatives.

Scarcity heuristic: Valuing options perceived as limited ("he's the only specialist in the city").

"I chose this hospital because my neighbour went there. If it was good enough for him, it's good enough for me. I didn't compare anyone else." (P4, 58-year-old female, completed procedure)

The Paradox of Choice: Several participants described decision paralysis when faced with multiple options:

"I had five doctor names. All were qualified, all had good reviews. I couldn't decide. For three weeks I did nothing. Finally my son said, 'Pick one, any one.' I picked the one closest to home." (P13, 67-year-old male, delayed decision)

Provider Perspectives: Physicians recognised patients' emotional vulnerability:

"When I give a serious diagnosis, the patient isn't processing information. They're processing fear. I've learned to say the diagnosis, then pause, then say 'I will explain everything. You don't need to decide today. Take this information home, talk to family, then come back with questions.'" (DR7, oncologist)

Triangulation: These findings align with sentiment analysis research showing that positive sentiment correlates with option identification, while negative sentiment correlates with decision implementation. The heuristics identified (authority, social proof, familiarity) are well-documented in behavioural economics and decision science. The paralysis finding echoes the "choice overload" literature in consumer behaviour.

4.5 Theme 5: The Convenience Imperative

Description: Beyond clinical quality and trust, convenience factors increasingly drive provider selection and care engagement. Digital access (online scheduling, telehealth, electronic records) has become a

key convenience dimension.

Interview Evidence: Working-age participants particularly emphasised convenience:

"I can't take multiple leaves. My office is strict. If I can schedule online, get a Saturday appointment, consult by video for follow-ups—that hospital wins my business." (P17, 35-year-old male, managing chronic condition)

Telehealth Adoption: Eight participants had used telehealth consultations, primarily for follow-ups or minor concerns:

"After the surgery, I had questions. The video call saved me a four-hour round trip. The doctor saw my incision, said it's healing fine, done. I wish everything could be like this." (P8, 49-year-old male, post-operative follow-up)

Digital Expectations: Younger participants expressed frustration with hospitals lacking digital capabilities:

"Why can't I book online? Why do I have to call during busy hours? Why can't I see my reports on an app? My bank does this. My grocery does this. My hospital should do this." (P21, 29-year-old female, companion for elderly parent)

Provider Perspectives: Physicians acknowledged convenience pressures but expressed concerns:

"Patients want everything fast, convenient, digital. But good medicine takes time. You can't do complex consultations in ten-minute video calls. You can't build relationships through portals. Convenience is good, but not at the cost of quality." (DR3, senior physician)

Administrator Perspectives: Hospital administrators described investment in digital front doors:

"We've spent ₹2 crores on our digital platform. Online appointments, report access, payment, teleconsultation. Why? Because patients demand it. If we don't provide it, they'll go to the hospital down the road that does." (AD1, medical superintendent, corporate hospital)

Case Study Observations: The two corporate hospitals offered comprehensive digital access (online scheduling, patient portals, telehealth). The trust-run hospital had limited online appointment functionality. The public hospital had no digital patient-facing services; patients queued physically for appointments.

Triangulation: Findings strongly align with literature documenting "widespread comfort with hybrid care delivery" and patient preferences for digital access. The digital divide observed across hospital types mirrors research on technology adoption disparities in emerging economies. Provider concerns about quality and relationship echo ongoing debates in telemedicine literature.

4.6 Theme 6: Systemic Barriers to Informed Choice

Description: Multiple systemic barriers prevent consumers from making truly informed, value-congruent healthcare decisions, even when motivated and digitally equipped.

Barrier 1: Complexity and Jargon

"The doctor used so many English words. I didn't understand half of them. But I was embarrassed to ask. I just nodded and signed." (P1, 65-year-old female, post-procedure)

Barrier 2: Information Fragmentation

"To understand the cost, I had to go to three different counters. First registration, then doctor consultation, then billing. Each gave different information. Nobody had the complete picture." (P14, 44-year-old male, diagnostic workup)

Barrier 3: Time Pressure

"The doctor said, 'Decide by tomorrow, or the slot will go to someone else.' How can anyone decide major surgery in one day?" (P10, 56-year-old female, declined recommended surgery)

Barrier 4: Health Literacy Gaps

"They gave me a form with terms and conditions. I signed. What else could I do? I can't read English properly. I trusted them." (P2, 70-year-old male, completed treatment)

Barrier 5: Administrative Friction

"The insurance approval took one week. By then, my condition worsened. The system created the delay, but I suffered." (P18, 47-year-old male, delayed admission)

Provider Perspectives on Barriers:

"I wish I could spend an hour with each patient explaining everything. But I see 50 patients in a morning. The system doesn't allow meaningful counselling." (DR6, busy specialist)

Administrator Perspectives:

"We know the process is fragmented. We know patients get confused. We're working on integrating services, having patient navigators. But change is slow. Hospitals are complex organisations." (AD4, quality manager)

Triangulation: These barriers are extensively documented: complexity, information asymmetry, health literacy disparities, and administrative friction. The convergence between patient experiences, provider acknowledgements, and literature suggests these are systemic features of healthcare delivery, not merely individual failures.

4.7 Cross-Stakeholder Perspectives: Convergence and Divergence

Triangulation across stakeholder groups revealed both convergent and divergent perspectives:

Theme	Patient Perspective	Provider Perspective	Administrator/Insurer Perspective	Convergence
Digital Information Seeking	Essential but overwhelming	Ambivalent; useful but often inaccurate	Invest heavily in digital presence	High
Trust	Foundational; based on reputation, recommendations	Recognise trust-signalling role	Deliberately build trust through experience design	High
Cost Sensitivity	Significant but often considered late	Aware but may not discuss proactively	Price transparency is challenging to implement	Medium
Emotional State	Overwhelming; impairs decision capacity	Recognise and attempt to accommodate	Limited organisational response	Medium
Convenience	Increasingly expected	Concern about quality impact	Investing in digital access	Medium
Systemic Barriers	Frustrating and consequential	Acknowledge but feel powerless	Working on improvements; change is slow	High

High convergence on digital trends, trust, and systemic barriers strengthens confidence in these findings. Divergence on cost and emotional dimensions suggests areas where stakeholder communication could be improved.

5. Discussion

5.1 Synthesis of Findings

This study investigated healthcare consumer decision-making in urban India through systematic literature review, multiple case studies, and multi-stakeholder interviews, triangulated to enhance validity. The findings paint a portrait of consumers navigating between traditional trust-based heuristics and emerging digital empowerment, between rational information-seeking and emotional vulnerability, and between cost sensitivity and pervasive opacity.

The six themes identified—digitally augmented information seeking, trust as foundational currency, price sensitivity amidst opacity, emotional heuristics under uncertainty, the convenience imperative, and

systemic barriers—interact dynamically to shape decision outcomes. Figure 2 presents an integrated framework synthesising these relationships.

5.2 Theoretical Contributions

This study makes several contributions to the theoretical understanding of healthcare consumer behaviour.

Extension of Consumer Decision Models to Healthcare: The findings validate and extend the Engel-Kollat-Blackwell model as applied to healthcare. The identification of "seeking value" and "assessing value" as distinct sub-stages within the evaluation and post-purchase phases resonates strongly with participant accounts of negotiation, price comparison attempts, and post-treatment cost disputes. However, the model requires adaptation to accommodate healthcare's unique features: the profound influence of emotional state on information processing, the role of family as collective decision-makers, and the persistence of heuristics even when consumers are motivated to engage systematically.

Integration of HBM and TPB with Digital Contexts: The findings demonstrate how classic health behaviour theories operate in digitally saturated environments. Perceived barriers in the HBM now include digital literacy and online credibility assessment. Subjective norms in TPB now encompass online reviews and social media influences alongside traditional family and community norms. The "cues to action" construct increasingly includes digital triggers—targeted health content, app notifications, online symptom checkers.

Contextualising Trust in Healthcare Decisions: The study deepens understanding of trust as a multi-dimensional construct in healthcare. Participants distinguished between trust in clinical competence ("is the doctor skilled?"), trust in communication ("does the doctor explain and listen?"), trust in institutional systems ("will the hospital coordinate my care?"), and trust in financial integrity ("will I be charged fairly?"). These dimensions operate semi-independently; a provider may be trusted clinically but distrusted financially, affecting decision outcomes.

Emotional Heuristics Framework: Building on behavioural economics, this study identifies four heuristics —authority, social proof, familiarity, and scarcity—that systematically shape healthcare decisions under uncertainty. The finding that emotional arousal triggers heuristic reliance (rather than systematic processing) has implications for interventions designed to support better decisions.

5.3 Practical Implications

The findings carry actionable implications for healthcare stakeholders.

5.3.1 For Healthcare Providers

Enhance the Digital Front Door: Given consumer expectations for digital access, providers must invest in user-friendly online scheduling, telehealth capabilities, patient portals, and digital communication.

However, technology should augment rather than replace human connection; the trust-building functions of interpersonal interaction cannot be fully digitised.

Communicate Trustworthiness Explicitly: Trust is signalled through multiple channels: clear communication, empathetic listening, courteous staff, clean facilities, transparent processes. Providers should audit the patient journey for trust signals and address trust deficits systematically.

Improve Cost Transparency: While full price transparency faces genuine challenges (variation in patient condition, insurance complexity, bundled services), providers can take incremental steps: offering package rates for common procedures, providing written cost estimates before service, training billing staff to explain costs clearly, and making price information available online where possible .

Simplify Without Paternalising: Patients need information they can understand without being infantilised. Plain language explanations, visual aids, decision aids, and teach-back techniques can bridge the gap between medical complexity and patient comprehension. Trained patient navigators or counsellors can support decision-making for complex, high-stakes choices.

Support Deliberate Decisions: Given the finding that time pressure leads to decision paralysis or suboptimal choices, providers should, where clinically appropriate, build in "cooling off" periods that allow patients to process information, consult family, and return with questions before finalising major decisions.

5.3.2 For Health Insurers

Demystify Coverage: Insurance complexity emerged as a major barrier. Insurers should simplify plan documents, provide real-time benefit checks (showing exact out-of-pocket costs before service), train customer service representatives to explain coverage clearly, and design digital tools that help consumers understand their benefits.

Align Incentives with Consumer Value: Rather than passive claims payment, insurers can actively support consumer decision-making through curated provider networks, quality and cost transparency tools, telehealth benefits, and value-added services that address whole-person needs.

Partner with Providers on Transparency: Insurers and providers should collaborate on standardising cost estimates, clarifying what is and isn't covered, and streamlining prior authorisation processes that currently create friction and delay.

5.3.3 For Policymakers

Mandate and Enforce Transparency: Building on the finding that opacity systematically impedes informed choice, policymakers should strengthen price transparency regulations, requiring hospitals to publish standardised package rates for common procedures and provide patients with personalised cost estimates before service.

Invest in Health Literacy: Given persistent health literacy disparities, public health investments should include community-based health literacy programmes, plain language requirements for health communications, and support for patient decision aids.

Promote Digital Inclusion: As healthcare increasingly digitises, policymakers must ensure equitable access through broadband infrastructure, digital literacy programmes, and regulations that prevent digital exclusion of vulnerable populations.

Support Interoperability: Fragmented information across providers, insurers, and patients impedes informed decision-making. Policies promoting health data interoperability can empower consumers with comprehensive information.

5.3.4 For Technology Developers

Design for Decision Support, Not Just Data Collection: Many health apps collect data but fail to support decision-making. Developers should design tools that synthesise information, clarify options, and guide users through decisions—not just track steps or store records.

Prioritise Credibility Cues: Given consumer struggles with online information credibility, platforms should incorporate clear source attribution, quality ratings, and cues that help users assess trustworthiness.

Ensure Accessibility: Health technologies must be usable by populations with varying digital literacy, language preferences, and visual or cognitive abilities. Universal design principles should guide development.

Protect Privacy and Build Trust: Data breaches and privacy violations disproportionately harm vulnerable populations and erode trust in digital health. Developers must prioritise security, transparency, and user control over personal data.

5.4 The Consumer-Centric Healthcare Organisation

Synthesising the implications, Table 2 outlines the characteristics of a consumer-centric healthcare organisation across multiple domains.

Table 2
Dimensions of Consumer-Centric Healthcare Organisations

Dimension	Traditional Approach	Consumer-Centric Approach
Information Provision	Reactive; provided when asked	Proactive; anticipate information needs
Digital Access	Optional; basic website	Comprehensive; online scheduling, telehealth, portal, app
Cost Communication	After service; unclear breakdown	Before service; written estimate, package options
Decision Support	Minimal; "doctor knows best"	Active; decision aids, counsellors, cooling-off periods
Patient Experience	Clinical focus only	Holistic; emotional support, convenience, dignity
Feedback Integration	Sporadic; complaints only	Systematic; regular surveys, patient councils
Trust Building	Assumed based on credentials	Deliberately cultivated through actions
Equity Focus	One-size-fits-all	Tailored to health literacy, language, culture

6. Limitations and Future Research

6.1 Limitations

This study has several limitations that should be considered when interpreting findings.

Geographic Scope: The empirical research was conducted in urban eastern India (Kolkata, Durgapur, and adjacent cities). While this addresses a gap in literature focused on Western contexts, findings may not generalise to rural India, other Indian regions with different cultural and economic profiles, or other countries. The urban, relatively educated sample may overrepresent digitally engaged consumers.

Sample Characteristics: Despite purposive sampling for diversity, the patient sample was predominantly middle-class, with 22 of 24 participants having completed secondary education. Lower-income, less-educated consumers—who face greater health literacy challenges and different decision constraints—are underrepresented. The sample also underrepresents non-English and non-Bengali speakers.

Cross-Sectional Design: Interviews captured retrospective accounts of decisions, subject to recall bias and post-decision rationalisation. The cross-sectional design cannot track how decision processes evolve over time or how preferences change with experience.

Focus on Elective/Planned Care: By design, the study excluded emergency decisions. Findings apply only to situations where consumers have time and capacity for deliberation, not to time-pressured, high-acuity contexts.

Rapid Digital Evolution: The digital health landscape evolves continuously; findings regarding specific technologies (telehealth platforms, apps, review sites) may require updating as new tools emerge and adoption patterns shift .

Single-Country Context: India's unique healthcare ecosystem—mixed public-private provision, high out-of-pocket expenditure, limited health insurance penetration, cultural emphasis on family decision-making—shapes findings in ways that may limit transferability to other settings.

6.2 Future Research Directions

The findings suggest multiple avenues for future investigation.

Longitudinal Decision Pathways: Research tracking consumers over time—from initial symptom recognition through information search, evaluation, decision, and post-decision experience—would illuminate how preferences evolve and how decisions are revised in light of new information. Such studies could identify critical junctures where interventions might be most effective.

Health Literacy Interventions: Given persistent health literacy disparities, intervention studies testing the effectiveness of decision aids, plain language communications, and patient navigation programmes would provide evidence for scalable solutions. Research should examine effects on decision quality, satisfaction, and outcomes, not just comprehension.

Vulnerable Population Focus: Deliberate over-sampling of low-income, low-literacy, and non-digital consumers would illuminate how these populations navigate (or are excluded from) consumer-directed healthcare. Such research could inform equity-focused policy and practice.

Treatment-Specific Decision Processes: Consumer behaviour likely varies by clinical condition. High-stakes, high-variation areas—oncology, cardiology, elective surgery, chronic disease management—warrant focused investigation to identify condition-specific decision dynamics.

AI and Personalisation: As artificial intelligence increasingly personalises health information and recommendations, research should examine how AI-driven tools affect decision quality, trust, and equity. Do personalisation algorithms reduce or amplify existing disparities? How do consumers perceive AI-generated versus human-generated recommendations?

Cross-Cultural Comparative Studies: Systematic cross-national research comparing healthcare consumer behaviour across cultures—varying in individualism/collectivism, trust in institutions, health system structures, and digital adoption—would illuminate which findings are universal and which are context-dependent.

Stakeholder Dynamics: This study examined stakeholders separately; future research could investigate stakeholder interactions directly—patient-provider consultations, family decision conferences, insurer-consumer communications—to understand how decisions are negotiated in real time.

Implementation Research: Given the practice implications identified, implementation science research examining how healthcare organisations successfully (or unsuccessfully) adopt consumer-centric practices would bridge the gap between evidence and action.

7. Conclusion

This study investigated healthcare consumer decision-making in urban India through a triangulated mixed-methods design encompassing systematic literature review, multiple case studies, and multi-stakeholder interviews. The findings reveal a complex landscape in which consumers navigate between traditional trust-based heuristics and emerging digital empowerment, between rational information-seeking and emotional vulnerability, and between cost sensitivity and pervasive opacity.

The six themes identified—digitally augmented information seeking, trust as foundational currency, price sensitivity amidst opacity, emotional heuristics under uncertainty, the convenience imperative, and systemic barriers—interact dynamically to shape decision outcomes. While consumers are increasingly empowered through digital access, meaningful decision-making remains constrained by complexity, trust deficits, and transparency gaps. The convergence of findings across stakeholder groups and with international literature strengthens confidence in these conclusions.

The theoretical contributions include extending consumer decision models to healthcare contexts, integrating classic health behaviour theories with digital realities, deepening understanding of trust as a multi-dimensional construct, and identifying emotional heuristics that guide decisions under uncertainty.

For practice, the implications are clear: healthcare stakeholders must embrace genuinely consumer-centric strategies that simplify choice, enhance cost transparency, leverage technology judiciously, and build authentic trust. This requires coordinated action across providers, insurers, policymakers, and technology developers—each playing complementary roles in supporting informed, value-congruent decisions.

The transition from paternalistic medicine to consumer-directed healthcare is neither complete nor uncontested. It carries risks alongside opportunities—risks of exacerbating inequities, of burdening vulnerable individuals with choices they cannot make, of commodifying care in ways that erode healing relationships. But the transition is underway regardless. Understanding and constructively shaping it is now imperative—not merely for improving individual health outcomes, but for building healthcare systems that are simultaneously more responsive, more equitable, and more sustainable.

As one participant reflected:

"I want to be a partner in my health, not just a patient. I want information I can trust, choices I can understand, costs I can plan for. Is that too much to ask?" (P22)

The findings of this study suggest that these aspirations are reasonable, achievable, and urgently deserving of response.

References

1. Crespo-Gonzalez C, Benrimoj SI, Frommer M, Dineen-Griffin S (2024) Navigating online health information: Insights into consumer influence and decision-making strategies—An overview of reviews. Digit Health 10:20552076241286815
2. Spacht WA, Lee SG, Varugheese M, Subramaniam S, McPartlin M, Tucci MR, Scirica BM (2025) Evaluating patient and provider experiences of enrolling in a remote cardiovascular health program: A qualitative interview study. BMC Prim Care 26:304
3. GMDS Working Group Consumer Health Informatics (2015) Metamodel for consumer health informatics. German Society for Medical Informatics, Biometry and Epidemiology
4. De G, Sondhi N, Bhattacharjee A, Joshi H (2024) Preventive healthcare behavior: A hybrid systematic literature review (1998–2023). Int J Consumer Stud, 48(1), e12989
5. Crespo-Gonzalez C, Benrimoj SI, Frommer M, Dineen-Griffin S (2024) Navigating online health information: Insights into consumer influence and decision-making strategies—An overview of reviews. Digit HEALTH, 10
6. Sinaiko AD, Rosenthal MB (2020) Examining consumer shopping behaviors and sentiment in healthcare purchasing. J Participatory Med, 12(2), e13924
7. Whitty JA, Rundle-Thiele SR, Scuffham PA (2012) Insights from triangulation of two purchase choice elicitation methods to predict social decision making in healthcare. Appl Health Econ Health Policy 10(2):113–126
8. Singh A, Padhy RK, Sahu A (2025) Bridging the healthcare gap: Understanding continued telemedicine-usage behaviour in emerging economies. *Global Knowledge, Memory and Communication*
9. Khatib R, Guzman I, Glowacki N, Lauffenburger JC, Anderson M, Mora N (2025) Adopting self-measured blood pressure monitoring in underserved primary care settings: A qualitative study using customer discovery and value proposition principles. BMC Prim Care 26(1):304
10. Whitty JA, Rundle-Thiele SR, Scuffham PA (2012) Insights from triangulation of two purchase choice elicitation methods to predict social decision making in healthcare. Appl Health Econ Health Policy 10(2):113–126
11. Berry LL, Bendapudi N (2007) Health care: A fertile field for service research. J Service Res 10(2):111–122
12. Hibbard JH, Greene J (2013) What the evidence shows about patient activation: Better health outcomes and care experiences; fewer data on costs. Health Aff 32(2):207–214
13. Rosenstock IM (1974) Historical origins of the health belief model. Health Educ Monogr 2(4):328–335
14. Ajzen I (1991) The theory of planned behavior. Organ Behav Hum Decis Process 50(2):179–211
15. Parasuraman A, Zeithaml VA, Berry LL (1988) SERVQUAL: A multiple-item scale for measuring consumer perceptions of service quality. J Retail 64(1):12–40

16. Deloitte (2023) 2023 Global Health Care Outlook. Deloitte Insights
17. Kaiser Family Foundation (2023) Employer Health Benefits Survey. KFF
18. Accenture (2022) *Digital Health Technology Vision*. Accenture
19. Agency for Healthcare Research and Quality (2023) Health Literacy Universal Precautions Toolkit, 3rd edn. AHRQ
20. Centers for Medicare & Medicaid Services (2023) *Hospital Compare*

Figures



Figure 1

Research Design Framework



Figure 2

Integrated Framework of Healthcare Consumer Decision-Making