

Supporting Information

Photo-activated Charge Transport and Unified Mechanism of Self-Trapped Excitons in Cs₂NaInCl₆ Double Perovskite

Fahad K. Alshammari ¹, Abdullah A. Alatawi ², Achref Jebnoui ³, Amjad Salamah Aljaloud ¹, Turki Alkathiri ^{4,*}, Ahmed F Almutairi ⁵, Bandar F. Alsaleh ¹, Mohamed Bouzidi ¹, Mohamed Ben Bechir ^{6,7}

¹ Department of Physics, College of Science, University of Ha'il, Ha'il P.O. Box 2440, Saudi Arabia

² King Abdulaziz City for Science and Technology, P.O. Box 6086, 11442 Riyadh, Saudi Arabia

³ Medical Surgical Nursing Department, College of Nursing, University of Ha'il, Ha'il, Saudi Arabia

⁴ Department of Electrical Engineering, College of Engineering, Al-Baha University, P.O. Box 65799, Albaha, Saudi Arabia

⁵ Department of Mechanical Engineering, College of Engineering at Yanbu, Taibah University, Yanbu Al-Bahr 41911, Saudi Arabia

⁶ Laboratory of Spectroscopic and Optical Characterization of Materials (LaSCOM), Faculty of Sciences, University of Sfax, BP1171 – 3000 Sfax, Tunisia

⁷ Department of Physics, Faculty of Sciences of Gafsa, University of Gafsa, 2112, Tunisia

* Corresponding author. E-mail address: talkathiri@bu.edu.sa, mohamedbenbechir@hotmail.com

Supplementary Figures




Fig. S1. X-ray diffraction refinement profile for Cs₂NaInCl₆, showing good agreement between experimental and simulated data.











Fig. S2. Equivalent circuit model fitted to the impedance spectra of $\text{Cs}_2\text{NaInCl}_6$, showing the contribution of space-charge polarization and bulk relaxation processes.