

2025 Internal Medicine Critical Care Triage Series

Please take a few minutes to reflect on the training and provide responses to the following questionnaire. This questionnaire will take you less than 5 minutes to complete. The information gathered will be helpful in planning future sessions.

Thank you!

Which session did you complete today?
Select ALL that apply.

- ☐ Anaphylaxis
- ☐ Complete Heart Block
- ☐ Cardiogenic Shock
- ☐ Ventricular Tachycardia
- ☐ Cardiac Tamponade
- ☐ Pulmonary Embolism
- ☐ Flash Pulmonary Edema
- ☐ Upper Gastrointestinal Bleed
- ☐ Lower Gastrointestinal Bleed
- ☐ Hemoptysis
- ☐ Small Bowel Obstruction
- ☐ Hypovolemic Shock
- ☐ Opioid Overdose
- ☐ Delirium
- ☐ Seizures
- ☐ Hypoglycemia

Please indicate the extent to which you agree with the following statements regarding the content of the training.

	Strongly Disagree	Disagree	Neutral	Agree	Strongly Agree
The content was relevant to my work.	☐	☐	☐	☐	☐
This training provided me with new information (or clarified existing information).	☐	☐	☐	☐	☐
I intend to use what I learned from this training.	☐	☐	☐	☐	☐

If you agree or strongly agree with this statement, please provide us with details as to which component(s) of the training you plan on utilizing.

Please indicate the extent to which you agree with the following statements regarding the facilitator(s) of the training.

	Strongly Disagree	Disagree	Neutral	Agree	Strongly Agree
The facilitator(s) was knowledgeable about the subject.	☐	☐	☐	☐	☐
The facilitator(s) was effective in helping me learn new information (or clarify existing information).	☐	☐	☐	☐	☐
The facilitator(s) was responsive to participants' needs and questions.	☐	☐	☐	☐	☐

Please indicate the extent to which you agree with the following statement regarding the learning environment.

	Strongly Disagree	Disagree	Neutral	Agree	Strongly Agree
The learning environment was conducive to learning.	☐	☐	☐	☐	☐

Please provide us with suggestions for improving the content, facilitation, delivery, environment and/or utility of the training.

Please indicate the extent to which you agree with the following statement regarding the use of simulation.

	Strongly Disagree	Disagree	Neutral	Agree	Strongly Agree
SIM based experience is more effective for my learning compared to traditional classroom-based lectures.	☐	☐	☐	☐	☐
This simulation helped me identify gaps in my knowledge.	☐	☐	☐	☐	☐
I would like more frequent case-based simulations instead of standard classroom-based lectures.	☐	☐	☐	☐	☐

Prior to today's Anaphylaxis activity, I was confident in my ability to:

	Strongly Disagree	Disagree	Neutral	Agree	Strongly Agree
Recognize signs and symptoms of the patient in anaphylaxis.	☐	☐	☐	☐	☐
Initiate appropriate management for treatment of anaphylaxis.	☐	☐	☐	☐	☐
Recognize and manage a difficult airway.	☐	☐	☐	☐	☐

As a result of today's Anaphylaxis activity, I am more confident in my ability to:

	Strongly Disagree	Disagree	Neutral	Agree	Strongly Agree
Recognize signs and symptoms of the patient in anaphylaxis.	☐	☐	☐	☐	☐
Initiate appropriate management for treatment of anaphylaxis.	☐	☐	☐	☐	☐
Recognize and manage a difficult airway.	☐	☐	☐	☐	☐

Prior to today's Complete Heart Block activity, I was confident in my ability to:

	Strongly Disagree	Disagree	Neutral	Agree	Strongly Agree
Recognize ECG findings of complete heart block.	☐	☐	☐	☐	☐
Initiate stabilization with transcutaneous pacing.	☐	☐	☐	☐	☐
Recognize need to escalate to transvenous pacing.	☐	☐	☐	☐	☐

As a result of today's Complete Heart Block activity, I am more confident in my ability to:

	Strongly Disagree	Disagree	Neutral	Agree	Strongly Agree
Recognize ECG findings of complete heart block.	☐	☐	☐	☐	☐
Initiate stabilization with transcutaneous pacing.	☐	☐	☐	☐	☐
Recognize need to escalate to transvenous pacing.	☐	☐	☐	☐	☐

Prior to today's Cardiogenic Shock activity, I was confident in my ability to:

	Strongly Disagree	Disagree	Neutral	Agree	Strongly Agree
Recognize signs and symptoms of cardiogenic shock.	☐	☐	☐	☐	☐
Order appropriate diagnostic evaluation to further differentiate shock.	☐	☐	☐	☐	☐
Initiate appropriate management for cardiogenic shock.	☐	☐	☐	☐	☐

As a result of today's Cardiogenic Shock activity, I am more confident in my ability to:

	Strongly Disagree	Disagree	Neutral	Agree	Strongly Agree
Recognize signs and symptoms of cardiogenic shock.	☐	☐	☐	☐	☐
Order appropriate diagnostic evaluation to further differentiate shock.	☐	☐	☐	☐	☐
Initiate appropriate management for cardiogenic shock.	☐	☐	☐	☐	☐

Prior to today's Ventricular Tachycardia activity, I was confident in my ability to:

	Strongly Disagree	Disagree	Neutral	Agree	Strongly Agree
Recognize ECG findings of ventricular tachycardia.	☐	☐	☐	☐	☐
Initiate appropriate management for stable ventricular tachycardia.	☐	☐	☐	☐	☐
Initiate appropriate management for unstable ventricular tachycardia.	☐	☐	☐	☐	☐

As a result of today's Ventricular Tachycardia activity, I am more confident in my ability to:					
	Strongly Disagree	Disagree	Neutral	Agree	Strongly Agree
Recognize ECG findings of ventricular tachycardia.	☐	☐	☐	☐	☐
Initiate appropriate management for stable ventricular tachycardia.	☐	☐	☐	☐	☐
Initiate appropriate management for unstable ventricular tachycardia.	☐	☐	☐	☐	☐

Prior to today's Cardiac Tamponade activity, I was confident in my ability to:

	Strongly Disagree	Disagree	Neutral	Agree	Strongly Agree
Recognize clinical and hemodynamic signs of cardiac tamponade.	☐	☐	☐	☐	☐
Initiate interventions for stabilization of cardiac tamponade.	☐	☐	☐	☐	☐
Identify need for definitive management with pericardiocentesis.	☐	☐	☐	☐	☐

As a result of today's Cardiac Tamponade activity, I am more confident in my ability to:

	Strongly Disagree	Disagree	Neutral	Agree	Strongly Agree
Recognize clinical and hemodynamic signs of cardiac tamponade.	☐	☐	☐	☐	☐
Initiate interventions for stabilization of cardiac tamponade.	☐	☐	☐	☐	☐
Identify need for definitive management with pericardiocentesis.	☐	☐	☐	☐	☐

Prior to today's Pulmonary Embolism activity, I was confident in my ability to:

	Strongly Disagree	Disagree	Neutral	Agree	Strongly Agree
Develop a differential diagnosis for respiratory failure based on history and physical examination.	☐	☐	☐	☐	☐
Order appropriate diagnostic evaluation of respiratory failure.	☐	☐	☐	☐	☐
Initiate appropriate management for a pulmonary embolism.	☐	☐	☐	☐	☐

As a result of today's Pulmonary Embolism activity, I am more confident in my ability to:

	Strongly Disagree	Disagree	Neutral	Agree	Strongly Agree
Develop a differential diagnosis for respiratory failure based on history and physical examination.	☐	☐	☐	☐	☐
Order appropriate diagnostic evaluation of respiratory failure.	☐	☐	☐	☐	☐
Initiate appropriate management for a pulmonary embolism.	☐	☐	☐	☐	☐

Prior to today's Flash Pulmonary Edema activity, I was confident in my ability to:

	Strongly Disagree	Disagree	Neutral	Agree	Strongly Agree
Recognize clinical signs and symptoms of hypertensive emergency.	☐	☐	☐	☐	☐
Initiate appropriate management for hypertensive emergencies.	☐	☐	☐	☐	☐
Identify respiratory failure as a complication of hypertensive emergencies.	☐	☐	☐	☐	☐

As a result of today's Flash Pulmonary Edema activity, I am more confident in my ability to:

Strongly Disagree Disagree Neutral Agree Strongly Agree

Recognize clinical signs and symptoms of hypertensive emergency.

☐☐☐☐☐

Initiate appropriate management for hypertensive emergencies.

☐☐☐☐☐

Identify respiratory failure as a complication of hypertensive emergencies.

☐☐☐☐☐

Prior to today's Upper GI Bleed activity, I was confident in my ability to:

	Strongly Disagree	Disagree	Neutral	Agree	Strongly Agree
Recognize clinical signs and symptoms of an upper gastrointestinal bleed.	☐	☐	☐	☐	☐
Initiate appropriate management based on clinical stability in the setting of a gastrointestinal bleed.	☐	☐	☐	☐	☐
Recognize need for more urgent diagnostic work-up or consultation.	☐	☐	☐	☐	☐

As a result of today's Upper GI Bleed activity, I am more confident in my ability to:

	Strongly Disagree	Disagree	Neutral	Agree	Strongly Agree
Recognize clinical signs and symptoms of an upper gastrointestinal bleed.	☐	☐	☐	☐	☐
Initiate appropriate management based on clinical stability in the setting of a gastrointestinal bleed.	☐	☐	☐	☐	☐
Recognize need for more urgent diagnostic work-up or consultation.	☐	☐	☐	☐	☐

Prior to today's Lower GI Bleed activity, I was confident in my ability to:

	Strongly Disagree	Disagree	Neutral	Agree	Strongly Agree
Recognize clinical signs and symptoms of a lower gastrointestinal bleed.	☐	☐	☐	☐	☐
Initiate appropriate management based on clinical stability in the setting of a gastrointestinal bleed.	☐	☐	☐	☐	☐
Recognize need for more urgent diagnostic work-up or consultation.	☐	☐	☐	☐	☐

As a result of today's Lower GI Bleed activity, I am more confident in my ability to:

	Strongly Disagree	Disagree	Neutral	Agree	Strongly Agree
Recognize clinical signs and symptoms of a lower gastrointestinal bleed.	☐	☐	☐	☐	☐
Initiate appropriate management based on clinical stability in the setting of a gastrointestinal bleed.	☐	☐	☐	☐	☐
Recognize need for more urgent diagnostic work-up or consultation.	☐	☐	☐	☐	☐

Prior to today's Hemoptysis activity, I was confident in my ability to:

	Strongly Disagree	Disagree	Neutral	Agree	Strongly Agree
Distinguish between massive and nonmassive hemoptysis.	☐	☐	☐	☐	☐
Initiate appropriate initial management of hemoptysis.	☐	☐	☐	☐	☐
Recognize impending respiratory compromise in the setting of massive hemoptysis.	☐	☐	☐	☐	☐

As a result of today's Hemoptysis activity, I am more confident in my ability to:

	Strongly Disagree	Disagree	Neutral	Agree	Strongly Agree
Distinguish between massive and nonmassive hemoptysis.	☐	☐	☐	☐	☐
Initiate appropriate initial management of hemoptysis.	☐	☐	☐	☐	☐
Recognize impending respiratory compromise in the setting of massive hemoptysis.	☐	☐	☐	☐	☐

Prior to today's Small Bowel Obstruction activity, I was confident in my ability to:

	Strongly Disagree	Disagree	Neutral	Agree	Strongly Agree
Recognize classic presentation and risk factors for small bowel obstruction.	☐	☐	☐	☐	☐
Initiate appropriate management for small bowel obstructions.	☐	☐	☐	☐	☐
Recognize complication of small bowel obstructions and indications for urgent surgical consultation.	☐	☐	☐	☐	☐

As a result of today's Small Bowel Obstruction activity, I am more confident in my ability to:

	Strongly Disagree	Disagree	Neutral	Agree	Strongly Agree
Recognize classic presentation and risk factors for small bowel obstruction.	☐	☐	☐	☐	☐
Initiate appropriate management for small bowel obstructions.	☐	☐	☐	☐	☐
Recognize complication of small bowel obstructions and indications for urgent surgical consultation.	☐	☐	☐	☐	☐

Prior to today's Hypovolemic Shock activity, I was confident in my ability to:

	Strongly Disagree	Disagree	Neutral	Agree	Strongly Agree
Identify early signs of hypovolemic shock from significant gastrointestinal loss.	☐	☐	☐	☐	☐
Initiate appropriate management and work-up for diarrhea.	☐	☐	☐	☐	☐
Recognize severe <i>Clostridioides difficile</i> infection and its complications.	☐	☐	☐	☐	☐

As a result of today's Hypovolemic Shock activity, I am more confident in my ability to:

	Strongly Disagree	Disagree	Neutral	Agree	Strongly Agree
Identify early signs of hypovolemic shock from significant gastrointestinal loss.	☐	☐	☐	☐	☐
Initiate appropriate management and work-up for diarrhea.	☐	☐	☐	☐	☐
Recognize severe <i>Clostridioides difficile</i> infection and its complications.	☐	☐	☐	☐	☐

Prior to today's Opioid Overdose activity, I was confident in my ability to:

	Strongly Disagree	Disagree	Neutral	Agree	Strongly Agree
Recognize clinical signs and symptoms of opioid toxidrome.	☐	☐	☐	☐	☐
Order appropriate diagnostic evaluation for altered mentation.	☐	☐	☐	☐	☐
Appropriately treat opioid overdose.	☐	☐	☐	☐	☐

As a result of today's Opioid Overdose activity, I am more confident in my ability to:

	Strongly Disagree	Disagree	Neutral	Agree	Strongly Agree
Recognize clinical signs and symptoms of opioid toxidrome.	☐	☐	☐	☐	☐
Order appropriate diagnostic evaluation for altered mentation.	☐	☐	☐	☐	☐
Appropriately treat opioid overdose.	☐	☐	☐	☐	☐

Prior to today's Delirium activity, I was confident in my ability to:

	Strongly Disagree	Disagree	Neutral	Agree	Strongly Agree
Recognize delirium and develop an appropriate differential diagnosis.	☐	☐	☐	☐	☐
Order appropriate diagnostic evaluation of delirium.	☐	☐	☐	☐	☐
Initiate treatment with non-pharmacological measures and recognize when pharmacological management is appropriate.	☐	☐	☐	☐	☐

As a result of today's Delirium activity, I am more confident in my ability to:

	Strongly Disagree	Disagree	Neutral	Agree	Strongly Agree
Recognize delirium and develop an appropriate differential diagnosis.	☐	☐	☐	☐	☐
Order appropriate diagnostic evaluation of delirium.	☐	☐	☐	☐	☐
Initiate treatment with non-pharmacological measures and recognize when pharmacological management is appropriate.	☐	☐	☐	☐	☐

Prior to today's Seizure activity, I was confident in my ability to:

	Strongly Disagree	Disagree	Neutral	Agree	Strongly Agree
Recognize clinical signs and symptoms of seizure.	☐	☐	☐	☐	☐
Initiate appropriate management of focal seizure.	☐	☐	☐	☐	☐
Initiate appropriate management of status epilepticus.	☐	☐	☐	☐	☐

As a result of today's Seizure activity, I am more confident in my ability to:

	Strongly Disagree	Disagree	Neutral	Agree	Strongly Agree
Recognize clinical signs and symptoms of seizure.	☐	☐	☐	☐	☐
Initiate appropriate management of focal seizure.	☐	☐	☐	☐	☐
Initiate appropriate management of status epilepticus.	☐	☐	☐	☐	☐

Prior to today's Hypoglycemia activity, I was confident in my ability to:

	Strongly Disagree	Disagree	Neutral	Agree	Strongly Agree
--	----------------------	----------	---------	-------	----------------

Recognize clinical signs and symptoms of hypoglycemia.

☐☐☐☐☐

Initiate appropriate initial treatment for hypoglycemia.

☐☐☐☐☐

Identify underlying etiology of hypoglycemia and tailor ongoing management appropriately.

☐☐☐☐☐

As a result of today's Hypoglycemia activity, I am more confident in my ability to:

	Strongly Disagree	Disagree	Neutral	Agree	Strongly Agree
Recognize clinical signs and symptoms of hypoglycemia.	☐	☐	☐	☐	☐
Order appropriate diagnostic evaluation of altered mental status.	☐	☐	☐	☐	☐
Identify underlying etiology of hypoglycemia and tailor ongoing management appropriately.	☐	☐	☐	☐	☐