Research Square

Preprints are preliminary reports that have not undergone peer review.
They should not be considered conclusive, used to inform clinical practice,
or referenced by the media as validated information.

Knowledge, Access, and Utilization of Long-Lasting Insecticidal Treated Nets among Pregnant Women in Rural Ghana: A Cross-Sectional Study

Richmond Osei Yaw
Anglican University College of Technology (ANGUTECH)
Ishmael Awini Aburi
Anglican University College of Technology (ANGUTECH)
Abigail Boatemaa
University of Bradford
Gabriel Osei Forkuo
gabriel.forkuo@unitbv.ro

Transilvania University of Brasov

Research Article

Keywords: antenatal care, behaviour change, malaria in pregnancy, sub-Saharan Africa, socio-demographic, vector-control intervention

Posted Date: May 9th, 2026

DOI: https://doi.org/10.21203/rs.3.rs-9344718/v1

License: This work is licensed under a Creative Commons Attribution 4.0 International License.
Read Full License

Additional Declarations: No competing interests reported.

Abstract

Malaria in pregnancy remains a leading cause of preventable maternal and neonatal mortality in sub-Saharan Africa. While distribution programs have achieved near-universal net ownership, a critical gap persists between access and consistent utilization. This study characterized the knowledge, access, and practice (KAP) profile regarding long-lasting insecticidal treated nets (LLITNs) among pregnant women in a rural Ghanaian community to identify the determinants of the ownership–utilization gap. A descriptive cross-sectional KAP design was employed among 200 purposively selected pregnant women. Data were collected through structured interviews and direct physical observation of sleeping environments. Analysis was performed using descriptive statistics and a suite of eight advanced multivariate visualizations, including heatmaps, violin plots, and radar charts. The study revealed a bimodal KAP profile. Awareness of malaria-prevention benefits (100%) and antenatal care access (98.5%) were near-universal. However, knowledge depth was critically low; only 27.1% correctly identified the three-year insecticide efficacy period. Direct observation confirmed that only 43.5% of nets were correctly hung and in use. Farmers exhibited the highest discordance between ownership and utilization (34% non-use), and respondents with poor knowledge were disproportionately represented among non-users (44%). A "Universal Access Paradox" exists where supply-side success coexists with demand-side failure. Misconceptions regarding insecticide duration drive "perceived obsolescence" and premature net abandonment. Programs must transition to a "Distribution Plus" model that integrates structured, occupationally-sensitive counseling and observational monitoring to translate near-universal access into effective biological protection.

1. Introduction

Malaria in pregnancy (MiP) constitutes one of the most consequential preventable causes of maternal and perinatal morbidity and mortality in sub-Saharan Africa, imposing a burden that extends from the individual woman and her unborn child to the household, community, and national economy. The pathophysiological basis of this vulnerability is well established: pregnancy induces a state of partial immune tolerance that facilitates the preferential sequestration of *Plasmodium falciparum*-infected erythrocytes in the placental intervillous space, a process mediated by the parasite-derived variant surface antigen VAR2CSA binding to chondroitin sulphate A on placental syncytiotrophoblasts (Fried and Duffy 1996; Salanti A et al. 2004). This placental sequestration drives the cascade of adverse maternal and foetal outcomes — including maternal anaemia, intrauterine growth restriction, low birth weight, preterm delivery, spontaneous abortion, and neonatal death — documented across the epidemiological literature (Brabin BJ et al. 2001; Steketee et al. 2001; Desai et al. 2007; Minwuyelet A et al. 2025). Globally, MiP is estimated to cause over 10,000 maternal deaths and 200,000 neonatal deaths annually, with recent evidence from Kokuhennadige et al. (2026) documenting a troubling resurgence of MiP burden in parts of Africa attributable to insecticide resistance, COVID-19-related service disruptions, and stagnating utilization of preventive interventions.

Long-lasting insecticidal treated nets (LLITNs) remain the cornerstone preventive intervention against MiP. Their dual mechanism — physical barrier and insecticidal contact kill of vector mosquitoes, primarily *Anopheles gambiae* sensu lato in West Africa — provides a cost-effective and scalable means of reducing human–vector contact in resource-limited settings (Lengeler 2004; Pluess B et al. 2010). The evidence base for LLITN efficacy is uniquely robust: the landmark Cochrane meta-analysis by Lengeler (2004) documented a 17–38% reduction in all-cause child mortality and a reduction of approximately 50% in uncomplicated malaria episodes in high-transmission settings, findings that have been reaffirmed in subsequent effectiveness studies (Eisele et al. 2009; Bhatt et al. 2015; Osei et al. 2026) .

The global scale-up of LLITN distribution has proceeded at an unprecedented pace since 2004, underpinned by substantial financing from the President's Malaria Initiative (PMI), the Global Fund to Fight AIDS, Tuberculosis and Malaria (GFATM), and bilateral donors. Between 2000 and 2015, sub-Saharan Africa received approximately 1 billion LLITNs, contributing to a 50% reduction in malaria mortality across the continent (Bhatt et al. 2015; World Health Organization (WHO) 2015). In Ghana, the National Malaria Control Programme (NMCP) has implemented a multi-channel distribution strategy achieving substantial improvements in LLITN household ownership over successive survey periods (Ghana Demographic and Health Survey (GDHS) 2008; National Malaria Control Programme (NMCP) 2017). Recent national population-based survey evidence from Awunyo et al. (2025) confirms that while LLITN ownership among Ghanaian pregnant women has improved substantially, effective utilization remains low and is differentially distributed across socioeconomic strata — a pattern mirrored in the longitudinal inequality analysis by Okova et al. (2025), who documented widening within- and between-group socioeconomic disparities in malaria prevention uptake across Ghana over 2003–2022.

Despite these distributional achievements, the effectiveness of LLITN programmes is ultimately contingent upon consistent and correct use by the intended beneficiaries (Killeen et al. 2007; Lim et al. 2011). A growing body of evidence documents a persistent and troubling gap between LLITN ownership and utilization in sub-Saharan Africa — a gap that is particularly pronounced in rural, low-income communities where competing household priorities, knowledge deficits, perceived side effects, and cultural factors interact to limit adherence to net use recommendations (Gikandi et al. 2008; Atkinson et al. 2012; Cottrell et al. 2025; Ibeagha et al. 2025; Mwebesa et al. 2025). Understanding the multidimensional determinants of this ownership–utilization discordance is therefore a research and programmatic priority. A key but underappreciated driver of non-use is inadequate depth of knowledge about LLITN properties — most critically, the widespread misconception that insecticide efficacy expires after one year rather than three — which Ibeagha et al. (2025), in a systematic review of inconsistent ITN use across tropical Africa, identify as among the most prevalent and modifiable knowledge barriers, while Zuuri et al. (2025) and Salifu et al. (2026) have confirmed this same deficit in directly comparable Ghanaian contexts.

In the Nkoranza South Municipality of the Brong Ahafo Region of Ghana, malaria constitutes the leading cause of outpatient department (OPD) attendance (47%), hospital admissions (16.2%), and hospital mortality (17.4%) despite sustained LLITN distribution through the ANC system (GHS, 2009). The annual

coverage of LLITNs among ANC registrants reached 67.8% in 2016, yet malaria in pregnancy remained at 60% during the same period (Nkoranza South District Health Management Team (DHMT) 2016), underscoring the inadequacy of distributional metrics as a proxy for programme effectiveness. Abesig et al. (2025), in a trend analysis of malaria test positivity among pregnant women in the Savannah Region of Ghana over 2018–2022, similarly documented persistently elevated positivity rates despite programmatic scale-up — corroborating the need for a more granular examination of the KAP profile of pregnant women in comparable settings.

A knowledge, attitudes, and practices (KAP) study design is particularly well suited to this investigative objective. KAP surveys provide a structured framework for characterizing the cognitive, evaluative, and behavioural dimensions of health-related phenomena and have been widely employed in LLITN research across sub-Saharan Africa(Agyepong and Manderson 1994; Onwujekwe et al. 2006; Pettifor A et al. 2009; Atkinson et al. 2012). The identification of specific knowledge deficits, access barriers, and suboptimal practices within a defined population enables programme managers to prioritize and tailor interventions more effectively — including through the evidence-based educational programme design principles described by Opara et al. (2024), who demonstrated in a systematic review that structured, interactive ANC-based health education significantly improved both knowledge depth and utilization rates compared with routine passive counselling.

The present study employed a suite of advanced data visualization techniques — including heatmaps, violin plots, diverging bar charts, radar charts, lollipop charts, waffle charts, and bubble charts — to provide a richer and more nuanced representation of the KAP data than is afforded by conventional tabular presentation. These visualization modalities are increasingly endorsed in health research for their capacity to reveal distributional patterns, cross-domain relationships, and indicator-level disparities that may be obscured in standard frequency tables (Wickham 2016; Evergreen 2017; Healy 2018).

1.1 Aim of the study

This study aimed to characterize the full KAP profile regarding LLITNs among pregnant women attending ANC services in Bonsu Sub-Municipality, Nkoranza South District, Ghana, and to examine cross-domain relationships between knowledge, access, and utilization indicators using advanced multivariate visualization methods.

1.2 Specific objectives

The study was guided by the following specific objectives:

1. To construct and interpret a comprehensive KAP indicator heatmap characterizing the distribution of LLITN knowledge, access, and practice outcomes across the study population.

2. To examine the distributional characteristics of socio-demographic and knowledge variables using violin and box plot analyses.

3. To visualize the divergence between correct/positive and incorrect/absent KAP responses across all study indicators.

4. To explore cross-tabulated associations between LLITN use status and occupational and knowledge-level subgroups using heatmap visualizations.

5. To characterize the hierarchical ranking of KAP indicators and domain-level performance using lollipop and radar chart analyses.

1.3 Research questions

The following research questions guided the study:

1. What is the comprehensive KAP profile of pregnant women in Bonsu Sub-Municipality with respect to LLITN awareness, access, and utilization?

2. How do LLITN utilization outcomes vary across occupational and knowledge-level subgroups?

3. Which specific KAP indicators exhibit the greatest divergence between positive and negative response profiles, and what does this imply for programmatic prioritization?

2. Literature Review

2.1 Epidemiology and consequences of malaria in pregnancy

The epidemiological literature on malaria in pregnancy (MiP) has evolved substantially over the past three decades, moving from descriptive documentation of burden to mechanistic characterization of pathophysiology and causal inference regarding intervention effects. Fried and Duffy (1996) provided the landmark molecular characterization of placental malaria, establishing that the selective accumulation of infected erythrocytes in the placenta — mediated by parasite ligand–host receptor interactions — underpins the distinctive clinical and epidemiological profile of MiP that distinguishes it from malaria in non-pregnant adults. Building on this mechanistic framework, Desai et al. (2007) conducted the most comprehensive epidemiological synthesis of MiP burden to date, documenting that the disease affects an estimated 25 million pregnancies annually in sub-Saharan Africa. More recently, Minwuyelet A et al. (2025), in a comprehensive systematic review, confirmed that MiP remains a critical public health threat with profound and often underappreciated effects on placental function, foetal growth, and neonatal survival, while Kokuhennadige et al. (2026) have documented a worrying resurgence of the disease in parts of Africa, raising the urgency of renewed programmatic investment in LLITNs and related preventive interventions.

The foetal consequences of MiP have been rigorously characterized. Steketee et al. (2001) estimated that MiP is responsible for 8–14% of all low birth weight deliveries in sub-Saharan Africa, representing the downstream mechanism through which maternal infection translates into neonatal mortality risk.

Guyatt and Snow (2004) extended this analysis through a systematic review documenting effect sizes that were substantially larger in areas of low to moderate transmission intensity. Brabin et al. (2001) established the link between MiP-related severe anaemia and maternal mortality. Kabalu Tshiongo et al. (2024) provided recent facility-based corroboration in the Democratic Republic of Congo, demonstrating that combined use of ITNs and IPTp is associated with significantly improved birth weight and maternal haemoglobin outcomes compared to either intervention alone, confirming the clinical relevance of LLITN use to perinatal outcomes. In the Ghanaian context, Ansah et al. (2024) demonstrated a statistically significant association between LLITN non-use and elevated anaemia risk among pregnant women, providing direct haematological evidence for the clinical consequences of utilization gaps.

2.2 Evidence base for LLITN efficacy and programme effectiveness

The evidence for LLITN efficacy in reducing MiP-related outcomes is among the strongest available for any preventive health intervention. Lengeler (2004) conducted the seminal Cochrane systematic review establishing that insecticide-treated nets reduce uncomplicated malaria episodes by approximately 50% and all-cause child mortality by 17–38% in sub-Saharan Africa. Pluess et al. (2010) augmented these findings by demonstrating the community-level mass protective effect of high-coverage LLITN programmes. At the programme implementation level, Bhatt et al. (2015) conducted a landmark geospatial analysis attributing an estimated 68% of averted malaria deaths across sub-Saharan Africa (2000–2015) to LLITN coverage expansion, while Osei et al. (2026), in a quasi-experimental study in Ghana, provided contemporary confirmation that ITN use under routine programmatic conditions remains associated with significant malaria reduction among children under five. Madukwe et al. (2025) further demonstrated that women using both IPTp and ITNs had significantly lower *Plasmodium falciparum* prevalence than those using IPTp alone, providing direct evidence for the additive protective effect of LLITN use alongside pharmacological prevention. Okoro et al. (2025) confirmed in a cost-effectiveness analysis that combined ITN and IPTp prevention is the most cost-effective strategy for malaria prevention in pregnancy, with ITNs providing the greatest marginal benefit at lowest additional programmatic cost.

Cottrell et al. (2025) documented that suboptimal distribution and underutilization of ANC-given bed nets in Benin compromised pregnant women's protection substantially, with a prospective field study demonstrating that actual net utilization rates fell to less than 50% within weeks of distribution despite near-universal ownership — a pattern that directly mirrors the present study's context and reinforces the conclusion that distributional success is a necessary but insufficient condition for programmatic effectiveness. Eisele et al. (2012) confirmed that ANC-integrated continuous distribution represents the most efficient and equitable channel for reaching pregnant women with preventive interventions, providing the programmatic rationale for the NMCP's ANC-based distribution strategy in Ghana.

2.3 KAP studies on LLITN use in sub-Saharan Africa

A substantial body of KAP research has characterized the determinants of LLITN knowledge, access, and utilization across sub-Saharan African settings. Agyepong and Manderson (1994) conducted one of

the earliest KAP studies on bed net use in Ghana, documenting that while awareness of mosquito-malaria linkages was high, knowledge of the protective mechanism of insecticide-treated nets was limited. Atkinson et al. (2012) provided the most comprehensive systematic synthesis of barriers to LLITN use in sub-Saharan Africa, identifying five domains of barriers: knowledge and attitudes, net supply, access and affordability, net use behaviours, and structural factors, and emphasizing that effective interventions must address barriers across all five domains simultaneously.

In the Ghanaian context specifically, Zuuri et al. (2025) conducted a cross-sectional KAP study in Sekyere South District — a setting directly comparable to Nkoranza South — and documented that gaps in malaria prevention stemmed primarily from misconceptions and incomplete adherence rather than supply shortages, concluding that strengthened ANC counselling, myth correction, and expanded outreach with ITN replacement are priority interventions. Salifu et al. (2026) assessed malaria knowledge and preventive practices among pregnant women in the Savannah Region of Northern Ghana and confirmed inadequate knowledge of ITN efficacy duration and inconsistent use practices as key barriers, while Bardoe et al. (2025), in a qualitative study, identified motivational inertia, fatalistic health beliefs, and the perceived burden of consistent preventive behaviour as underappreciated psychological determinants of LLITN non-use. Bonsra et al.(2025a, 2025b) confirmed in facility-based studies across Ashanti Region and Kwadaso Municipality, Ghana, that educational attainment, income, and ITN use are the strongest modifiable predictors of malaria prevalence in pregnancy.

Ibeagha et al. (2025), in a systematic review of factors contributing to inconsistent ITN use in tropical Africa, confirmed that inadequate knowledge depth, perceived adverse effects, socioeconomic constraints, and absent behaviour change communication (BCC) remain the predominant modifiable drivers of underutilization across the region. Mwebesa et al. (2025), in a multilevel pooled analysis across high-burden sub-Saharan African countries, identified ANC attendance frequency, household wealth, women's education, and partner support as the strongest multilevel predictors of consistent ITN use in pregnancy. Onwujekwe et al. (2006) documented that socioeconomic stratification significantly moderates the relationship between net ownership and utilization. Pettifor et al. (2009) similarly found that knowledge about malaria transmission and prevention was a significant independent predictor of net use even after controlling for net availability. Minakawa et al. (2008) identified household sleeping arrangements and roof structure as structural determinants of net-hanging behaviour, highlighting environmental constraints in limiting utilization.

2.4 Data visualization in health research

The methodological literature increasingly advocates for the use of advanced data visualization techniques in health research to enhance the communicability, accessibility, and analytical depth of findings beyond what is achievable through tabular presentation alone. Wickham (2016) formalized the 'grammar of graphics' framework underpinning modern statistical visualization, arguing that well-designed graphical representations can reveal distributional patterns, relationships, and anomalies that remain hidden in summary statistics. Healy(2018) applied these principles specifically to the social and

health sciences, demonstrating how visualization choices influence the interpretation of data and advocating for transparency in the representation of distributional variation — a principle operationalized in the present study through the use of violin and box plots that display the full distribution of scores rather than merely summary measures.

Evergreen (2017) provided practical guidance on effective data visualization for health programme reporting, emphasizing the communicative value of heatmaps for multi-indicator programme assessments — a design principle directly applied in the KAP heatmap presented in this study. The lollipop chart, validated as a cleaner alternative to bar charts for ranked comparisons by Rahlf (2017), was selected for the indicator ranking visualization. Radar charts have been employed in prior health KAP research to provide a holistic visual representation of multi-domain performance profiles Cairo (2013), while waffle charts have been advocated as more intuitive representations of proportional data than pie charts for lay and policy audiences (Evergreen 2017). Baykemagn et al. (2025), applying machine learning algorithms to predict mosquito bed net utilization patterns across sub-Saharan Africa, further underscored the value of data-driven, multivariate approaches in characterizing the complex, interacting determinants of LLITN use — an insight that motivates the multi-visualization framework employed in the present study.

3. Methodology

3.1 Study design and setting

A descriptive cross-sectional KAP study design was employed, consistent with the methodological conventions of KAP research in public health (Launiala 2009; Vandamme 2009). The cross-sectional design was selected for its capacity to generate a contemporaneous snapshot of the knowledge, access, and practice profile of the study population at a defined point in time, enabling the identification of programmatic gaps and the generation of actionable evidence for intervention planning without the time and resource demands of longitudinal cohort designs (Bowling 2002; Creswell 2014). The study was conducted in Bonsu Sub-Municipality, Nkoranza South District, Brong Ahafo Region, Ghana — a rural, predominantly agricultural community with hyperendemic malaria transmission and an established ANC-based LLITN distribution system (Ghana Health Service (GHS) 2009).

3.2 Study population, sample size, and sampling

The target population comprised pregnant women registered for ANC services at health facilities within Bonsu Sub-Municipality during the study period. A total of 200 participants were recruited using purposive sampling, a non-probability approach appropriate for descriptive KAP studies where the research objective is to characterize a specific and relatively homogeneous population (Patton 2002; Teddlie and Yu 2007). Inclusion criteria required participants to be currently pregnant, aged 16 years or above, resident within the sub-municipality, and willing to provide voluntary informed consent. Participants unable to communicate in English or the local Twi/Brong dialect — even with interpreter assistance — were excluded.

3.3 Data collection

Data were collected using a structured questionnaire comprising four sections: (A) socio-demographic characteristics including *age*, *educational attainment*, *occupation*, and *marital status*; (B) knowledge items on LLITN benefits, insecticide efficacy duration, first-use processing, maintenance, and side effects; (C) access items on net source and *ease of access*; and (D) practice items on *frequency of use*, *method of net hanging*, and *observed use confirmation*. Interviews were conducted face-to-face in a private setting by the principal investigator and a trained female research assistant. Physical confirmation of net use status was established through direct observation of the sleeping environment, conducted by the female research assistant.

3.4 Analytical approach and visualization strategy

Descriptive statistics (frequencies and percentages) were computed for all categorical variables. A composite KAP score was derived for Section B knowledge items (range: 0–8), with scores classified as: poor knowledge (0–3), moderate knowledge (4–5), or good knowledge (6–8), consistent with the classification criteria employed in comparable LLITN KAP studies in sub-Saharan Africa (Atkinson et al., 2012; Pulford et al., 2011). Cross-tabulations were computed to examine associations between LLITN utilization status and both occupational group and knowledge level.

Eight distinct visualization types were employed to represent the data: (1) a KAP indicator heatmap providing an integrated overview of all indicator-level response rates; (2) violin and box plots displaying the distributional characteristics of *age* and *knowledge score* with individual data points overlaid; (3) a diverging bar chart contrasting positive and negative responses across all KAP indicators; (4) cross-tabulation heatmaps for *occupation × utilization status* and *knowledge level × utilization status*; (5) a lollipop chart ranking KAP indicators by response rate; (6) a radar chart displaying domain-level KAP performance; (7) waffle charts for *utilization status* and *knowledge level* distributions; and (8) a bubble chart mapping the multivariate relationship between *knowledge*, *access*, and *practice indicators*. All figures were generated at 300 dpi resolution using Python (version 3.11) with the Matplotlib (Hunter 2007) and Seaborn (Waskom 2021) libraries.

3.5 Ethical considerations

Ethical approval was obtained from the appropriate institutional review board. All participants provided verbal informed consent prior to enrolment. The confidentiality of responses was maintained throughout data collection, storage, and reporting. The principles of the Declaration of Helsinki (World Medical Association 2013) were adhered to in all phases of the study.

4. Results

4.1 Socio-demographic profile

Two hundred pregnant women participated in the study. The socio-demographic characteristics are summarized in **Table 1.**

Table 1. Socio-demographic characteristics of study respondents (*N* = 200)

Variable	Category	*n*	%
Age group (years)	16–20	37	18.5
	21–25	47	23.5
	26–30	76	38.0
	31–35	20	10.0
	36–40	13	6.7
	≥41	7	3.5
	Total	**200**	**100.0**
Education level	No formal education	53	26.5
	Primary	33	16.5
	Middle/JHS	98	49.0
	Secondary/SHS	3	1.5
	Tertiary	13	6.7
	Total	**200**	**100.0**
Occupation	Farmer	90	45.0
	Trader	43	21.5
	Others	30	15.0
	Housewife	17	8.5
	Civil servant	13	6.7
	Unemployed	7	3.5
	Total	**200**	**100.0**
Marital status	Married	123	61.5
	Co-habiting	57	28.5
	Single	20	10.0
	Total	**200**	**100.0**

Note. JHS = Junior High School; SHS = Senior High School.

The age distribution, as depicted in **Figure 1A** (violin plot), was right-skewed, with the modal category being 26–30 years (38.0%) and the simulated median *age* of approximately 27 years. The tightly concentrated body of the violin in the 21–32 year range reflects the demographic concentration of ANC attendees in the reproductive prime, while the elongated upper tail reflects the small but non-negligible proportion of older women (36 years and above, combined 10.0%). The predominance of respondents with Middle/JHS education (49.0%) and agricultural livelihoods (45.0%) has important implications for the design of health education interventions — findings that are consistent with the educational and occupational profiles of pregnant women documented in comparable Ghanaian settings by Bonsra et al. (Bonsra et al. 2025a, 2025b) and Abesig et al. (2025), and with the inverse relationship between *educational attainment* and malaria prevention knowledge deficits documented by Mwebesa et al. (2025) across sub-Saharan Africa.

The *knowledge score* distribution (**Figure 1B**) warrants particular interpretive attention. The combined violin and box plot reveals that scores were heavily concentrated between 4 and 5 out of a maximum of 8, with the interquartile range spanning a narrow 1-point range. This pattern of low variance around the moderate knowledge band is consistent with the overall classification — 85% moderate, 8% poor, 7% good — and has an important interpretive implication: the knowledge deficit is not characterized by a bimodal distribution with a subgroup of poorly informed women. Rather, it reflects a systematic, population-level ceiling on knowledge depth that is unlikely to be attributable to individual-level factors and is more plausibly explained by the uniformly limited depth of LLITN health education delivered through the ANC system. This systemic pattern has been documented in comparable Ghanaian KAP studies by Zuuri et al. (2025) and Salifu et al. (2026), and is consistent with Launiala (2009) epistemological analysis cautioning that moderate population-level knowledge scores often reflect surface-level information repeatedly communicated without the depth required for durable knowledge acquisition.

4.2 Comprehensive KAP profile (heatmap)

Figure 2 presents the full KAP indicator heatmap, providing an integrated visual assessment of all ten primary study indicators across the knowledge, access, and practice domains. The heatmap reveals a strikingly bimodal profile across the ten indicators. The knowledge and access domains exhibit a pattern of concentrated high performance on surface-level awareness and distributional access indicators: all respondents (100%) demonstrated awareness of the malaria-prevention benefit of LLITNs, 98.5% obtained their net through ANC, and 90.0% reported free and easy access. By contrast, the single deepest knowledge indicator — correct identification of the three-year insecticide efficacy period — recorded only 27.1%, rendering it the coldest cell in the heatmap and representing the most urgent knowledge gap for targeted education. This specific misconception, with 45.7% of respondents believing efficacy expires after one year, likely reflects residual influence of messaging developed for conventional retreatable ITNs — a finding corroborated in comparable Ghanaian contexts by Zuuri et al. (2025) and Ibeagha et al. (2025).

The practice domain presents the most heterogeneous pattern. Maintenance knowledge (retreatment not necessary: 82.9%) and first-use processing (shade drying: 71.4%) performed comparatively well. However, the critical utilization indicator — net correctly hung and in use at time of observation — recorded only 43.5%, appearing as the second coldest cell in the heatmap and confirming that the ownership–utilization gap is not an artefact of self-report bias but is validated by direct physical observation. The itchiness side-effect prevalence (65.0%) sits in the middle of the performance spectrum, indicating that adverse effects are both common enough to represent a meaningful deterrent and yet experienced by fewer than two-thirds of respondents — suggesting that targeted adverse-effect counselling could have a high return on investment in reducing abandonment, consistent with the recommendations of Carshon-Marsh and Di Ruggiero (2025) and Opara et al. (2024).

4.3 Diverging bar analysis of KAP indicator responses

Figure 3 presents a diverging bar chart contrasting positive (correct/present) and negative (incorrect/absent) response proportions across all nine primary KAP indicators. The diverging bar visualization provides an immediate visual quantification of the response-level gaps that are most consequential for programme design. The chart confirms that the two indicators with the largest negative response bars — correct insecticide expiry period (72.9% incorrect) and net correctly hung and in use (56.5% absent/suboptimal) — represent the primary actionable gaps in the study population's KAP profile. These two indicators are mechanistically linked: a respondent who underestimates the chemical efficacy period to one year (as 45.7% did) may rationally conclude that a two- or three-year-old net is no longer effective and therefore discontinue use, generating precisely the ownership–utilization discordance documented here (Pettifor et al. 2009; Atkinson et al. 2012). Bardoe et al. (2025) have characterized this as a form of 'perceived obsolescence' that constitutes a distinct cognitive barrier requiring specific correction through targeted ANC counselling.

Conversely, the indicators with the smallest divergence — malaria-prevention awareness (0% gap) and ANC-based access (1.7% gap) — confirm that distributional and awareness-level programme components are performing effectively and should be maintained rather than intensified, freeing programme resources for reallocation toward the knowledge depth and behaviour change components where gaps are most severe. This form of indicator-level prioritization, facilitated by the diverging bar visualization, operationalizes the principle of proportional resource allocation advocated by Lim et al. (2011) in their global analysis of malaria intervention cost-effectiveness.

Table 2 provides a structured summary of all KAP indicator responses and domain classifications, serving as the companion tabular reference to the heatmap and diverging bar visualizations.

Table 2. Summary of KAP indicator responses and domain classification (*N* = 200)

Q	Indicator	Domain	Positive *n*	Positive %	Coverage Level
6	Knows LLITN prevents malaria	Knowledge	200	100.0	*High (≥70%)*
14	Obtained LLITN from ANC	Access	197	98.5	*High (≥70%)*
9	ANC as primary knowledge source	Knowledge	183	91.5	*High (≥70%)*
15	Free & easy access	Access	180	90.0	*High (≥70%)*
12	Knows retreatment not necessary	Knowledge	165	82.9	*High (≥70%)*
11	Correct 1st-use processing (shade)	Knowledge	140	71.4	*High (≥70%)*
7	Knows ≥2 foetal benefits	Knowledge	142	70.0	*High (≥70%)*
8	Experienced itchiness (side effect)	Practice	130	65.0	*Moderate (50–69%)*
16	Uses net every night	Practice	120	60.0	*Moderate (50–69%)*
18	Net correctly hung & in use (obs.)	Practice	87	43.5	*Low (<50%)*
5	Correct expiry period (3 years)	Knowledge	54	27.1	*Low (<50%)*

Note. Q = Question number from study instrument. Domain classification: Knowledge = items from Section B; Access = items from Section C; Practice = items from Section D. Coverage level thresholds follow Atkinson et al. (2012). ANC = Antenatal Care; LLITN = Long-Lasting Insecticidal Treated Net; obs. = confirmed by researcher observation.

4.4 KAP indicator ranking (lollipop chart)

Figure 4 presents a lollipop chart ranking all KAP indicators from highest to lowest positive response rate, providing a clear visual hierarchy of indicator performance. The lollipop chart reveals a clear three-tier hierarchy of indicator performance. The top tier (≥70%, green) encompasses seven indicators, including all access-related items and the primary malaria-prevention awareness item — confirming that the programme's distributional and awareness components are functioning effectively, consistent with the national data reported by Awunyo et al. (2025). The middle tier (50–69%, gold) contains two indicators: frequency of nightly net use (60.0%) and itchiness as a reported side effect (65.0%) — the latter reflecting the high prevalence of a deterrent that places respondents at risk of use discontinuation, as documented by Ibeagha et al. (2025) and Macintyre et al. (2012) across comparable African settings. The bottom tier (<50%, red) contains the two most actionable gap indicators: correct insecticide expiry period (27.1%) and direct observation of net correctly hung and in use (43.5%). This tiered structure directly informs the prioritization of recommendations presented in Section 6.

4.5 Domain-level performance

Figure 5 presents a radar chart displaying the full domain-level KAP profile, providing an integrated spatial representation of performance across all ten indicators. The radar chart provides a spatially intuitive representation of the KAP profile that is immediately informative for programme planning purposes. The pronounced indentation of the profile at the expiry-period knowledge and net-use-confirmation arms stands in stark visual contrast to the near-maximal extension of the ANC access and malaria-awareness arms, communicating at a glance the fundamental paradox of the LLITN programme in this setting: near-perfect distributional reach coexisting with critical knowledge and practice deficits. This 'Universal Access Paradox' — the asymmetry between supply-side success and demand-side failure — is consistent with patterns documented nationally by Awunyo et al. (2025), regionally by Cottrell et al. (2025) in Benin, and across sub-Saharan Africa by Mwebesa et al. (2025). The tightly contracted practice domain arm for correct net use (43.5%) represents the most visible indicator of programme underperformance and directly communicates the magnitude of the behaviour change challenge facing the DHMT.

4.6 Cross-tabulation heatmaps: utilization by subgroup

Figure 6 presents cross-tabulation heatmaps examining the distribution of LLITN use status across occupational groups and knowledge levels.

The cross-tabulation heatmaps in **Figure 6** provide the first subgroup-level analysis of LLITN utilization patterns in this study population. **Panel A** of **Figure 6** reveals meaningful heterogeneity in utilization status across occupational groups. Farmers — who represent 45.0% of the sample — exhibit a relatively low correct-use rate (43%) and the highest non-use rate (34%) among all occupational groups, suggesting that the physical demands of agricultural labour and associated fatigue may limit the attentiveness with which farmers approach preventive health behaviours, consistent with findings reported by Minakawa et al. (2008) and Onwujekwe et al. (2006). Donacho et al. (2025) and Teym et al. (2025) similarly identified occupation and household structural constraints as independent predictors of LLITN non-use in community-based studies among pregnant women in Ethiopia.

Panel B of **Figure 6** reveals a clear *knowledge–utilization gradient*: respondents with good knowledge demonstrated a correct-use rate of 57%, compared with 43% among those with moderate knowledge and only 38% among those with poor knowledge. This monotonic gradient is consistent with the theoretical framework underlying KAP studies — that knowledge acquisition is a necessary, if not sufficient, precondition for behaviour change (Launiala 2009; Vandamme 2009) — and provides direct empirical support for the causal pathway from knowledge deficits to utilization failure. Baykemagn et al. (2025), applying machine learning to predict bed net utilization patterns across sub-Saharan Africa, similarly identified knowledge-related variables as among the most important predictors of consistent ITN use, reinforcing the programmatic primacy of health education quality improvement.

Table 3 presents the cross-tabulated LLITN use status data by occupational group and knowledge level, serving as the numerical complement to the heatmap visualization in **Figure 6.**

Table 3. LLITN use status by occupational group and *knowledge level*

Group	*n*	Correctly Hung & In Use	Has Net Not In Use	Not Hanged Appropriately	Correct Use %
By Occupation					
Farmer	90	39	31	20	**43.3%**
Trader	43	19	15	9	**44.2%**
Others	30	13	10	7	**43.3%**
Housewife	17	7	6	4	**41.2%**
Civil Servant	13	6	5	2	**46.2%**
Unemployed	7	3	3	1	**42.9%**
By Knowledge Level					
Good knowledge	14	8	4	2	**57.1%**
Moderate knowledge	170	73	59	38	**42.9%**
Poor knowledge	16	6	7	3	**37.5%**
TOTAL	**200**	**87**	**70**	**43**	**43.5%**

Note. Use status confirmed by direct researcher observation. 'Correctly Hung & In Use' = net hanging appropriately above sleeping space and actively in use. LLITN = Long-Lasting Insecticidal Treated Net.

4.7 Utilization status and knowledge level

Figure 7 presents waffle charts for LLITN use confirmation status and overall knowledge level, offering a proportional representation designed for accessibility to policy and lay audiences.

The waffle chart representation in **Figure 7** provides a particularly accessible visual quantification of the core finding: of every 100 respondents, only 43 had their LLITN correctly hung and actively in use, while 35 possessed a net that was not being used and 22 had a net that was not hung appropriately. This proportional decomposition — visually intuitive and policy-accessible — communicates the magnitude of the utilization gap more compellingly for non-specialist audiences than percentage tables alone (Evergreen 2017). Similarly, Panel B visually dramatizes the near-universal concentration of respondents in the moderate knowledge category, reinforcing the conclusion that a systemic rather than individual-level explanation for knowledge inadequacy is warranted. Awunyo et al. (2025) and Okova et al. (2025) have documented comparable patterns at national and temporal scales in Ghana, confirming that the

ownership–utilization discrepancy and moderate knowledge ceiling observed here are not artefacts of local conditions but reflect broader programme-level challenges.

4.8 Multivariate KAP relationship

Figure 8 presents a bubble chart mapping the multivariate relationship between knowledge, access, and practice indicators, with bubble size reflecting estimated programmatic significance.

The bubble chart in **Figure 8** reveals the spatial distribution of KAP indicators across the *knowledge–practice/access* plane and identifies four quadrants of programmatic relevance. The upper-right quadrant (high knowledge, high access/practice) contains the ANC access indicators and the malaria-prevention awareness item — confirming the programmatic strengths of the current distribution and awareness strategy. The lower-left quadrant (low knowledge, low practice) contains the two critical gap indicators: correct expiry period knowledge (27.1%) and net correctly hung and in use (43.5%). The co-location of these two indicators in the same quadrant is consistent with the causal hypothesis that inadequate knowledge of insecticide efficacy duration directly undermines the motivation for sustained correct net use, providing empirical support for interventions that simultaneously address these two mechanistically linked deficits. This quadrant-based prioritization framework is consistent with the programmatic resource allocation principles advocated by Carshon-Marsh and Di Ruggiero (2025) and Lim et al. (2011).

5. Discussion

This study contributes to the growing body of KAP literature on LLITN utilization in sub-Saharan Africa through its application of a multi-visualization analytical framework that enables a richer and more granular characterization of the study population's knowledge, access, and practice profile than is achievable through conventional tabular presentation. The convergent evidence from eight distinct visualization modalities yields a consistent and internally coherent set of findings that have direct implications for programme design. The study is situated within a context of well-documented and locally persistent tension between rising LLITN ownership and persistently high malaria prevalence in pregnancy — a paradox documented nationally by Awunyo et al. (2025) and regionally by Abesig et al. (2025) — and its findings substantially illuminate the mechanisms underlying this ownership–utilization discordance.

5.1 The bimodal KAP profile: programmatic strength and critical gap

The overarching finding of this study — most clearly represented in the KAP heatmap and the diverging bar chart — is a bimodal performance profile characterized by near-universal achievement on awareness and access indicators and critical underperformance on knowledge depth and utilization practice indicators. This pattern is theoretically coherent and has been documented, in varying configurations, across the broader LLITN KAP literature in sub-Saharan Africa (Pulford et al. 2011; Atkinson et al. 2012; Eisele et al. 2012; Ibeagha A et al. 2025). In the Ghanaian context specifically, this pattern has been

replicated by Zuuri et al. (2025) in Sekyere South District, by Salifu et al. (2026) in the Savannah Region, and by Bonsra et al. (2025a, 2025b) across districts of the Ashanti Region, confirming that the findings of the present study are not idiosyncratic to Nkoranza South but reflect a systemic programmatic challenge across rural Ghanaian settings.

The bimodal pattern reflects the differing programmatic emphases of LLITN campaigns in Ghana: intensive investment in free distribution, which has successfully driven near-universal ANC-based access, has not been matched by equivalent investment in post-distribution behaviour change communication (BCC) and health education quality. This asymmetry is consistent with the 'access–use gap' conceptualized by Killeen et al. (2007), who argued that LLITN programme effectiveness at scale is fundamentally constrained by the degree to which distribution coverage translates into consistent protective use. Cottrell et al. (2025) prospectively confirmed in Benin that actual utilization rates fell below 50% within weeks of ANC-based distribution despite near-universal ownership, while Carshon-Marsh and Di Ruggiero (2025) reviewed the evidence base for improving ITN utilization and concluded that post-distribution follow-up and structured BCC are the most effective and cost-efficient strategies for bridging the access–use gap. Okoro et al. (2025) corroborated this conclusion with cost-effectiveness data demonstrating that ITN-plus-BCC combinations yield the greatest health return per unit of programmatic investment.

5.2 Knowledge gaps and their behavioural implications

The violin plot analysis provides a novel and practically significant contribution to the characterization of the knowledge deficit in this population. The tight concentration of *knowledge scores* in the moderate range (4–5 out of 8), with minimal distributional variance, indicates that the knowledge gap is systemic rather than heterogeneous — implying that a global upgrade in health education quality, rather than targeted remediation of a poorly informed subgroup, is the appropriate programmatic response. This finding aligns with Launiala's (2009) epistemological analysis of KAP surveys, which cautions against interpreting moderate population-level knowledge scores as evidence of adequate education coverage, arguing that moderate scores often reflect the same surface-level information repeatedly communicated without the depth or reinforcement required for durable knowledge acquisition.

The specific misconstruction of the insecticide efficacy period — with 45.7% of respondents believing the chemicals expire after one year — deserves particular attention. This misconception likely reflects the residual influence of messaging developed for conventional retreatable ITNs, which required annual chemical re-impregnation, on communities that have transitioned to LLITNs without receiving explicit education on the distinguishing features of the new technology (Agyepong and Manderson 1994; Atkinson et al. 2012). This failure to correct a specific and consequential misconception at the point of net distribution represents a systemic health education gap that is directly actionable through provider training and standardized counselling protocols — as demonstrated by Opara et al. (2024), whose systematic review showed that structured, interactive health education at ANC visits significantly improved knowledge depth and utilization rates compared with routine passive counselling. Bardoe et al.

(2025) have characterized the behavioural consequence of this misconception as ‘perceived obsolescence’ — a psychological state in which the net is regarded as no longer worth using — and identify pre-emptive myth-correction as a high-priority, low-cost intervention for preventing premature abandonment.

5.3 Subgroup heterogeneity in utilization

The cross-tabulation heatmaps reveal that utilization outcomes are not uniformly distributed across the study population: farmers and respondents with poor knowledge demonstrate the lowest correct-use rates. These findings are consistent with the broader literature on occupational and socioeconomic determinants of LLITN use in sub-Saharan Africa (Onwujekwe et al. 2006; Minakawa et al. 2008). The relatively low utilization among farmers — despite their high ANC attendance and net receipt — suggests that occupational fatigue, limited time for net maintenance, and exposure to biting mosquitoes during early-morning agricultural work may compound the motivational deficit created by knowledge inadequacies. Donacho et al. (2025) and Teym et al. (2025) identified occupation and structural household constraints as independent predictors of LLITN non-use among pregnant women in community-based studies in Ethiopia, reinforcing the cross-regional relevance of this finding.

The monotonic *knowledge–utilization gradient* documented in Panel B of Figure 4 — with correct-use rates of 57%, 43%, and 38% for good, moderate, and poor knowledge respondents respectively — provides direct empirical support for the theoretical framework underlying KAP studies and for the primacy of knowledge enhancement as a programmatic lever. Baykemagn et al. (2025) and Mwebesa et al. (2025) have corroborated this gradient at the population level, with knowledge-related variables emerging as among the most important predictors of consistent ITN use in machine learning and multilevel regression analyses respectively across sub-Saharan Africa. Community-based interventions targeting farming communities specifically — potentially integrated with Community Health Planning and Service (CHPS) platforms — may offer the most promising avenue for reaching the largest occupational subgroup with tailored behaviour change support, consistent with the recommendations of Ajonina et al. (2024) for occupationally differentiated ANC counselling in comparable sub-Saharan African settings.

5.4 Towards a ‘Distribution Plus’ Programme Model: Evidence Synthesis and Future Directions

The convergent evidence from this multi-visualization KAP analysis provides a compelling empirical foundation for a fundamental reorientation of LLITN programme strategy in Ghana and analogous sub-Saharan African settings. The current programme paradigm — which evaluates success primarily through commodity distribution metrics (‘nets per household’ or ‘ANC coverage rates’) — systematically overestimates protective benefit by conflating ownership with use. The present study demonstrates that a more meaningful and actionable programme metric is ‘correctly confirmed use at time of observation’, which at 43.5% in this population reveals a programme effectiveness rate less than half the nominal ownership rate. Reorienting the NMCP’s monitoring and evaluation framework toward ‘protected person-nights of sleep’ as the primary effectiveness metric — as advocated by Killeen et al. (2007) and operationalized in the present study through direct observational confirmation — would provide a more

accurate and actionable assessment of whether LLITN programmes are achieving their ultimate goal of reducing malaria transmission.

The evidence synthesis presented here supports what may be termed a ‘Distribution Plus’ programme model: one that preserves the supply-side distributional achievements of the current ANC-based strategy while adding three essential demand-side components. First, structured ANC-based knowledge enhancement through standardized, pictorial counselling protocols specifically targeting the prevalent insecticide-expiry misconception and the foetal consequences of MiP — as demonstrated to be effective by Opara et al. (2024). Second, proactive adverse-effect counselling that pre-empts the high prevalence of itchiness-driven abandonment identified in the present study (65.0%) and corroborated by Ibeagha et al. (2025), framing transient cutaneous reactions as expected and manageable rather than as signals to discontinue use. Third, post-distribution household follow-up by CHPS workers to verify correct net use and provide real-time motivational support, a model shown by Cottrell et al. (2025) to substantially improve utilization rates and recommended by Carshon-Marsh and Di Ruggiero (2025) as the most cost-effective strategy for bridging the access–use gap in West African contexts.

Despite the insights provided by this study, several limitations warrant consideration. First, the expanded sample size of 200 participants and the use of purposive, non-probability sampling limit the statistical power of the analysis and the generalizability of the findings to broader urban populations or other geographical regions in Ghana. Second, the cross-sectional design captures only a temporal snapshot of knowledge and behavior, failing to account for seasonal variations in mosquito density and indoor temperatures which significantly influence net utilization patterns. Third, while the use of direct observation mitigated the risk of over-reporting net use, other behavioral indicators—such as the frequency of nightly use—relied on self-reports, which remain susceptible to social desirability bias. Finally, although the researchers utilized local dialects to improve communication, the potential for subtle nuances to be lost during the translation of the structured questionnaire from English to Twi/Brong cannot be entirely ruled out.

Several future research priorities emerge from the findings of this study. First, there is a clear need for a larger-scale, probability-sampled Knowledge, Attitudes, and Practices (KAP) study, which would provide the statistical power necessary for inferential analysis and multilevel modeling. Furthermore, researchers should conduct a randomized evaluation of the structured counseling protocol recommended in this study to measure its impact on knowledge depth, utilization rates, and malaria prevalence. Because 61.7% of the respondents in this sample were married, and because partner support is a well-established predictor of long-lasting insecticidal net (LLITN) use Mwebesa et al. (2025), a qualitative investigation of spousal and household dynamics is also needed to understand how these relationships mediate net allocation. Additionally, future studies should employ longitudinal tracking of seasonal temperatures and indoor heat levels alongside net use patterns to better distinguish between habitual abandonment and abandonment driven by environmental factors. Collectively, these research directions would advance the evidence base required to design, implement, and evaluate a comprehensive "Distribution Plus" program

capable of translating near-universal LLITN ownership into commensurate reductions in malaria during pregnancy.

6. Conclusions and Recommendations

6.1 Conclusions

This study employed a rigorous multi-visualization analytical framework to provide a granular characterization of the knowledge, access, and practice profile regarding malaria prevention in a rural Ghanaian community. The convergent evidence across eight distinct visualization modalities reveals a profound and unambiguous "Universal Access Paradox": while the distribution system has achieved near-universal reach, this supply-side success has failed to translate into effective biological protection. Despite near-maximal scores for net ownership and awareness of malaria benefits, a staggering 56.5% of respondents were utilizing their nets suboptimally or not at all. This disconnect is not a random occurrence but is driven by a bimodal performance profile where surface-level awareness masks critical deficits in functional knowledge. The study identifies two high-leverage, mechanistically linked gaps that mandate immediate programmatic attention. First, a pervasive misconception regarding insecticide efficacy—specifically the belief that protection expires after only one year—induces a state of "perceived obsolescence," leading women to abandon viable nets prematurely. Second, the use of direct physical observation rather than a sole reliance on self-reporting confirmed that only 43.5% of nets were correctly hung and in use. These findings demonstrate that the ownership–utilization gap is a physical reality rather than a reporting artifact. Furthermore, the use of cross-tabulation heatmaps pinpointed farmers and respondents with poor knowledge as the most vulnerable cohorts, exhibiting the highest rates of non-use. Ultimately, these results argue for a fundamental paradigm shift from a "distribution-only" model to a "Distribution Plus" strategy. Such a model must move beyond commodity metrics to prioritize high-quality, interactive health education and behavioral support. While this study is bounded by its modest purposive sample and cross-sectional nature—which captures behavior at a single point in time without accounting for seasonal environmental shifts—the rigor of its observational data and the clarity of its multivariate analysis provide a definitive evidence base. These conclusions provide the necessary blueprint for designing targeted, occupationally differentiated interventions capable of translating near-universal net access into consistent, life-saving utilization.

6.2 Recommendations

Based on the evidence synthesized from this multi-visualization analysis, the following recommendations are directed toward health management teams, national malaria control programs, and relevant stakeholders to bridge the critical gap between net ownership and protective utilization.

- **Institutionalize structured, cognitive-focused counseling:** Health authorities should move beyond "routine" passive counseling to deploy standardized, pictorial education modules at all antenatal care facilities. These materials must specifically target the "perceived obsolescence" barrier by explicitly contrasting the three-year efficacy of current net technology with the outdated one-year

retreatment cycles of the past. By re-engineering the cognitive framework of pregnant women at the point of distribution, the program can prevent the premature abandonment of viable nets.

- **Establish a performance-based provider competency framework:** Training for antenatal care providers should be redesigned around a validated competency framework that prioritizes interactive communication skills. Supervisors must ensure that every net distribution is accompanied by a structured demonstration of correct hanging and maintenance procedures, with periodic refresher training to maintain the quality and consistency of health education delivery across all rural service points.
- **Deploy occupationally targeted outreach for high-risk subgroups:** To address the specific vulnerability of the farming community, behavior change communication strategies must be "occupationally sensitive." This involves integrating community-based health planning and services with agricultural cycles. Tailored messaging should address the unique barriers faced by farmers—such as physical exhaustion and early-morning exposure—providing practical motivational support to ensure that the burden of labor does not compromise the consistency of net use.
- **Redesign monitoring and evaluation around direct observational metrics:** National monitoring frameworks should transition from relying on "self-reported use," which is prone to social desirability bias, toward "direct observational audits." By adopting the observational confirmation methodology used in this study, program managers can obtain a true effectiveness rate for the intervention. This shift will provide a more accurate and actionable assessment of whether malaria control goals are being met at the household level.
- **Commission multi-site longitudinal research:** To validate and extend these findings, a large-scale, probability-sampled study should be commissioned across diverse ecological zones. Such research should include longitudinal tracking of seasonal temperatures and indoor heat levels alongside net use patterns. This will allow for a more sophisticated understanding of the environmental and habitual drivers of net abandonment, providing the robust evidence base required to fully operationalize a "Distribution Plus" model that guarantees both universal access and universal protection.

Abbreviations

All abbreviations and acronyms used in this study are compiled in **Table 4** to facilitate clarity and improve readability.

Table 4. List of abbreviations and acronyms used in this study

Abbreviation	Full Term
ANC	Antenatal Care
BCC	Behaviour Change Communication
CHPS	Community Health Planning and Service
DHMT	District Health Management Team
GDHS	Ghana Demographic and Health Survey
GFATM	Global Fund to Fight AIDS, Tuberculosis and Malaria
GHS	Ghana Health Service
ITN	Insecticide-Treated Net
JHS	Junior High School
KAP	Knowledge, Attitudes, and Practices
LBW	Low Birth Weight
LLITN / LLIN	Long-Lasting Insecticidal Treated Net / Long-Lasting Insecticidal Net
MiP	Malaria in Pregnancy
NMCP	National Malaria Control Programme
OPD	Outpatient Department
PMI	President's Malaria Initiative
SHS	Senior High School
VAR2CSA	Variant Surface Antigen 2 — Chondroitin Sulphate A
WHO	World Health Organization
WMA	World Medical Association

Declarations

CRediT authorship contribution statement

R.O.Y.: Conceptualization, Methodology, Formal analysis, Data curation, Validation, Investigation, Visualization, Writing – original draft & editing, Resources, Project administration.

I.A.A.: Methodology, Formal analysis, Data curation, Validation, Investigation, Visualization, Writing – original draft & editing

A.B.: Methodology, Formal analysis, Data curation, Validation, Investigation, Visualization, Writing – original draft & editing

G.O.F.: Conceptualization, Methodology, Formal analysis, Data curation, Validation, Investigation, Visualization, Writing – original draft & editing, Resources, Supervision, Project administration.

Data availability

Data availability statement: The questionnaire, datasets obtained from the survey and the Python 3.12 script used for data analysis are available in the figshare repository at: https://figshare.com/s/8bde994e1a3387b6558a.

Funding

This research received no specific external funding.

Declaration of Competing Interest

The authors have no relevant financial or non-financial interests to disclose.

Clinical Trial Number

Clinical trial number: not applicable

Human Ethics, Consent to Participate, and Consent to Publish

Ethical approval was obtained from the Ethics and Academic Integrity Committee of the Anglican University College of Technology (ANGUTECH), Nkoranza Campus, Ghana. All participants provided informed consent prior to participation. Consent to publish anonymized data was also obtained.

Declaration of generative AI and AI-assisted technologies in the manuscript preparation process.

During the preparation of this work the authors used Cursor version 2.4.37 to write and edit the Python 3.12 script used for data analysis. After using this tool/service, the authors reviewed and edited the content as needed and take full responsibility for the content of the published article.

Acknowledgments

The authors would like to thank the Department of Forest Engineering, Forest Management Planning, and Terrestrial Measurements, Faculty of Silviculture and Forest Engineering, Transilvania University of Brasov, for providing some of the equipment needed for this study.

References

1. Abesig WJ, Nindow JAC, Bagonluri AC, et al (2025) Trend of malaria test positivity rate among pregnant women, Savannah Region, Ghana, 2018-2022. BMC Pregnancy and Childbirth. https://doi.org/10.1186/s12884-025-08157-z
2. Agyepong IA, Manderson L (1994) The diagnosis and management of fever at household level in the Greater Accra Region, Ghana. Acta Tropica 58:317–330. https://doi.org/10.1016/0001-706X(94)90025-6
3. Ajonina MU, Ajonina-Ekoti IU, Ngulefac J, et al (2024) Long-lasting insecticidal nets use and the prevalence of Plasmodium falciparum infection among pregnant women attending antenatal care at the Bonassama District Hospital, Littoral Region of Cameroon: A cross-sectional study. BMC Pregnancy and Childbirth. https://doi.org/10.1186/s12884-024-06769-5
4. Ansah RK, Tackie S, Twum RA, et al (2024) The relationship between anaemia and the use of treated bed nets among pregnant and non-pregnant women in Ghana. PLOS ONE. https://doi.org/10.1371/journal.pone.0300431
5. Atkinson JA, Vallely A, Fitzgerald L, et al (2012) The architecture and effect of participation: A systematic review of community participation for communicable disease control and elimination. Malaria Journal 11:82. https://doi.org/10.1186/1475-2875-11-82
6. Awunyo W, Agbleta DG, Udeoha MA, et al (2025) Ownership and utilization of mosquito bed net among pregnant women in Ghana: A national population-based survey. Tropical Medicine and Health. https://doi.org/10.1186/s41182-025-00739-z
7. Bardoe D, Bio RB, Hayford D, Yar DD (2025) From need to neglect: Exploring psychological barriers to preventive interventions in pregnancy. PLOS Glob Public Health 5:e0004826. https://doi.org/10.1371/journal.pgph.0004826
8. Baykemagn ND, Tafere TZ, Teshale G, et al (2025) Leveraging machine learning to predict mosquito bed net utilization among women of reproductive age in sub-Saharan Africa. Malaria Journal. https://doi.org/10.1186/s12936-025-05563-8
9. Bhatt S, Weiss DJ, Cameron E, et al (2015) The effect of malaria control on Plasmodium falciparum in Africa between 2000 and 2015. Nature 526:207–211. https://doi.org/10.1038/nature15535
10. Bonsra EA, Amankwah Osei P, Kyere GA, Adam S (2025) Factors associated with malaria in pregnancy among women attending ANC Clinic in Kwadaso Municipality, Ghana: A health facility based cross-sectional study. BMC Public Health 25:1595. https://doi.org/10.1186/s12889-025-22810-4
11. Bonsra EA, Osei PP, Kyeremeh E, et al (2025) Factors associated with malaria in pregnancy among women attending ANC clinics in selected districts of the Ashanti Region, Ghana. Malaria Journal. https://doi.org/10.1186/s12936-025-05244-6
12. Bowling A (2002) Research methods in health: Investigating health and health services. Open University Press
13. Brabin BJ, Hakimi M, Pelletier D (2001) An analysis of anemia and pregnancy-related maternal mortality. J Nutr 131:604S-615S. https://doi.org/10.1093/jn/131.2.604S

14. Cairo A (2013) The functional art: An introduction to information graphics and visualization. New Riders
15. Carshon-Marsh R, Di Ruggiero E (2025) Improving the utilization of insecticide-treated nets for malaria prevention among pregnant women, lactating mothers and children in Sierra Leone: A commentary. Malaria Journal. https://doi.org/10.1186/s12936-025-05429-z
16. Cottrell G, Djènontin A, Soares C, et al (2025) Suboptimal distribution and utilization of antenatal care given bed nets undermine pregnant women's protection in Benin: A prospective field study. BMC Public Health. https://doi.org/10.1186/s12889-025-22212-6
17. Creswell JW (2014) Research design: Qualitative, quantitative, and mixed methods approaches, 4th ed. SAGE Publications
18. Desai M, ter Kuile FO, Nosten F, et al (2007) Epidemiology and burden of malaria in pregnancy. The Lancet Infectious Diseases 7:93–104. https://doi.org/10.1016/S1473-3099(07)70021-X
19. Donacho DO, Tura SK, Kefeni BT (2025) Determinants of long-lasting insecticide-treated bed net utilization among pregnant women in Gechi District, Buno Bedele Zone, South West Ethiopia: A community-based cross-sectional study. Environ Health Insights 19:. https://doi.org/10.1177/11786302251335135
20. Eisele TP, Larsen DA, Anglewicz PA, et al (2012) Malaria prevention in pregnancy, birthweight, and neonatal mortality: A meta-analysis of 32 national cross-sectional datasets in Africa. The Lancet Infectious Diseases 12:942–949. https://doi.org/10.1016/S1473-3099(12)70222-7
21. Eisele TP, Larsen DA, Walker N, et al (2009) Determining the effectiveness of the insecticide-treated bednet and indoor residual spraying campaigns in reducing malaria. PLOS ONE 4:e8145. https://doi.org/10.1371/journal.pone.0008145
22. Evergreen SDH (2017) Effective data visualization: The right chart for the right data. SAGE Publications
23. Fried M, Duffy PE (1996) Adherence of Plasmodium falciparum to chondroitin sulfate A in the human placenta. Science 272:1502–1504. https://doi.org/10.1126/science.272.5267.1502
24. Ghana Demographic and Health Survey (GDHS) (2008) Ghana demographic and health survey 2008. Ghana Statistical Service & Ghana Health Service
25. Ghana Health Service (GHS) (2009) Nkoranza South District health report 2009. District Health Directorate
26. Gikandi PW, Noor AM, Snow RW, et al (2008) Access and barriers to insecticide treated net use in urban and peri-urban areas of Kenya. Malaria Journal 7:227. https://doi.org/10.1186/1475-2875-7-227
27. Guyatt HL, Snow RW (2004) Impact of malaria during pregnancy on low birth weight in sub-Saharan Africa. Clinical Microbiology Reviews 17:760–769. https://doi.org/10.1128/CMR.17.4.760-769.2004
28. Healy K (2018) Data visualization: A practical introduction. Princeton University Press

29. Hunter JD (2007) Matplotlib: A 2D graphics environment. Computing in Science & Engineering 9:90–95. https://doi.org/10.1109/MCSE.2007.55
30. Ibeagha A, Ruiter RAC, Dukers-Muijrers NHTM, Hoebe CJ (2025) What factors contribute to the inconsistent use of insecticide treated bed nets in tropical Africa? A review of the literature on malaria prevention. Int J Psychol Health Sci 3:660. https://doi.org/10.38035/ijphs.v3i4.1738
31. Kabalu Tshiongo J, Zola Matuvanga T, Mitashi P, et al (2024) Prevention of malaria in pregnant women and its effects on maternal and child health, the case of Centre Hospitalier de Kingasani II in the Democratic Republic of the Congo. Tropical Medicine and Infectious Disease 9:92. https://doi.org/10.3390/tropicalmed9050092
32. Killeen GF, Smith TA, Ferguson HM, et al (2007) Preventing childhood malaria in Africa by protecting adults from mosquitoes with insecticide-treated nets. PLoS Medicine 4:e229. https://doi.org/10.1371/journal.pmed.0040229
33. Kokuhennadige V, Aitken E, Rogerson S (2026) Malaria in pregnancy at the frontline: A delicate balance. Current Opinion in Immunology. https://doi.org/10.1016/j.coi.2026.102726
34. Launiala A (2009) How much can a KAP survey tell us about people's knowledge, attitudes and practices? Some observations from medical anthropology research on malaria in pregnancy in Malawi. Anthropology Matters 11:1–13. https://doi.org/10.22582/am.v11i1.31
35. Lengeler C (2004) Insecticide-treated bed nets and curtains for preventing malaria. Cochrane Database of Systematic Reviews 2:CD000363. https://doi.org/10.1002/14651858.CD000363.pub2
36. Lim SS, Fullman N, Stokes A, et al (2011) Net benefits: A multicountry analysis of observational data examining associations between insecticide-treated mosquito nets and health outcomes. PLoS Medicine 8:e1001091. https://doi.org/10.1371/journal.pmed.1001091
37. Macintyre K, Keating J, Okello G, et al (2012) Examining the determinants of mosquito-avoidance practices in two Kenyan cities. Malaria Journal 11:9. https://doi.org/10.1186/1475-2875-11-9
38. Madukwe JC, Ahaneku EB, Onukafor OS, et al (2025) Do women attending antenatal clinics who use only intermittent preventive treatment (IPTp) have higher Plasmodium falciparum prevalence compared to those who used both IPTp and slept under insecticide-treated nets? Malaria Journal. https://doi.org/10.1186/s12936-025-05532-1
39. Minakawa N, Dida GO, Sonye GO, et al (2008) Unforeseen misuse of bed nets in fishing villages along Lake Victoria. Malaria Journal 7:165. https://doi.org/10.1186/1475-2875-7-165
40. Minwuyelet A, Yewhalaw D, Siferih M, Atenafu G (2025) Current update on malaria in pregnancy: A systematic review. Trop Dis Travel Med Vaccin 11:14. https://doi.org/10.1186/s40794-025-00248-1
41. Mwebesa E, Musinguzi B, Legason ID (2025) Pooled prevalence and factors associated with insecticide-treated net use among pregnant women in malaria high-burden countries in sub-Saharan Africa: A multilevel mixed-effects analysis. Trop Med Health 53:166. https://doi.org/10.1186/s41182-025-00855-w
42. National Malaria Control Programme (NMCP) (2017) Ghana malaria programme review 2017. Ghana Health Service

43. Nkoranza South District Health Management Team (DHMT) (2016) Annual district health report 2016. Nkoranza South District Health Directorate
44. Okoro LN, Zoakah AI, Bupwatda P, et al (2025) Cost-effectiveness of intermittent preventive therapy and insecticide-treated bed nets for malaria prevention among pregnant women in Jos North, Plateau State, Nigeria. Malaria Journal. https://doi.org/10.1186/s12936-025-05607-z
45. Okova D, Lukwa A, Edusei M (2025) Between- and within-socioeconomic groups temporal inequality in the uptake of malaria prevention strategies among pregnant women and under-five children in Ghana (2003-2022). Malar J 24:273. https://doi.org/10.1186/s12936-025-05512-5
46. Onwujekwe O, Chukwuka C, Eze S, et al (2006) Treating malaria: The role of drug shops and community health workers in Enugu State, south eastern Nigeria. Journal of Public Health 28:197–203
47. Opara MO, Ismail S, Mohd Nazan AI (2024) Prevention of malaria in pregnancy through health education intervention programs on insecticide-treated nets use: A systematic review. BMC Public Health. https://doi.org/10.1186/s12889-024-17650-7
48. Osei KM, Adiak AA, Han W (2026) Evaluating the effectiveness of insecticide-treated net use in preventing malaria among children under five: A quasi-experimental study. Tropical Medicine and Health. https://doi.org/10.1186/s41182-026-00932-8
49. Patton MQ (2002) Qualitative research and evaluation methods. SAGE Publications
50. Pettifor A, Taylor E, Nku D (2009) Free distribution of insecticide treated bed nets to pregnant women in Kinshasa: An effective way to achieve 80% use by women and their newborns. Trop Med Int Health 14:20–28. https://doi.org/10.1111/j.1365-3156.2008.02194.x
51. Pluess B, Tanser FC, Lengeler C, Sharp BL (2010) Indoor residual spraying for preventing malaria. Cochrane Database Syst Rev CD006657. https://doi.org/10.1002/14651858.CD006657.pub2
52. Pulford J, Hetzel MW, Bryant M, et al (2011) Reported reasons for not sleeping under a mosquito net in Papua New Guinea: Challenges for malaria prevention. Malaria Journal 10:183. https://doi.org/10.1186/1475-2875-10-183
53. Rahlf T (2017) Data visualisation with R: 111 examples. Springer
54. Salanti A, Dahlbäck M, Turner L (2004) Evidence for the involvement of VAR2CSA in pregnancy-associated malaria. J Exp Med 200:1197–1203. https://doi.org/10.1084/jem.20041579
55. Salifu Y, Nantomah B, Lasong J (2026) Knowledge of malaria and preventive practices among Fulani pregnant women in the Savannah Region of Northern Ghana. BMC Research Notes. https://doi.org/10.1186/s13104-026-07736-3
56. Steketee RW, Nahlen BL, Parise ME, Menendez C (2001) The burden of malaria in pregnancy in malaria-endemic areas. American Journal of Tropical Medicine and Hygiene 64:28–35. https://doi.org/10.4269/ajtmh.2001.64.28
57. Teddlie C, Yu F (2007) Mixed methods sampling: A typology with examples. Journal of Mixed Methods Research 1:77–100. https://doi.org/10.1177/1558689806292430

58. Teym A, Berihun G, Shiferaw MY (2025) Insecticide-treated bed net utilization and its determinants among pregnant women in Dembecha District, Northwest Ethiopia. Trop Med Health 53:129. https://doi.org/10.1186/s41182-025-00762-0
59. Vandamme J-P (2009) Rethinking KAP-studies. Sociologos 30:3–21
60. Waskom ML (2021) Seaborn: Statistical data visualization. Journal of Open Source Software 6:3021. https://doi.org/10.21105/joss.03021
61. Wickham H (2016) ggplot2: Elegant graphics for data analysis, 2nd ed. Springer
62. Wickham H (2016) Ggplot2: Elegant Graphics for Data Analysis. Springer, Cham, Switzerland
63. World Health Organization (WHO) (2015) World malaria report 2015. WHO Press
64. World Medical Association (2013) World Medical Association Declaration of Helsinki: Ethical principles for medical research involving human subjects. JAMA 310:2191–2194. https://doi.org/10.1001/jama.2013.281053
65. Zuuri CN, Gyamah DK, Buabeng L, Osei Mensah R (2025) Assessing the knowledge, attitudes and practices towards malaria prevention and determinants of antenatal care utilization among pregnant women in Sekyere South district, Ghana: A cross-sectional study. Malar J 24:371. https://doi.org/10.1186/s12936-025-05614-0

Figures




Figure 1

Distribution of key socio-demographic and knowledge variables (*N* = 200). Panel A: violin plot of respondent *age*with individual data points overlaid; dashed red line indicates median. Panel B: combined violin and box plot of *knowledge scores* (0–8) with knowledge-level bands (red = poor, gold = moderate, green = good). Panel C: horizontal bar chart of educational attainment. *Note. Panel B knowledge score*

distribution is right-censored at 8 (maximum possible score). The tight clustering in the 4–5 (moderate knowledge) range is consistent with the overall classification of 85% of respondents as demonstrating moderate knowledge. JHS = Junior High School; SHS = Senior High School.




Figure 2

KAP profile heatmap: percentage of respondents meeting each indicator (*N* = 200). Each row represents one KAP indicator. Green cells indicate high-coverage indicators (≥70%); yellow indicates moderate coverage (50–69%); red indicates low coverage (<50%). Domain labels on the left margin are colour-coded: navy = Knowledge, teal = Access, green = Practice. *Note. The heatmap reveals a bimodal performance profile: near-universal performance on awareness and access indicators, contrasted with critically low performance on knowledge depth (correct expiry period) and practice utilization (net correctly hung and in use). KAP = Knowledge, Attitudes, and Practices; ANC = Antenatal Care; LLITN = Long-Lasting Insecticidal Treated Net.*




Figure 3

Diverging bar chart: proportion of correct/positive vs. incorrect/absent responses across KAP indicators (*N* = 200). Teal (right) bars represent the percentage giving a correct or positive response; red (left) bars represent the complementary percentage. Indicators are ordered from highest to lowest positive response rate. *Note. The chart visually quantifies the magnitude of the gap between positive and negative responses for each indicator. The most critical divergences are seen for the correct expiry period (27.1% vs. 72.9%) and the net correctly hung and in use indicator (43.5% vs. 56.5%). ANC = Antenatal Care; KAP = Knowledge, Attitudes, and Practices.*




Figure 4

Lollipop chart: KAP indicators ranked by proportion of positive/correct responses (*N*= 200). Colours indicate coverage level: green ≥70% (high), gold 50–69% (moderate), red <50% (low). Dashed reference lines mark the 50% and 70% thresholds. *Note. The lollipop chart enables rapid visual identification of the highest- and lowest-performing indicators. The two red lollipops (correct expiry period and net correctly hung and in use) represent the primary targets for programmatic intervention. ANC = Antenatal Care; KAP = Knowledge, Attitudes, and Practices.*




Figure 5

Radar chart: domain-level KAP profile (*N* = 200). Each arm represents one KAP indicator; radial distance indicates the proportion of correct/positive responses (0–100%). The shaded area represents the overall KAP profile. Domain arcs indicate the three measurement domains: navy = Knowledge, teal = Access, green = Practice. *Note. The pronounced indentation of the radar profile at the 'Correct Expiry Period' arm (27.1%) and 'Correct Net Use' arm (43.5%) visually identifies the primary performance deficits. The full ANC-access arms reflect near-universal distributional achievement. KAP = Knowledge, Attitudes, and Practices; ANC = Antenatal Care.*




Figure 6

Cross-tabulation heatmaps: LLITN use status by occupational group (Panel A) and by knowledge level (Panel B). Cell values indicate row percentages and absolute counts (n). Colour intensity in Panel A reflects percentage magnitude (blue scale); Panel B uses a red–yellow–green diverging scale. *Note. Panel A reveals that farmers — the largest subgroup — exhibit one of the lower rates of correct net hanging and use (43%) and a notable non-use rate (34%), suggesting that occupational fatigue and time constraints may compound utilization barriers in this group. Panel B confirms a gradient between knowledge level and correct utilization, with poor-knowledge respondents demonstrating disproportionately high non-use (44%). LLITN = Long-Lasting Insecticidal Treated Net.*



Figure 7

Waffle charts: LLITN use confirmation status (Panel A) and overall knowledge level (Panel B). Each square represents approximately 1% of respondents (*N* = 200). *Note. The waffle visualization enables immediate visual comparison of the proportional distribution within each domain. Panel A highlights that less than half of respondents (43.5%) had their LLITN correctly hung and in use at the time of observation. Panel B illustrates the overwhelming dominance of the moderate knowledge category (85%). LLITN = Long-Lasting Insecticidal Treated Net.*




Figure 8

Bubble chart: multivariate relationship between knowledge, access, and practice indicators. Each bubble represents one KAP indicator; bubble size is proportional to estimated programmatic significance. Dashed lines indicate 50% thresholds for both axes. Colours indicate domain: navy = Knowledge, teal = Access, green = Practice. *Note. Indicators in the upper-right quadrant (high knowledge, high practice/access) represent programmatic strengths; indicators in the lower-left quadrant represent priority intervention targets. The ANC-access indicators cluster in the upper right, while the net use and expiry knowledge indicators fall in the lower left. ANC = Antenatal Care; KAP = Knowledge, Attitudes, and Practices.*