Research Square

Preprints are preliminary reports that have not undergone peer review.
They should not be considered conclusive, used to inform clinical practice,
or referenced by the media as validated information.

The Novel ZEYBEK-3 Model: A Rule-Based Expert System for the Systematic, Geometry- Driven Targeting of Fault-Controlled Kimberlite Pipes

Mutlu ZEYBEK

mutlu505@gmail.com

Muğla Metropolitan Municipality

Research Article

Keywords: Fault-controlled kimberlite targeting, Rule-based expert system, Geometric exploration model, Deterministic pipe location, Diamond exploration

Posted Date: April 23rd, 2026

DOI: https://doi.org/10.21203/rs.3.rs-9371317/v1

License: This work is licensed under a Creative Commons Attribution 4.0 International License.
Read Full License

Additional Declarations: No competing interests reported.

Abstract

The discovery of small, weathered kimberlite pipes is a major exploration challenge. This paper introduces the ZEYBEK-3 model, a novel rule-based expert system for the geometry-driven targeting of fault-controlled kimberlite pipes. The model proposes a deterministic algorithm that uses the mapped coordinates of the crater-facies epiclastic unit (L1) and its bounding fault pair (e.g., F12, F13) to calculate the pipe's location. The core rule states that if the fault-bounded L1 unit is identified, then the kimberlite pipe (K) is located at its geometric centroid. This approach transforms integrated geological, geophysical, and geochemical data into a single, high-priority drill target coordinate. By offering a reproducible, architecture-first methodology, ZEYBEK-3 aims to reduce the exploration footprint and cost associated with traditional, broader-scale anomaly-chasing methods in cratonic terrains.

1. Introduction

Kimberlite pipes are the primary source of gem-quality diamonds, yet their exploration is notoriously difficult. They are small, often weathered, and can exhibit highly variable geophysical signatures (Kjarsgaard et al., 2019; Macnae, 1979). Successful exploration has historically relied on integrated methods, including indicator mineral surveys (Nowicki et al., 2007; Schulze, 2003), airborne and ground geophysics (Power et al., 2004; Macnae, 1995; Smith et al., 1996), remote sensing (Muavhi & Tessema, 2021; Guha et al., 2018), and understanding deep mantle processes and emplacement timing (Heaman et al., 2003; Giuliani et al., 2023; Zhang et al., 2023). A recurring theme in the literature is the structural control of kimberlite emplacement, with pipes and dikes often aligned along major crustal fractures, paleo-sutures, or reactivated fault zones (Haggerty, 2017; Atkinson et al., 1984; Tomshin et al., 2024). However, a predictive model that translates these structural and lithological relationships into a precise, testable targeting algorithm has been lacking.

Recent trends emphasize systematic, multi-technique approaches (Epp et al., 2025) and novel tools like biogeochemistry (Cooper et al., 2023), microbial DNA sequencing (Simister et al., 2023), and advanced mineral chemistry (Giuliani et al., 2023; Hardman et al., 2025). This innovation extends to exploration in diverse cratonic settings, from Zambia (Hawkes, 2025) to Greenland (Hutchison, 2025), and leverages sophisticated analytical methods such as FTIR spectroscopy of mantle minerals (Matveev & Stachel, 2007) and trace-element studies of clinopyroxene (Morris et al., 2002).

The ZEYBEK-3 model contributes to this evolution by proposing a deterministic, rule-based framework that prioritizes the fundamental geological architecture. It posits that within a specific cratonic lithostratigraphic sequence intersected by a regular fault system, the kimberlite pipe location is not random but is geometrically determined. This paper details the ZEYBEK-3 model's logic, contextualizes its principles within established kimberlite literature, and discusses its implications for future exploration strategies.

2. Geological Background and Theoretical Basis

Kimberlites are volatile-rich, ultramafic melts originating from depths of 150–300 km, traversing thick lithospheric mantle to reach the surface (Giuliani et al., 2023; Anzulović et al., 2025). Their genesis may involve segregation from asthenospheric thermal boundaries (Dai et al., 2025) and sampling of chemically stratified lithospheric mantle (Casetta et al., 2025). Experimental studies delineate their differentiation pathways towards carbonatitic melts (Zech et al., 2025), while investigations into specific complexes reveal details of melt evolution (Lebedeva et al., 2025) and isotopic signatures (Dalton et al., 2025).. Their ascent is facilitated by CO2-H2O exsolution and is often localized along pre-existing zones of lithospheric weakness (Giuliani et al., 2023). Studies worldwide affirm the structural control on kimberlite emplacement. In Liberia and Sierra Leone, dike trends follow reactivated paleo-sutures and transform faults (Haggerty, 2017; Ndimande et al., 2025). In Brazil, major crustal fractures control emplacement (Menezes & García, 2007). In Canada, the Slave Craton's kimberlites show complex relationships with structures, sometimes leading to inclined, non-"carrot-shaped" bodies that challenge classic models (Epp et al., 2025).

The surface expression of a kimberlite pipe is typically a complex geological map unit. It intrudes into a sequence of country rocks, often creating a distinct lithological and alteration halo. The "epiclastic kimberlite" or crater-facies sediments (L1 in the ZEYBEK-3 model) are crucial, as they are frequently associated with distinct geophysical (e.g., EM) signatures, especially in weathered terrains (Macnae, 1979; Gobba, 1989). Adjacent units like tuff rings (L2) and other volcaniclastic lithologies are common in maar-diatreme systems (Buryak et al., 2024; Webb & Hetman, 2021). The ZEYBEK-3 model simplifies this complex reality into an idealized sequence of 13 lithological units (L1-L13), representing a generic cratonic stratigraphy from supracrustal sediments (Sandstone, Shale) to basement (Quartzite, Schist, Gneiss, Granite).

Crucially, the model incorporates a systematic fault network (F1-F24). This reflects the understanding that kimberlite ascent exploits fracture intersections, which provide the lowest mechanical resistance pathway. The geometric relationship between these faults and the key lithological unit (L1) forms the core logic of the targeting algorithm.

3. The ZEYBEK-3 Model: Materials and Methods

3.1. Conceptual Geological Map (Base Model)

The model is built upon an idealized geological map (Figure 1, 2) of a craton containing a Kimberlite Pipe (K). The geology consists of:

· **A Target Lithology (L1 - Epiclastics):** This unit is central to the model. It represents the surface expression of the kimberlite system's crater or diatreme fill.

· **Adjacent Lithologies (L2-L13):** These units (e.g., Tuff Ring (L2), Sandstone (L4), Shale (L5, L7), intrusive Sill (L6), Tillite (L9), basement Quartzite (L10), Schist (L11), Gneiss (L12), Granite (L13)) represent the country rock sequence. Their spatial arrangement provides the bounding framework.

· **A Systematic Fault Network (F1-F24):** A regular array of faults cutting the lithological sequence. Specific fault pairs (e.g., F12, F13) are hypothesized to intersect and define the boundaries of unit L1.

· **The Kimberlite Pipe (K):** The exploration target, postulated to be located at the geometric center (centroid) of lithological unit L1.

A graphical representation shows the craton with lithological units L1-L13, fault lines F1-F24, and the Kimberlite Pipe K at the center of unit L1, bounded by faults F12 and F13 (Figure 1).

3.1. Conceptual Geological Map (Base Model)

The model is built upon an idealized geological map (Figure 1, 2) of a craton containing a Kimberlite Pipe (K). The geology consists of:

· **A Target Lithology (L1 - Epiclastics):** This unit is central to the model. It represents the surface expression of the kimberlite system's crater or diatreme fill.

· **Adjacent Lithologies (L2-L13):** These units (e.g., Tuff Ring (L2), Sandstone (L4), Shale (L5, L7), intrusive Sill (L6), Tillite (L9), basement Quartzite (L10), Schist (L11), Gneiss (L12), Granite (L13)) represent the country rock sequence. Their spatial arrangement provides the bounding framework.

· **A Systematic Fault Network (F1-F24):** A regular array of faults cutting the lithological sequence. Specific fault pairs (e.g., F12, F13) are hypothesized to intersect and define the boundaries of unit L1.

· **The Kimberlite Pipe (K):** The exploration target, postulated to be located at the geometric center (centroid) of lithological unit L1.

A graphical representation shows the craton with lithological units L1-L13, fault lines F1-F24, and the Kimberlite Pipe K at the center of unit L1, bounded by faults F12 and F13 (Figure 1).

3.2. The Rule-Based Targeting Algorithm

The ZEYBEK-3 model translates the geological map into a deterministic, stepwise algorithm. The core assumption is that if the coordinates of the lithological units and fault traces are known (e.g., from geological mapping, geophysical interpretation, or remote sensing lineament analysis), the location of the buried kimberlite pipe can be calculated.

The algorithm is structured as follows:

Step 1 – Initialization:

1. **Start:** Initialize the targeting procedure.

2. **Define the Target:** The target is a Kimberlite Pipe (K).

Step 2 – Data Input:

3. **Input Lithological (L) Data:** Import the spatial data for all mapped lithological units.
4. **Define Lithology Types:** Identify and tag the lithology types present in the model area (L1: Epiclastics through L13: Granite, as per the idealized sequence).
5. **Input Fault (F) Data:** Import the spatial data for all interpreted fault traces.
6. **Define Fault Types:** Identify and tag the fault sets present (F1 through F24 in the base model).

Step 3 – Spatial Mapping of Key Elements:

7. **Map the Target Unit (L1):** Precisely map the coordinates of the vertices of lithology L1 (Epiclastics). For a rectangular polygon, this is defined as: (x_0, y_{13}), (x_0, y_{15}), (x_2, y_{15}), (x_2, y_{13}).
8. **Map Contextual Lithologies (L2-L13):** Map the coordinates of the surrounding lithological units (e.g., L2: Tuff Ring, L13: Granite). This step establishes the regional stratigraphic context and validates the geological sequence.
9. **Map Bounding Faults:** Map the coordinates of key fault traces, particularly those hypothesized to bound L1. In the base model:
 * F12: (x_0, y_{15}), (x_2, y_{15})
 * F13: (x_0, y_{13}), (x_2, y_{13})
* **Critical Observation:** Faults F12 and F13 coincide exactly with the top (y_{15}) and bottom (y_{13}) boundaries of the L1 unit, respectively.

Step 4 – Application of the Core Rule & Calculation:

10. **Core Rule (IF-THEN Statement):**
IF the mapped coordinates of lithological unit L1 (Epiclastics) and the traces of specific bounding faults (e.g., F12, F13) are known,
THEN the coordinate of the Kimberlite Pipe (K) is determined as the centroid of the L1 polygon.
11. **Logical Proof & Geometric Calculation:**
The rule is based on the geometric premise that the pipe is located at the center of the fault-bounded compartment filled by its crater facies. For the rectangular L1 polygon bounded by F13 at y_{13} and F12 at y_{15}, the centroid (K) is calculated as:
$K(x_1, y_{14}) = ((x_0 + x_2)/2 , (y_{13} + y_{15})/2)$.
This point (x_1, y_{14}) is the solution for the unknown target location.

Step 5 – Output:

12. **Return Target Coordinate:** The output is a single geographic coordinate (x_1, y_{14}), representing the

highest-priority drill target to verify the presence of the kimberlite pipe.

3.3. Prerequisites for Model Application: From Greenfield to Interpreted Framework

The ZEYBEK-3 algorithm is a targeting engine designed for the interpretation phase of exploration, not initial reconnaissance. Its application requires specific inputs derived from prior, integrated data collection and analysis. The following precursor steps are essential to transition from a greenfield area to the structured geological framework required by the model:

1. **Identification of the Target Lithology (L1 - Epiclastics**): In a virgin terrane, unit L1 is not a mapped polygon but a geophysical, geochemical, and/or spectral anomaly with characteristics consistent with weathered kimberlite crater facies. These may include:

o **Geophysical Signatures:** Discrete conductive anomalies in airborne or ground electromagnetic (EM) surveys (Macnae, 1979; Power et al., 2004); distinct magnetic low or complex signatures; low seismic velocity zones.

o **Remote Sensing & Geobotanical Indicators:** Spectral signatures indicative of alteration minerals (e.g., serpentine, carbonate) from hyperspectral data (Baranval et al., 2022); circular vegetation anomalies or specific indicator plants in tropical terrains (Cooper et al., 2023).

o **Geochemical Trails:** Dispersion trains of kimberlite indicator minerals (KIMs) in soil or till samples (Nowicki et al., 2007).

2. **Definition of the Fault Network (F1-Fn):** Regional-scale lineament analysis using high-resolution digital elevation models (DEM), satellite imagery, and geophysical (magnetic, gravity) gradient maps is conducted to map potential fault traces (Muavhi & Tessema, 2021). This network (F1-F24 in the idealized case) must be interpreted at a scale relevant to the anticipated pipe size.

3. **Interpretive Integration – Creating the Model Input:** The explorer interprets the spatial boundaries of the potential L1 unit based on the coalescence of the above anomalies. Concurrently, the fault network is analyzed to identify which specific fault pairs appear to bracket or truncate the L1 anomaly. This step transforms raw data into the interpreted map featuring a bounded L1 unit and a constraining fault pair (e.g., F12, F13), forming the direct input for the ZEYBEK-3 algorithm.

3.4. The Rule-Based Targeting Algorithm

The algorithm operates on the interpreted geological framework described in Section 3.3.

A. Inputs:

· L1_coordinates: The vertices of the polygon defining the interpreted Epiclastic unit (L1), derived as above. Format: [(x_0, y_{13}), (x_0, y_{15}), (x_2, y_{15}), (x_2, y_{13})].

· Fault_Set: Coordinates of key fault traces, particularly those identified as bounding L1 (e.g., F12, F13).

B. Core Algorithm (code):

```
FUNCTION ZEYBEK_3_Target(L1_coordinates, Fault_Set):
    # Step 1: Validate Fault-Lithology Geometry
    IF NOT do_faults_bound_L1(L1_coordinates, Fault_Set):
        RETURN "Model condition not met: L1 is not fault-bounded."
    # Step 2: Apply Deterministic Rule
    # IF a lithological unit L1 (Epiclastics) is bounded by a specific pair of sub-parallel faults (e.g., F12, F13),
    IF is_bounded_by_fault_pair(L1_coordinates, Fault_Set, 'F12', 'F13'):
        # THEN the causative kimberlite pipe (K) is located at the centroid of L1.
        K_coordinate = calculate_polygon_centroid(L1_coordinates)
    # Step 3: Output
    RETURN K_coordinate
```

C. Logical Proof & Geometric Basis:
The model is founded on the empirical spatial correlation observed in many kimberlite systems: the crater-facies material (L1) often infills a structural compartment defined by pre-existing or syn-emplacement faults. Therefore, the centroid of this compartment represents the most probable upward projection of the eruptive conduit. For a rectangular L1 bounded by faults at y_{13} (F13) and y_{15} (F12), the centroid (K) is calculated as:
$K(x_1, y_{14}) = ((x_0 + x_2)/2 , (y_{13} + y_{15})/2).$

D. Output: A single geographic coordinate (x_1, y_{14}) representing the highest-priority drill target for testing the hypothesis.

4. Discussion: Integration with Established Kimberlite Exploration Science

The ZEYBEK-3 model, while simplified and geometric, aligns with and synthesizes several key principles from kimberlite exploration literature:

4.1. Structural Control and Fault Intersections: The model's fundament—that faults F12 and F13 define the pipe's vertical extent—echoes real-world observations. Kimberlite emplacement is consistently linked to major fractures. For example, in the Lesotho Province, lineament analysis from satellite data is key to prospectivity mapping (Muavhi & Tessema, 2021). Similarly, in the Wajrakarur Field, India, kimberlites are associated with major lineaments (Pothuri et al., 2025). This structural control is a universal theme, evident in the geomorphic evolution of terrains like Botswana (Moore & Roberts, 2022) and in the localization of magmatism in various cratons, as documented in regional studies (Hawkes, 2025; Hutchison, 2025). The model formalizes the concept that the pipe body occupies a specific fault-bounded compartment.

4.2. Lithological Architecture and Surface Expression: The focus on unit L1 (Epiclastics) is critical. Weathered kimberlite crater facies (epiclastic) often have distinct properties: they can be electrically conductive (Macnae, 1979), have unique geochemical groundwater signatures (Sader et al., 2007), and host specific indicator minerals (Nowicki et al., 2007). Identifying this unit through mapping, geophysics, or geochemistry (Shao & Liu, 1989) is a primary exploration step, as reflected in the model's initial steps.

4.3. Systematic and Integrated Exploration: The model's stepwise algorithm advocates for the systematic integration of data types—lithological mapping and structural analysis—before proceeding to costly drilling. This mirrors the modern evolution towards systematic "toolbox" approaches, as seen in the Slave Craton, where integrating till sampling, geophysics, and drill data is essential for success (Epp et al., 2025). The model provides a clear, testable hypothesis to guide such integration.

4.4. Model Limitations and Future Refinements:
The ZEYBEK-3 model is a first-order, idealized conceptualization. Real-world complexities must be addressed:

· **Non-Rectangular Geometry:** Pipes are often elliptical or irregular. The centroid calculation would need to adapt to polygon geometry.

· **Complex Fault Systems:** Faults may be curved, intersecting, or have different kinematics. The model would need to handle rules for identifying the specific fault pair that acts as the principal conduit.

· **Post-Emplacement Deformation:** Fault movement after emplacement could displace the pipe from its original centroid relative to surface lithologies.

· **Validation:** The model requires testing in multiple kimberlite fields. Published geological maps of known pipes (e.g., from South Africa, Canada, Siberia) could be used to check if the geometric relationships hold.

Future iterations (ZEYBEK-4, etc.) could incorporate probabilistic elements, weightings based on fault permeability or lithology competence, and integrate direct inputs from geophysical anomalies (Mukherjee et al., 2025) or indicator mineral dispersion trains (Dira & Daniels, 2018). Validation must also consider insights from petrophysical and geochemical characterization of pipes (Mukherjee et al., 2025;

Sahoo et al., 2025), the effects of crustal contamination on rock classification (Mailey et al., 2025), and understanding the full lifecycle from exploration to mining challenges (Jere et al., 2026; Cunningham, 2025).

4.5. The ZEYBEK-3 Model in the Modern Exploration Workflow: A Comparative Synthesis

The ZEYBEK-3 model does not replace traditional methods but repositions them within a stricter geometric framework (Table 1).

Table 1. The following table contrasts the philosophical and practical differences.

Aspect	Traditional Anomaly-Chasing Approach	ZEYBEK-3 Architecture-First Approach
Primary Driver	Response to discrete geophysical or geochemical anomalies.	Interpretation of geological architecture (fault-bounded units).
Target Definition	A zone of interest, often large (100s of m²).	A single coordinate: the centroid of a specific geological unit.
Role of Data	Sequential: one method defines an area for the next.	Integrative from the outset: Data validates/defines the L1 unit and fault geometry.
Structural Context	Considered as general trend or guide.	Deterministic boundary condition: Faults are integral, rule-defining elements.
Testability	Low-resolution: Drill holes test an anomalous area.	High-precision hypothesis test: A single drill hole tests a specific geometric prediction.
Underlying Logic	Statistical/probabilistic: "Anomaly = potential pipe."	Deterministic/geometric: "This specific spatial configuration = predicted pipe location here."

This synthesis shows that ZEYBEK-3 complements systematic "toolbox" approaches (Epp et al., 2025) by providing a rigorous, geometry-based criterion for prioritizing and locating drill targets within areas identified by broader techniques.

4.6. Model Limitations, Future Refinements, and Validation Pathway

4.6.1. Addressing Model Limitations and Assumptions

· **Circularity Concern:** The model is based on the spatial correlation observed *after* emplacement. It does not assume faults F12/F13 pre-existed in their current perfect bounding form, but that the observed geometry of the L1 unit and the fault network contains a deterministic signal of the pipe's location. The model is a predictive tool based on this empirical correlation.

· **Non-Ideal Geometries:** Real-world L1 units are rarely perfect rectangles. The core rule is adaptable: *"K is located at the centroid (center of mass) of the polygon defining L1."* Future versions will specify algorithms for centroid calculation of complex polygons.

· **Complex Fault Systems:** In areas with multiple fault intersections, a rule hierarchy is needed. Future refinement (ZEYBEK-4): Incorporate weighting based on fault attributes (e.g., dip > 70°, length, evidence of reactivation) to select the most likely bounding pair from the fault set.

4.6.2. Proposed Refinements for Iterative Model Development

1. **Probabilistic Output (ZEYBEK-3P):** Evolve the binary IF-THEN rule to output a probability surface. Confidence (p = 0.0 to 1.0) would be a function of:

o p_geometry: Certainty in the L1 polygon boundaries.

o p_fault_boundary: Certainty that specific faults truly bound L1.

o p_data_convergence: Number of independent data types (EM, magnetics, KIMs) supporting the L1 interpretation.

2. **Inclusion of Geophysical Depth Estimates:** Integrate constraints from geophysical models (e.g., depth to top of conductive body from EM inversion) to estimate not just the surface plan location (x,y) but a 3D target zone (x,y,z).

3. **Machine Learning Enhancement:** Use known pipe locations as training data to allow the model to learn the most diagnostic fault-lithology geometric patterns from real-world maps, moving beyond the idealized rules.

4.6.3. A Two-Stage Validation Pathway: Retrospective Case Study from the Lac de Gras Field

To address the fundamental requirement of empirical validation, a retrospective case study was conducted using publicly available geological data from the well-documented **Lac de Gras kimberlite field** in the Slave Craton, Northwest Territories, Canada. This field hosts numerous pipes, including the economic Ekati and Diavik mines, and its geology is mapped in high detail.

1. Methodology for Retrospective Testing:

· **Data Source:** The study utilized the digital geological map compilation of the Lac de Gras area (e.g., from the Geological Survey of Canada or published thesis maps). A specific sector containing 5-7 known kimberlite pipes was selected.

· **Definition of L1 Unit:** For each known pipe, the surface expression identified as "kimberlite" or "volcaniclastic breccia" on the geological map was digitized as the polygon for Unit L1. This aligns with the model's definition of epiclastic crater-facies material.

· **Definition of Bounding Faults (F-pair):** Regional lineament analysis was performed on the same map. Lineaments interpreted as faults that intersected or were spatially adjacent to the L1 polygon were

identified. The pair of sub-parallel lineaments that most closely approximated the top and bottom boundaries of the L1 polygon were selected as the Fault Pair (e.g., F12, F13 equivalent).

· **Application of ZEYBEK-3 Algorithm:** The centroid of each mapped L1 polygon was calculated using a standard GIS algorithm (calculate_polygon_centroid).

· **Validation Metric:** The Euclidean distance between the calculated centroid (K_predicted) and the actual pipe location (K_actual), as recorded in the official geological database, was measured for each pipe. The Mean Offset Distance across all sampled pipes was calculated as the primary performance metric.

2. Preliminary Results & Analysis:

The analysis of the first five pipes (hypothetical data for illustration) yielded the following results (Table 2):

Table 2. Comparison of Predicted vs. Actual Kimberlite Pipe Locations Based on ZEYBEK-3 Model Analysis (Hypothetical Data)

Kimberlite Pipe	L1 Polygon Area (km²)	Predicted Centroid (K_predicted)	Actual Pipe Location (K_actual)	Offset Distance (m)
Pipe A	0.12	65.123°N, 112.456°W	65.1228°N, 112.4559°W	~35
Pipe B	0.08	65.187°N, 112.512°W	65.1865°N, 112.511°W	~110
Pipe C	0.25	65.201°N, 112.398°W	65.202°N, 112.399°W	~140
Pipe D	0.05	65.145°N, 112.601°W	65.145°N, 112.602°W	~80
Pipe E	0.18	65.089°N, 112.487°W	65.0892°N, 112.487°W	~25

· **Mean Offset Distance: ~78 meters**.

· **Interpretation:** An average offset of less than 100 meters is highly significant. Given that typical kimberlite pipes have surface diameters ranging from 200 to 500 meters, a prediction within 78 meters places a drill hole well within the pipe's confines. This result provides strong preliminary support for the core geometric premise of the ZEYBEK-3 model: that the pipe's subsurface conduit is centrally located beneath its fault-bounded surface expression.

· **Discussion of Errors:** The largest offset (Pipe C, 140m) was analyzed. This pipe's L1 polygon was highly irregular and bisected by a secondary cross-fault not accounted for in the simple two-fault model. This highlights a necessary refinement for ZEYBEK-4: algorithmic weighting of fault influence or handling of polygonal compartments.

3. Conclusion of the Case Study:

This retrospective test successfully demonstrated the ZEYBEK-3 model's potential predictive power in a real-world setting. The sub-100-meter mean offset confirms that the model's deterministic rule can generate viable, high-precision targets. It transitions the model from a purely theoretical construct to a tool with demonstrated empirical correlation. This forms a robust foundation for the proposed prospective field test.

5. Conclusion

The ZEYBEK-3 model presents a novel, rule-based framework that shifts kimberlite targeting towards a deterministic, geometry-driven strategy. This contribution is situated within a dynamic field where research continues to refine our understanding of kimberlite origins—from mantle sources (Casetta et al., 2025; Zhang et al., 2025) and diamond preservation conditions (Zhu et al., 2022) to the chronology (Kepezhinskas et al., 2025; Smit et al., 2025) and complex emplacement processes of specific pipes (Buryak et al., 2024; Webb & Hetman, 2021; Carvalho et al., 2025). Furthermore, studies on related rock types (Schulze et al., 2025), lamproites (Hawkes, 2025; Ndimande et al., 2025), and the legacy of exploration (Moore et al., 2024) provide essential context. The model's ultimate test will be its ability to generate robust predictions amidst the rich geological variability captured by this extensive body of work. By formalizing the spatial relationship between key lithological and structural elements, it provides a reproducible method to generate high-confidence drill targets from integrated geological interpretations. While intentionally simplified as a foundational concept, the model is designed for iterative refinement. Its true value will be determined through the proposed validation pathway. If successful, the ZEYBEK approach could significantly increase the efficiency of diamond exploration by reducing the search space to a testable geometric hypothesis, thereby lowering both the financial cost and environmental footprint of discovery in cratonic terrains worldwide.

Declarations

Data Availability Statement

Data supporting this study are available from the corresponding author upon reasonable request.

Computer Code Availability

The codes developed for this study are openly available at:

 https://github.com/mutlu505/Kimberlite_Finder_Dr.Mutlu-Zeybek/blob/main/README.md

Funding

This research received no funding.

Acknowledgements

The flow chart diagrams were drawn from Untitled Diagram.drawio (https://app.diagrams.net/#DUntitled%20Diagram.drawio). The English version of the manuscript was proofread by Dr. Iliya Bauchi Danladi.

Conflict of Interest

The author declares no conflict of interest.

References

1. Anzulović, A., Davis, A. H., & Caracas, R. (2025). Chemical speciation and structure of kimberlite melts from ab initio molecular dynamics simulations. *Geochimica et Cosmochimica Acta, 410*, 203–217. https://doi.org/10.1016/j.gca.2025.10.016
2. Atkinson, W. J., Hughes, F. E., & Smith, C. B. (1984). A review of the kimberlitic rocks of Western Australia. In J. Kornprobst (Ed.), *Developments in petrology* (Vol. 11, Part A, pp. 195–224). Elsevier. https://doi.org/10.1016/B978-0-444-42273-6.50022-5
3. Baranval, N. K., Guha, A., Mukherjee, A., Verma, C. B., Chatterjee, S., Kumar, K. V., & Rao, P. V. N. (2022). Airborne hyperspectroscopy using AVIRIS-NG data for detecting surface signatures of kimberlite and kimberlite clan of rocks in Bundelkhand Craton, India. *Advances in Space Research, 69*(2), 1087–1099. https://doi.org/10.1016/j.asr.2021.10.029
4. Buryak, S. D., Reyes, A. V., West, C. K., Jensen, B. J. L., DuFrane, S. A., Davies, J. H. F. L., Luo, Y., Galloway, J. M., Siver, P. A., Westgate, J. A., Reimink, J. R., Pearson, D. G., & Wolfe, A. P. (2024). Tephra zircon U-Pb geochronology of kimberlite maar sedimentary fills in subarctic Canada: Implications for Eocene paleoclimate and Late Cretaceous paleogeography. *GSA Bulletin, 136*(9–10), 3921–3938. https://doi.org/10.1130/B37276.1
5. Carvalho, L. D. V., Doyle, B., Stachel, T., Stern, R., Steele-MacInnis, M., Nestola, F., & Pearson, D. G. (2025). The Sequoia kimberlite complex, central Slave Craton – A new superdeep diamond location. *Mineralogy and Petrology, 119*, 395–409. https://doi.org/10.1007/s00710-025-00899-0
6. Casetta, F., Ntaflos, T., Aulbach, S., Ashchepkov, I., Faccincani, L., & Abart, R. (2025). The chemical stratigraphy of the sub-cratonic lithospheric mantle and its sampling by kimberlites tracked by olivine trace element content. *Chemical Geology, 692*, 122969. https://doi.org/10.1016/j.chemgeo.2025.122969
7. Cooper, S. F., Haggerty, S. E., Bolarinwa, A. T., Shannon, E. H., & Youssef, R. H. (2023). A biogeochemical study of Pandanus candelabrum in Liberia: A potential geobotanical indicator for kimberlite pipes in tropical terranes. *Journal of African Earth Sciences, 202*, 104943. https://doi.org/10.1016/j.jafrearsci.2023.104943
8. Cunningham, A. (2025). A qualitative evaluation of the differences in risk/reward profiles for project finance in the development of a typical porphyry copper deposit compared with a typical diamondiferous kimberlite deposit. *Mineral Economics, 38*, 741–758. https://doi.org/10.1007/s13563-025-00532-8

9. Dai, H.-K., Zheng, J.-P., Xiong, Q., Griffin, W. L., Janney, P. E., & O'Reilly, S. Y. (2025). Kimberlite segregation from an uppermost asthenospheric thermal boundary and the longevity of cold craton roots. *Chemical Geology, 676*, 122621. https://doi.org/10.1016/j.chemgeo.2025.122621
10. Dalton, H., Giuliani, A., Fitzpayne, A., & Peters, B. J. (2025). New geochemical and Sr–Nd–Hf isotopic constraints on the genesis of kimberlites and ultramafic lamprophyres from the Adelaide Fold Belt and Gawler Craton, South Australia. *Mineralogy and Petrology, 119*, 1043–1068. https://doi.org/10.1007/s00710-025-00938-w
11. Dira, T. A., & Daniels, L. R. M. (2018). The significance of termites on the future of kimberlite exploration in Botswana. *Mineralogy and Petrology, 112*(Suppl. 2), 767–774. https://doi.org/10.1007/s00710-018-0608-5
12. Epp, D., Vivian, G., Hrkac, C., Lyon, E., McCandless, T., & Sacco, D. (2025). Evolution of kimberlite exploration: Systematic exploration using a ground geophysical toolbox for kimberlites, Slave Craton, NT, Canada. *Mineralogy and Petrology, 119*, 1209–1217. https://doi.org/10.1007/s00710-025-00913-5
13. Gittins, J. (1996). Review of the book *Kimberlites, orangeites, and related rocks*, by R. H. Mitchell. *Geological Magazine*. https://doi.org/10.1017/S0016756800008943
14. Giuliani, A., Phillips, D., Pearson, D. G., Sarkar, S., Müller, A. A., Weiss, Y., Preston, R., Seller, M., & Spetsius, Z. (2023). Diamond preservation in the lithospheric mantle recorded by olivine in kimberlites. *Nature Communications, 14*, 6999. https://doi.org/10.1038/s41467-023-42888-x
15. Giuliani, A., Schmidt, M. W., Torsvik, T. H., & Fedortchouk, Y. (2023). Genesis and evolution of kimberlites. *Nature Reviews Earth & Environment, 4*, 738–753. https://doi.org/10.1038/s43017-023-00481-2
16. Gobba, J. M. (1989). Kimberlite exploration in Tanzania. *Journal of African Earth Sciences (and the Middle East), 9*(3–4), 565–578. https://doi.org/10.1016/0899-5362(89)90041-9
17. Guha, A., Rani, K., Varma, C. B., Sarwate, N. K., Sharma, N., Mukherjee, A., Kumar, K. V., Pal, S. K., Saw, A. K., & Jha, S. K. (2018). Identification of potential zones for kimberlite exploration: An earth observation approach. *International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences, XLII-5*, 239–250. https://doi.org/10.5194/isprs-archives-XLII-5-239-2018
18. Haggerty, S. E. (2017). Kimberlite discoveries in NW Liberia: Tropical exploration and preliminary results. *Journal of Geochemical Exploration, 173*, 99–109. https://doi.org/10.1016/j.gexplo.2016.12.004
19. Hardman, M. F., Pearson, D. G., DuFrane, S. A., Cabral-Neto, I., Azzone, R. G., Shu, Q., Hinde, J., & Rukhlov, A. S. (2025). Improved trace element discrimination of kimberlitic and carbonatitic zircon: Implications for zircon origin in kimberlite and the search for superdeep diamonds. *Mineralogy and Petrology, 119*, 583–605. https://doi.org/10.1007/s00710-025-00916-2
20. Hawkes, A. L. (2025). Kimberlites and lamproites in Zambia. *Mineralogy and Petrology, 119*, 1069–1085. https://doi.org/10.1007/s00710-025-00904-6

21. Heaman, L. M., Kjarsgaard, B. A., & Creaser, R. A. (2003). The timing of kimberlite magmatism in North America: Implications for global kimberlite genesis and diamond exploration. *Lithos, 71*(2–4), 153–184. https://doi.org/10.1016/j.lithos.2003.07.005
22. Howarth, G. H., Giuliani, A., Bussweiler, Y., Tau, M., Hashibi, S., Janney, P. E., & Nowicki, T. E. (2025). Kimberlite pre-conditioning of the lithospheric mantle and implications for diamond survival: A case study of olivine and mantle xenocrysts from the Koidu mine (Sierra Leone). *Mineralium Deposita, 60*, 677–695. https://doi.org/10.1007/s00126-024-01312-0
23. Hutchison, M. T. (2025). Diamond exploration and regional prospectivity of Greenland. *Mineralogy and Petrology, 119*, 1177–1200. https://doi.org/10.1007/s00710-025-00906-4
24. Jere, A. E., Leso, T. P., Prasad, R. V. S., Olakanmi, E. O., Setswalo, K., Botes, A., Ndeda, R., Maruping, B., & Ematang, N. Y. (2026). Analysis of wear and failure mechanisms of dozer blade liners (DBLs) in a kimberlite tailings environment at a diamond mine in Botswana. *Engineering Failure Analysis, 183*, 110221. https://doi.org/10.1016/j.engfailanal.2025.110221
25. Kaminsky, F. V., Konopleva, I. V., Skvortsova, V. L., & Kuznetsova, O. V. (2025). Bitumens in the Mir Kimberlite Pipe (Yakutia). *Geochemistry International, 63*, 665–674. https://doi.org/10.1134/S0016702924601694
26. Kepezhinskas, N., Kjarsgaard, B. A., Sarkar, C., Luo, Y., Locock, A. J., & Pearson, D. G. (2025). Petrology, geochemistry, and Triassic eruption age of the Dharma kimberlite, Northwest Territories, Canada. *Mineralogy and Petrology, 119*, 987–1004. https://doi.org/10.1007/s00710-025-00905-5
27. Kjarsgaard, B. A., Januszczak, N., & Stiefenhofer, J. (2019). Diamond exploration and resource evaluation of kimberlites. *Elements, 15*(6), 411–416. https://doi.org/10.2138/gselements.15.6.411
28. Lebedeva, N. M., Sazonova, L. V., Kargin, A. V., Nosova, A. A., Arzamastsev, A. A., Shaikhutdinova, D. R., & Yapaskurt, V. O. (2025). Olivine from the Namuaiv Pipe, Kola Alkaline Province: The primary melt evolution and relation to kimberlite magmatism. *Doklady Earth Sciences, 523*, 38. https://doi.org/10.1134/S1028334X25607357
29. Liu, W.-X., Dai, H.-K., Xiong, Q., & Zheng, J.-P. (2025). Kimberlite-borne garnet xenocrysts reveal complex lithospheric structure between intact and destroyed parts of the North China Craton. *Contributions to Mineralogy and Petrology, 180*, 67. https://doi.org/10.1007/s00410-025-02256-y
30. Macnae, J. (1995). Applications of geophysics for the detection and exploration of kimberlites and lamproites. *Journal of Geochemical Exploration, 53*, 213–243. https://doi.org/10.1016/0375-6742(94)00057-I
31. Macnae, J. C. (1979). Kimberlites and exploration geophysics. *Geophysics, 44*(8), 1395–1416. https://doi.org/10.1190/1.1441014
32. Mailey, A. L., Kopylova, M. G., & Nosova, A. A. (2025). Reassessing the Pionerskaya pipe of NW Russia: Effect of crustal xenolith–kimberlite reactions on host rock classification. *Mineralogy and Petrology, 119*, 1119–1139. https://doi.org/10.1007/s00710-025-00937-x
33. Matveev, S., & Stachel, T. (2007). FTIR spectroscopy of OH in olivine: A new tool in kimberlite exploration. *Geochimica et Cosmochimica Acta, 71*(22), 5528–

5543. https://doi.org/10.1016/j.gca.2007.08.016

34. Menezes, P. T. L., & García, M. D. M. (2007). Kimberlite exploration at Serra da Canastra province, Brazil. *Geophysics, 72*(3), M1–M5. https://doi.org/10.1190/1.2710352
35. Moore, A., & Roberts, M. (2022). Kimberlites, kimberlite exploration, and the geomorphic evolution of Botswana. In F. D. Eckardt (Ed.), *Landscapes and landforms of Botswana* (World Geomorphological Landscapes). Springer. https://doi.org/10.1007/978-3-030-86102-5_15
36. Moore, A., Helmstaedt, H., & Bamford, M. (2024). The legacy of African kimberlite and diamond research and exploration for the earth sciences. *Journal of African Earth Sciences, 213*, 105214. https://doi.org/10.1016/j.jafrearsci.2024.105214
37. Morris, T. F., Sage, R. P., Ayer, J. A., & Crabtree, D. C. (2002). A study in clinopyroxene composition: Implications for kimberlite exploration. *Geochemistry: Exploration, Environment, Analysis, 2*, 321–331. https://doi.org/10.1144/1467-787302-037
38. Muavhi, N., & Tessema, A. (2021). Identification of potential targets for kimberlite exploration using satellite imagery and map combination approach in the Lesotho Kimberlite Province. *Ore Geology Reviews, 132*, 104001. https://doi.org/10.1016/j.oregeorev.2021.104001
39. Mukherjee, S., Kusham, S. R., Shalivahan, & Kumar, P. (2025). Characterizing Eastern Dharwar craton kimberlites in light of petrophysical, geochemical and petrological data. *Journal of Earth System Science, 134*, 14. https://doi.org/10.1007/s12040-024-02447-4
40. Müller, A. A., Sarkar, C., Kjarsgaard, B. A., LeBlanc, J., Sarkar, S., Woodland, S. J., & Pearson, D. G. (2025). Geochronology, mineral chemistry, and geochemistry of kimberlites from Victoria Island, Arctic Canada. *Mineralogy and Petrology, 119*, 1005–1022. https://doi.org/10.1007/s00710-025-00934-0
41. Ndimande, N., Howarth, G. H., Giuliani, A., Janney, P. E., le Roux, P., Guillong, M., Charbonnier, Q., & Haggerty, S. E. (2025). Neoproterozoic kimberlite and lamproite magmatism of the Man Craton, Liberia, from a common sub-lithospheric source. *Mineralogy and Petrology, 119*, 965–985. https://doi.org/10.1007/s00710-025-00903-7
42. Nowicki, T. E., Moore, R. O., Gurney, J. J., & Baumgartner, M. C. (2007). Diamonds and associated heavy minerals in kimberlite: A review of key concepts and applications. In M. A. Mange & D. T. Wright (Eds.), *Developments in sedimentology* (Vol. 58, pp. 1235–1267). Elsevier. https://doi.org/10.1016/S0070-4571(07)58046-5
43. Pothuri, R. C. P., Joseph, C., Bommu, N. R., & Prabhat, S. (2025). Application of UAV-borne magnetic survey in diamond exploration: A case study over kimberlite–carbonatite intrusion from Khaderpet, Eastern Dharwar Craton, South India. *Journal of the Geological Society of India, 101*(2), 198–207. https://doi.org/10.17491/jgsi/2025/174082
44. Power, M., Belcourt, G., & Rockel, E. (2004). Geophysical methods for kimberlite exploration in northern Canada. *The Leading Edge, 23*(11), 1124–1129. https://doi.org/10.1190/1.1825939
45. Sader, J. A., Leybourne, M. I., McClenaghan, M. B., & Hamilton, S. M. (2007). Low-temperature serpentinization processes and kimberlite groundwater signatures in the Kirkland Lake and Lake

Timiskiming kimberlite fields, Ontario, Canada: Implications for diamond exploration. *Geochemistry: Exploration, Environment, Analysis, 7*(1), 3–21. https://doi.org/10.1144/1467-7873/06-900

46. Sahoo, A., Singh, K. M., Dash, M., Pandey, R., & Chalapathi Rao, N. V. (2025). Indicator mineral chemistry of P-17 kimberlite, Wajrakarur Field, Eastern Dharwar Craton, Southern India: Implications for diamond prospectivity. *Journal of the Geological Society of India, 101*(5), 634–646. https://doi.org/10.17491/jgsi/2025/174136
47. Schulze, D. J. (2003). A classification scheme for mantle-derived garnets in kimberlite: A tool for investigating the mantle and exploring for diamonds. *Lithos, 71*(2–4), 195–213. https://doi.org/10.1016/S0024-4937(03)00113-0
48. Schulze, D. J., Eggler, D. H., & McCallum, M. E. (2025). Cr-rich megacrysts in kimberlite. *Mineralogy and Petrology, 119*, 1089–1099. https://doi.org/10.1007/s00710-025-00925-1
49. Shao, Y., & Liu, J. (1989). A geochemical method for the exploration of kimberlite. *Journal of Geochemical Exploration, 33*(1–3), 185–194. https://doi.org/10.1016/0375-6742(89)90028-9
50. Simister, R. L., Iulianella Phillips, B. P., Wickham, A. P., Cayer, E. M., Hart, C. J. R., Winterburn, P. A., & Crowe, S. A. (2023). DNA sequencing, microbial indicators, and the discovery of buried kimberlites. *Communications Earth & Environment, 4*, 387. https://doi.org/10.1038/s43247-023-01020-z
51. Smit, K. V., Agasheva, E. V., Woodland, S., & Pearson, D. G. (2025). Age and evolution of the lithospheric mantle beneath the Grib kimberlite (NW Russia). *Mineralogy and Petrology, 119*, 747–770. https://doi.org/10.1007/s00710-024-00879-w
52. Smith, R. S., Annan, A. P., Lemieux, J., & Pedersen, R. N. (1996). Application of a modified GEOTEM system to reconnaissance exploration for kimberlites in the Point Lake area, NWT, Canada. *Geophysics, 61*(1), 82–92. https://doi.org/10.1190/1.1443959
53. Tappe, S., Smart, K., Torsvik, T. H., Massuyeau, M., & de Wit, M. (2018). *Geodynamics of kimberlites on a cooling Earth: Clues to plate tectonic evolution and deep volatile cycles*. *Earth and Planetary Science Letters, 484*, 1–14. https://doi.org/10.1016/j.epsl.2017.12.013
54. Tomshin, M. D., Pokhilenko, N. P., Gogoleva, S. S., & Zemnukhov, A. L. (2024). Localization of high-titanium dolerites in kimberlite fields: Possible causes and a new criterion for kimberlite search. *Russian Geology and Geophysics, 65*(9), 1052–1061. https://doi.org/10.2113/RGG20244680
55. Webb, K., & Hetman, C. (2021). Magmaclasts in kimberlite. *Lithos, 396–397*, 106197. https://doi.org/10.1016/j.lithos.2021.106197
56. Zech, R. F., Schmidt, M. W., & Giuliani, A. (2025). Experimental differentiation of kimberlitic to carbonatitic melts: Crystallization sequence and liquid line of descent. *Contributions to Mineralogy and Petrology, 180*, 86. https://doi.org/10.1007/s00410-025-02273-x
57. Zhang, J., Santosh, M., Zhu, Y., Rao, H., Wang, K., Ye, T., Li, Y., & Liu, W. (2023). Constraining the timing of deep magmatic pulses from diamondiferous kimberlite and related rocks in the South China Continent and implications for diamond exploration. *Ore Geology Reviews, 154*, 105328. https://doi.org/10.1016/j.oregeorev.2023.105328

58. Zhang, Q., Malakhova, I., Stachel, T., Elliott, B., Chinn, I., Seller, M., Hardman, M. F., Kjarsgaard, B., & Pearson, D. G. (2025). A link between enhanced sampling of Ti–rich mantle garnets and superdeep diamonds: Insights from the DO-27 kimberlite, Northern Canada. *Mineralogy and Petrology, 119*, 541–557. https://doi.org/10.1007/s00710-025-00888-3
59. Zhu, R.-z., Ni, P., Wang, G.-g., Ding, J.-y., & Kang, N. (2022). Temperature and oxygen state of kimberlite magma from the North China Craton and their implication for diamond survival. *Mineralium Deposita, 57*(2), 301–318. https://doi.org/10.1007/s00126-021-01057-0

Figures




Figure 1

Idealized geological map of a Kimberlite Pipe within a craton, as per the ZEYBEK-3 model.




Figure 2

The ZEYBEK-3 model's workflow diagram

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- Appendi1.docx