

Parasitological efficacy of sulfadoxine-pyrimethamine and amodiaquine for Seasonal Malaria Chemoprevention (SMC) in Africa

Chayanin Sararat

`chayanin@tropmedres.ac`

Mahidol-Oxford Tropical Medicine Research Unit <https://orcid.org/0000-0002-5486-6024>

Craig Bonnington

Centre for Tropical Medicine and Global Health

Paul Sondo

Infectious Diseases Data Observatory

Jane Achan

Malaria Consortium

Chukwudi Nnaji

Malaria Consortium

Anna Burkhard

Malaria Consortium

Gauthier Tougri

Programme National de Lutte contre le Paludisme

Adama Traore

Malaria Consortium

Aimé Yaro

Malaria Consortium

Anthony Nuwa

Malaria Consortium

Richard Kajubi

Malaria Consortium

Kanokon Suwanansin

Mahidol-Oxford Tropical Medicine Research Unit,

Stephane Proux

Mahidol University

Urairat Koesukwiwat

Mahidol-Oxford Tropical Medicine Research Unit

Sonia Maria Enosse

118 Avenida Elias Lucas Kumato

Ivan Alejandro Pulido Tarquino

Malaria Consortium

Stije Leopold

Mahidol-Oxford Tropical Medicine Research Unit

Arjen Dondorp

Mahidol-Oxford Tropical Medicine Research Unit <https://orcid.org/0000-0001-5190-2395>

Halidou Tinto

Institute de Recherche en Sciences de la Santé

James Tibenderana

Malaria Consortium

Joel Tarning

Mahidol University <https://orcid.org/0000-0003-4566-4030>

Mallika Imwong

Mahidol-Oxford Tropical Medicine Research Unit, Faculty of Tropical Medicine, Mahidol University

<https://orcid.org/0000-0003-0857-1855>

François Nosten

Shoklo Malaria Research Unit <https://orcid.org/0000-0002-7951-0745>

James Watson

University of Oxford

Nicholas White

Mahidol University <https://orcid.org/0000-0002-1897-1978>

Article

Keywords:

Posted Date: May 4th, 2026

DOI: <https://doi.org/10.21203/rs.3.rs-9444378/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Additional Declarations: There is **NO** Competing Interest.

Parasitological efficacy of sulfadoxine-pyrimethamine and amodiaquine for Seasonal Malaria Chemoprevention (SMC) in Africa

Chayanin Sararat^{1,2,*}, Paul Sondo^{3,4}, Craig Bonnington², Jane Achan⁵, Chukwudi A Nnaji⁵, Anna Y. Burkhard⁵, Gauthier Tougri⁶, Adama Traore⁷, Aimé Parfait Yaro⁷, Anthony Nuwa⁸, Richard Kajubi⁸, Kanokon Suwannasin¹, Stephane Proux⁹, Uairat Koesukwiwat¹, Sonia Enosse¹⁰, Ivan Alejandro Pulido Tarquino¹⁰, Stije J Leopold^{1,2}, Arjen M Dondorp^{1,2}, Halidou Tinto⁴, James K. Tibenderana⁵, Joel Tarning^{1,2}, Mallika Imwong^{1,2}, Francois H Nosten^{2,9}, James A Watson^{2,3,*}, Nicholas J White^{1,2,†}

1. Mahidol-Oxford Tropical Medicine Research Unit, Faculty of Tropical Medicine, Mahidol University, Bangkok 10400, Thailand; 2. Centre for Tropical Medicine and Global Health, Nuffield Department of Medicine, University of Oxford, OX3 7LJ, UK; 3. Infectious Diseases Data Observatory, Oxford, UK; 4. Institut de Recherche en Sciences de la Santé- Clinical Research Unit of Nanoro (CRUN), Burkina Faso; 5. Malaria Consortium, London, UK; 6. Programme National de Lutte contre le Paludisme, Ouagadougou, Burkina Faso; 7. Malaria Consortium, Ouagadougou, Burkina Faso; 8. Malaria Consortium, Kampala, Uganda; 9. Shoklo Malaria Research Unit (SMRU), Mahidol Oxford Tropical Medicine Research Unit, Faculty of Tropical Medicine, Mahidol University, Mae Sot, Thailand; 10. Malaria Consortium, Maputo, Mozambique.

*For correspondence: james.watson@wwarn.org; chayanin@tropmedres.ac; †deceased;

Abstract

Seasonal malaria chemoprevention (SMC) with sulfadoxine-pyrimethamine plus amodiaquine (SPAQ) is given to millions of African children to prevent malaria during the peak transmission season, yet the contribution of the individual drugs to preventive efficacy remains unclear. We pooled individual-level data on drug concentrations and qPCR positivity from 934 children under 5 years from three prospective studies in Uganda, Mozambique, and Burkina Faso. We used natural variation in drug exposures to estimate the concentration-protection relationships against malaria infection over one month. The concentration of desethylamodiaquine, the active amodiaquine metabolite, was the primary determinant of malaria infection (odds ratio [OR] 0.55, 95% CI 0.44–0.68 per standard deviation increase in log concentration). Children with low day 28 desethylamodiaquine exposure (<24 ng/mL) had nearly twice the risk of malaria infection compared to those with high exposure (risk ratio 1.9, 95% CI 1.5–2.4). Only weak associations were observed with sulfadoxine and pyrimethamine concentrations. Concentration-protection relationships were consistent across countries despite marked differences in parasite resistance markers. These findings suggest that the preventive efficacy of SPAQ is primarily driven by amodiaquine, highlighting the importance of adherence to the 3-day amodiaquine regimen. The contribution of single-dose sulfadoxine-pyrimethamine appears modest and merits further evaluation in randomised trials.

Introduction

Malaria remains the leading cause of preventable death in young children in sub-Saharan Africa. In parts of sub-Saharan Africa such as the Sahel region, malaria transmission is highly seasonal with most cases concentrated in 4–5 months of the year during the rainy season. Seasonal malaria chemoprevention (SMC) consists of giving monthly treatment doses of slowly eliminated antimalarials to young children to prevent infection and illness during the peak risk period [1]. SMC has been recommended by the World Health Organisation (WHO) since 2012 [2] and in 2025, over 50 million children received SMC. In 2022, WHO broadened their recommendation around SMC allowing countries to choose how and where to implement it [3]. SMC can now be used in a wider age group, in areas where there is high grade resistance to sulfadoxine-pyrimethamine, and over more cycles each year. There is little evidence to support this broader recommendation.

The drug combination currently recommended by WHO for SMC is sulfadoxine-pyrimethamine plus amodiaquine (SPAQ) [1]. Sulfadoxine-pyrimethamine (SP) has been used as an antimalarial for over 50 years; a single treatment dose reliably prevented recurrence of infection with sensitive parasites within one month, although its efficacy is now compromised by multiple mutations in the *Pf dhfr* gene (conferring resistance to pyrimethamine) and the *Pf dhps* gene (conferring resistance to sulfadoxine) [4]. Nearly all *P. falciparum* parasites in West Africa are “quadruple mutants”, characterised by the *Pf dhps* 487G mutation but lacking the *Pf dhps* 540E mutation. However, in East and Southern Africa, nearly all *P. falciparum* parasites are “quintuple mutants” (540E) [5] or sextuple mutants, with substantially less susceptibility to sulfadoxine. The other component of SMC, amodiaquine, is rapidly converted to its active metabolite desethylamodiaquine which has a relatively long elimination half-life (approximately 10 days) [6]. Amodiaquine is given over three days to obtain sufficiently high drug concentrations for one month. For SMC to work effectively, children must maintain adequate drug concentrations in the blood throughout the protection window. Low drug exposure would result in reduced protection [7].

To better characterise the relationship between drug exposure and protection against malaria infection, in 2022 Malaria Consortium conducted three Chemoprevention Pharmacokinetic-Efficacy Studies (CPES) of SPAQ in Uganda [8], Mozambique [9], and Burkina Faso. These studies were embedded in larger implementation studies of SMC. The studies enrolled children under 5 years who were eligible for SMC; the children were administered their first cycle of SPAQ and then were followed up for one month with dried blood spots taken at baseline and then on days 7 and 28 after start of treatment to assess drug exposure and malaria infection by PCR. This approach allows for characterisation of the relationship between drug levels and protective efficacy against blood parasitaemia. In this pooled analysis, we look at the determinants of parasitological efficacy for SPAQ and assess whether current dosing regimens of sulfadoxine, pyrimethamine, and amodiaquine administered during standard SMC campaigns provide adequate protection over one month against *Plasmodium* infections in children under 5. If protection is inadequate, an important question is whether this is due to underdosing or drug resistance. This will contribute to understanding whether the dosage or choice of drugs needs to be modified to achieve the full potential for SMC to prevent malaria infection in the future.

Results


Participant Characteristics

A total of 1513 children were enrolled into SPAQ arms across the three studies: Uganda (n = 504), Mozambique (n = 489), and Burkina Faso (n = 520) (Figure 1). After excluding children with drug concentrations suggesting possible additional or off-schedule dosing, no biological samples for laboratory evaluation, or lacking post-hoc re-consent (Burkina Faso), 934 children were included in the analysis population (Figure 1 and Table 1). We restricted the analysis population to children with blood samples available at all three time points (days 0, 7 and 28) in order to have a fair comparison across the three drugs which have different pharmacokinetic profiles. The proportion of complete follow-up varied substantially by site. Only 104 children (21%) in Mozambique had complete sampling, compared with 81% in Uganda and 82% in Burkina Faso. Overall, participants were broadly comparable across sites with respect to age, sex, and nutritional status. However, the participants from Uganda tended to be younger and showed higher levels of malnutrition, particularly when assessed using mid-upper arm circumference-for-age (z-MUAC; Table 1 and Figure S1).

Table 1: Demographic and anthropometric characteristics of participants in the analysis population by country. For continuous variables we give the median values.

Site	Uganda	Mozambique	Burkina Faso
Enrolled participants (n)	504	489	520
Analysis population (n, %)	407 (81%)	104 (21%)	423 (81%)
Age (months, IQR)	33 (12 - 47)	33 (21 - 46)	32 (18 - 47)

Male sex (%)	212 (52%)	53 (51%)	223 (53%)
Weight (kg, IQR)	11.4 (9.0 - 14.0)	11.3 (8.6 - 15.0)	11.4 (9.1 - 13.5)
z-weight	-0.8 (-1.7 to 0.0)	-1.1 (-2.9 to 0.3)	-1.2 (-1.8 to -0.4)
MUAC (cm)	13 (12 - 14)	15 (14 - 16)	15 (14 - 16)
z-MUAC	-2.1 (-2.9 to -1.2)	-0.1 (-0.8 to 0.6)	-0.6 (-1.1 to 0.0)
Malaria infection at baseline (n, %)	239 (59%)	68 (65%)	131 (31%)



* DP: dihydroartemisinin-piperaquine

Figure 1: Flow of participants through enrolment, exclusion, follow-up, and inclusion in the analysis population across the three study sites.

Malaria infection prevalence

Half of the children in the analysis population (438/934) had qPCR detectable malaria infections at baseline (Figure 2). The proportion of children infected at baseline was lower in Burkina Faso (31%), likely due to the timing of the first SMC dose relative to the start of the transmission season. Infection prevalence increased with age, with a higher proportion of older children harbouring detectable infections. Most infections were caused by *Plasmodium falciparum*, which was detected in 91% (435/478) of children overall. Among PCR-positive samples, the proportion attributable to *P. falciparum* was similar across sites: 223/239 (93%) in Uganda, 63/68 (93%) in Mozambique, and 149/171 (87%) in Burkina Faso (Table S1 and Figure S2). Estimates of overall and *P. falciparum* prevalence differed only slightly when calculated using all available baseline samples (Table S2).

28 days after SMC administration, the overall proportion of children with malaria infection declined modestly to 30% (310/935), with consistent reductions observed across all age groups (Table 2). During the 28-day follow-up period, 44 children experienced intercurrent clinical malaria, including 24 (6%) in Uganda, 3 (3%) in Mozambique, and 17 (4%) in Burkina Faso. *P. falciparum* remained the predominant species at day 28, detected in 238/284 (84%) of PCR-positive children, corresponding to 78% in Uganda, 93% in Mozambique, and 86% in Burkina Faso.

A substantial burden of *Plasmodium ovale* infections were observed, accounting for 86/478 (18%) of baseline infections and 29/284 (10%) by day 28 (Table S1 and Figure S2). *Plasmodium malariae* was detected infrequently (11 patients, 2%), with 10 cases in Uganda, and one in Burkina Faso and was almost entirely cleared by day 28, with only a single case remaining in Uganda.




Figure 2: Malaria positivity at baseline and day 28 of follow-up period. Malaria infection was defined as either PCR positivity at day 28 or intercurrent clinical malaria. Children under one year of age received half the total dose of SPAQ relative to those aged 1-5 (shaded area). Prevalence point estimates are shown with 95% binomial confidence intervals.

Table 2: Summary of efficacy outcomes by country. *Malaria infection is defined as PCR positive by day 28 or intercurrent malaria illness (fever or history of fever within 24 hours with a positive RDT).

Site	Uganda	Mozambique	Burkina Faso
Malaria infection (n, %)*	125/407 (31%)	45/104 (43%)	140/423 (33%)
PCR positive on day 28 (n, %)	109/407 (27%)	44/104 (42%)	131/423 (31%)
Intercurrent clinical malaria* (n, %)	24/407 (6%)	3/104 (3%)	17/423 (4%)
Cycle threshold (Ct) values			
Range (min-max)	18.7-40	19.8-40	18.8-40
Mean	38.4	36.9	37.5
Median (IQR)	40 (37.7-40)	40 (34.3-40)	40 (35.7-40)

SMC drug concentration profiles

The concentrations of sulfadoxine, pyrimethamine, desethylamodiaquine, and amodiaquine were measured on days 7 and 28. Day 7 concentrations showed relatively limited variability and were evenly distributed across age groups. In contrast, day 28 concentrations declined with increasing age, most likely reflecting slower clearance in young children. Age-related effects were most apparent for sulfadoxine and desethylamodiaquine (Figure 3).

For sulfadoxine, the median day 28 concentration among children under 1 year was 5,045 ng/ml with undetectable concentrations in 8% of children, whereas the median sulfadoxine level was 849 ng/ml and undetectable in 50% of children above 1 year. Across sites, sulfadoxine levels were broadly comparable, whereas pyrimethamine concentrations were almost twice as high in Uganda (Figure 3). Amodiaquine concentrations fell below the threshold for reliable laboratory quantification (1.87 ng/mL) in 42% of children by day 7 and 94% by day 28, consistent with its rapid elimination half-life [10]. In contrast, for its active metabolite, desethylamodiaquine, levels were highest in Uganda at day 7. By day 28, desethylamodiaquine levels were similar in Uganda and Burkina Faso, while remaining lower in Mozambique. Given that amodiaquine was administered daily for three days, the observed variation in desethylamodiaquine levels across sites is likely to be explained by differences in adherence [11].




Figure 3: Drug concentration profiles of desethylamodiaquine, sulfadoxine, and pyrimethamine at day 7 and day 28 post-treatment, as a function of age in years with added jitter, colored by study site. Lines show spline trends. Horizontal dashed lines denote the lower limit of quantification (LLOQ) for each drug.

Drug exposure and protection against malaria

Across all three main SMC drug exposures (sulfadoxine, pyrimethamine, and desethylamodiaquine), higher post-treatment concentrations were generally associated with lower risk of malaria by day 28 (Figure 4). Although breakthrough infections occurred among some children with high drug exposure, parasite densities in these cases were lower than among children with lower drug exposure (Figure S5). We compared four exposure metrics for each drug: the day 7 concentration, day 28 concentration, AUC from day 7 to day 28, and terminal elimination half-life. The day 28 concentration was the best predictor of malaria infection for desethylamodiaquine and the day 7 concentration for sulfadoxine. For pyrimethamine, only the day 7 concentration was evaluable, as day 28 concentrations were below the LLOQ in 83% of children overall (64% in Uganda, 100% in Mozambique, and 99% in Burkina Faso).

Under a multivariable model including these three selected drug exposures, desethylamodiaquine exposure showed the strongest association with malaria infection by day 28, after adjustment for site, calendar time, age, and malaria infection at baseline. There was weaker evidence of an association with sulfadoxine exposure, and no clear protective effect was observed for the pyrimethamine exposure (Figure S6). For the desethylamodiaquine exposure, the relationship with parasitological malaria on day 28 was approximately linear on the logit scale (best fit in a generalised additive model). After standardising each drug exposure so the effects were comparable, one standard deviation increase in the log desethylamodiaquine exposure was associated with an odds ratio for malaria infection by day 28 of 0.55 (95% CI 0.44-0.68) (Figure S6). The corresponding odds ratios for one standard deviation increase in the log sulfadoxine and pyrimethamine concentrations were 0.80 (95% CI 0.63–1.00) and 0.88 (95% CI 0.67–1.15), respectively.

We further evaluated whether the protective effect of drug exposure varied by age. Interaction models were fitted separately for each drug. There was no clear evidence that the associations for desethylamodiaquine or pyrimethamine differed across the age range of enrolled children. In contrast, younger children appeared to derive greater benefit from higher sulfadoxine exposure, with the protective effect diminishing with increasing age. The interaction between age (in years) and day 28 sulfadoxine concentration corresponded to an approximately 20% attenuation in the protective odds ratio per additional year of age (interaction coefficient = 0.18 on the log-odds scale, $p = 0.0007$). Despite this age-dependent effect, the overall conclusions regarding the dominant role of desethylamodiaquine exposure remained unchanged.

The predicted probability of malaria infection by day 28 declined markedly with increasing day 28 desethylamodiaquine concentrations (Figure S7). The risk fell to approximately below 40% once concentrations exceeded the population geometric mean (z-score = 0; corresponding to approximately 24 ng/mL). Notably, fewer than 25% of children in Mozambique achieved concentrations above this threshold, compared with more than 50% of children in Uganda and Burkina Faso (Figure S7).

To estimate the causal effect of low (<24 ng/mL: population mean exposure) versus high day 28 desethylamodiaquine exposure, we used inverse probability weighting based on the known confounders: age, nutritional status, sex, and site, allowing for an unbiased comparison of low versus high exposure. Age was the primary determinant of low day 28 desethylamodiaquine exposure (Figure S8). Children with low day 28 desethylamodiaquine exposure had a risk ratio for malaria infection by day 28 of 1.92 (95% CI: 1.53-2.41).






Figure 4: Relationship between drug concentrations and malaria infection after SMC. Participants were ranked by the drug concentrations and divided into quintiles. For each site and quintile, the proportion of malaria-positive children (PCR+ or clinical malaria) is shown with 95% exact binomial confidence intervals.



Uganda Mozambique Burkina Faso

Higher day 28 desethylamodiaquine concentrations were associated with lower parasitaemia on day 28, as reflected by higher Ct values (Figure 5). As for the primary outcome, associations between the Ct values and the day 7 sulfadoxine and pyrimethamine concentrations were weaker. Strong evidence of drug–drug interaction was observed between desethylamodiaquine and sulfadoxine, and between desethylamodiaquine and pyrimethamine, suggesting diminishing incremental effects of higher desethylamodiaquine concentrations at higher levels of partner drug exposure (saturation of effect). In left-censored Tobit models accounting for Ct values at the limit of quantification, one standard deviation increase in desethylamodiaquine concentrations was associated with a 2 Ct value increase in day 28 Ct values (equivalent to an approximately 4-fold reduction in parasite density, $p < 0.001$). Each standard deviation increases in sulfadoxine concentration reduced desethylamodiaquine effect by approximately 1 Ct (approximately 2-fold reduction in the parasite density reduction, $p < 0.001$). Baseline Ct was independently associated with day 28 Ct, whereas age and nutritional status were not.




Figure 5: Relationships between *Plasmodium spp.* Ct values and drug concentrations. Scatter plots show observed D28 Ct values in relation to D28 desethylamodiaquine, D7 sulfadoxine and D7 pyrimethamine. Ct values were analysed as left-censored at a Ct value of 40. Solid lines represent estimated trends from Tobit regression models, adjusting for age, study site, baseline Ct value, mid-upper arm circumference, and enrolment date.

Molecular Resistance Markers

Known molecular markers associated with SP and AQ resistance were genotyped in samples collected at baseline (Figure 6, Table S3, and Figure S9). Mutations in *pfdhps* (A437G, K540E) and *pfdhfr* (N51I, C59R, S108N) associated with reduced sulfadoxine and pyrimethamine susceptibility, respectively [12, 13], were highly prevalent across sites, predominantly as pure mutant genotypes. The *pfdhps* 437G mutation was fixed or near-fixed in all settings: Uganda 97% pure + 3% mixed, Mozambique 78% + 20%, and Burkina Faso 88% + 8%, respectively. At *pfdhps* 540, Uganda and Mozambique had high frequencies of the pure mutant 540E allele, whereas Burkina Faso remained almost entirely pure wild-type, with rare mixed infections.

The *pfdhfr* mutations were also widespread. The 51I and 108N alleles were fixed as pure mutants in Uganda and Mozambique, and predominant in Burkina Faso (73% and 83%, respectively), where a small proportion of mixed infections was detected. The 59R mutation occurred at high prevalence, 91% in Burkina Faso and 98% in Uganda, with mixed genotypes more common at these sites (24% and 33%, respectively). In Mozambique, most 59R infections were pure mutants (89%), with only a minority mixed (7%).

For AQ-resistance-associated markers, we genotyped *pfcr*t codons 72–76 [14] and *pfmdr1* codons 86, 184, and 1246 [12]. All samples carried the wild-type *pfcr*t CVMNK haplotype at codons 72–76, with only a small number of mixed or mutant haplotypes observed in Burkina Faso. *pfmdr1* mutations were generally uncommon. At codon 86, the wild-type N allele predominated at all sites, with only rare mixed or pure 86Y mutants found in Burkina Faso. Similarly, at codon 1246, both pure and mixed mutant infections were only

sporadically observed in Uganda and Burkina Faso. The 184F allele occurred at moderate frequencies across sites, appearing mainly as pure mutant or mixed 184F/Y genotypes.




Figure 6: Proportion of key SPAQ-resistance-associated mutations at baseline across study sites. Stacked bars coloured by genotype status (pure wild-type, mixed mutant, and pure mutant). Markers include key positions in *pfdhps*, *pfdhfr*, *pfprt*, and *pfmdr1*.

Discussion

Effective malaria chemoprevention relies on sustaining sufficiently high drug exposures throughout the at-risk period to prevent reinfection and illness. Current SMC deployment consists of monthly administration of two drugs, sulfadoxine-pyrimethamine given as a single dose, and amodiaquine given once a day over three days. The initial randomised trials of SMC did not determine the relative contribution of each drug to the overall protective efficacy. Knowing the relative contribution of amodiaquine to the overall protective efficacy is important because it determines whether full adherence to the three-day course is important. This pharmacometric assessment of SMC uses natural variation in drug exposures due to variation in body weight, absorption and elimination to compare the contribution of sulfadoxine-pyrimethamine versus amodiaquine for parasitological efficacy. We show that the day 28 desethylamodiaquine concentration was the primary determinant of parasitological efficacy. Sulfadoxine (the slower eliminated component of sulfadoxine-pyrimethamine) contributed little to the protection against malaria infection. A notable and somewhat surprising result was the consistency of the exposure–response relationships across the three study sites. This is particularly striking given the different sulfadoxine-pyrimethamine resistance profiles across these settings. The consistency across sites strengthens the conclusion that amodiaquine exposure is the main determinant of parasitological efficacy in SMC.

Sulfadoxine-pyrimethamine was once used widely for the treatment of malaria in Africa in the 1990s and early 2000s but rapidly succumbed to resistance. In Africa, sulfadoxine-pyrimethamine is now only used for intermittent preventive treatment. Resistance to sulfadoxine-pyrimethamine is thought to be generally worse in East and Southern Africa where the *pfdhps* K540E mutation (quintuple mutants) is highly prevalent. However, parasites in West Africa are also partially resistant to sulfadoxine-pyrimethamine with very high prevalence of the *Pfdhps* A437G mutation (quadruple mutant). Data from treatment studies done over 20 years suggest that the duration of protection from recurrent infection with quadruple mutants is ~30 days, compared with ~16 days for the quintuple mutant [15]. This would suggest that sulfadoxine should have a much greater contribution in Burkina Faso compared to Uganda and Mozambique. The data from these studies do not support this prediction and show little contribution of the sulfadoxine component in West or in East and Southern Africa in terms of parasitological efficacy. Prevention of infection at day 28 was only observed in children who had high desethylamodiaquine levels (e.g. >24 ng/ml: population geometric mean exposure). This implies that adherence to the three-day course of amodiaquine is essential for the effectiveness of SMC as currently deployed.

Amodiaquine is an old antimalarial drug that was previously used extensively for prophylaxis in travellers because it retained efficacy against chloroquine resistant *P. falciparum*. However, its use in prophylaxis was discontinued in the 1980s because of rare toxicity, notably agranulocytosis and liver injury, in some cases fatal [16]. These effects appear to follow prolonged exposure (weekly or daily prophylaxis) and have not been observed in either treatment or seasonal malaria chemoprevention [17]. However, the dose given in SMC is likely to be the maximum tolerated dose [18]. It could be dangerous to increase the daily dose to obtain higher day 28 concentrations. Given that desethylamodiaquine concentrations are below optimal for many children even after fully observed treatment, this implies that amodiaquine alone cannot achieve very high protective efficacy. This supports additional interventions such as seasonal vaccination [19].

This pharmacometric analysis has some limitations. First, the quality of the data collection and follow-up was variable across three studies. There were substantial issues in data collection and follow-up in Mozambique where complete follow-up data was available in only 21% of participants. Because the exposures of interest are the drug concentrations at day 7 and 28, and the outcome (malaria infection) is measured in the same samples, this is unlikely to introduce bias. Second, it is not possible to entirely differentiate between the causal contribution of higher desethylamodiaquine concentrations versus higher sulfadoxine concentrations. Exposure to both drugs is highly correlated in the same individual (same mg/kg dose; same food effects on absorption and age effects on elimination). The correlation is not perfect, but it is possible that a stronger predictive relationship between desethylamodiaquine and malaria infection compared with sulfadoxine is due to higher accuracy of the desethylamodiaquine measurement (less noise). The effects can only be differentiated fully if the dosage of each drug were randomised independently (e.g. a factorial design). Third, the exposure-response relationships inferred in this analysis were derived primarily from qPCR detection of malaria parasites in blood. The primary therapeutic goal of SMC is not to prevent infection but to prevent illness. Parasitological and clinical efficacies are related by definition (illness cannot occur without infection), but the exact relationship has not been characterised. Age is a key effect modifier: any given infection is more likely to become symptomatic in the younger children. It could be the case that the clinical efficacy of amodiaquine is higher than its parasitological efficacy if sub-therapeutic concentrations do not prevent infection but sufficiently restrict parasite multiplication thus preventing symptomatic illness. However, higher parasitological efficacy would imply higher clinical efficacy, as well as providing greater spillover benefit in terms of reducing transmission.

This meta-analysis of three studies of SPAQ as SMC in West and East Africa suggest that the parasitological efficacy of SPAQ is sub-optimal. Desethylamodiaquine concentrations are too low in around half of all children. Sulfadoxine-pyrimethamine appears to contribute little to the overall parasitological efficacy. Randomised studies of alternatives to amodiaquine and of the added benefit of sulfadoxine-pyrimethamine should be conducted in the Sahel and in other countries considering SMC deployment.

Methods

Data

This pooled analysis used individual-level data from three observational studies evaluating seasonal malaria chemoprevention (SMC) with sulfadoxine-pyrimethamine plus amodiaquine conducted in Uganda (NCT05323721), Mozambique (NCT05186363), and Burkina Faso (NCT05478954). The study designs and methods for the Uganda and Mozambique components have been described in detail in the original papers [8, 9]. The Mozambique enrolment count differs slightly from the previously published trial report due to use of an updated participant list. Briefly, across all sites, children aged 3–59 months were enrolled at the time of SMC administration and followed through a standardised sampling schedule comprising blood sample collection at day 0 (pre-dose), day 7, and day 28. At enrolment, blood samples were used to quantify *Plasmodium* parasitaemia and, where sufficient DNA was available, to perform molecular genotyping of drug resistance markers (*Pfcr*t, *Pfmdr*1, *PfAAT*, *Pfdhfr*, and *Pfdhps*). Day 7 and day 28 samples were used to measure plasma concentrations of sulfadoxine, pyrimethamine, amodiaquine, and desethylamodiaquine, with day 28 samples additionally used to assess parasitaemia. Drug administration was fully observed in Uganda, but only the first dose of SP and AQ was directly observed in Burkina Faso and Mozambique.

All three studies were sponsored by the Malaria Consortium. The Mbale Regional Referral Hospital-REC (reference: MRRH-2022-168) and the Uganda National Council for Science and Technology (UNCST; reference: HS2212ES) approved the study in Uganda; the Comité Nacional de Bioética par a Saúde, Mozambique Ministry of Health (reference 130/CNBS/22) and the Research Ethics Committee of the Hospital Clinic of Barcelona (reference 16/2021) approved the study in Mozambique; the ethics committee for health research in Burkina Faso (CERS) approved the study in Burkina Faso (reference N° 2022-07-167). In all three studies, the parents or carers gave fully informed written consent and declared that they were willing and able to comply with the study protocol and blood sampling. In Burkina Faso, due to lack of clarity in the initial information form regarding blood sample collection and processing, a second consent was re-obtained after ethics committee approval to address incomplete information regarding the planned use of blood analysis results. The procedures were successfully implemented, and data from participants who could not be traced were excluded from the dataset prior to analysis.

Analysis population

The analysis population comprised children who received SMC and had complete pharmacokinetic and parasitological data across the scheduled follow-up visits (days 7 and 28). To ensure appropriate interpretation of drug exposure, any samples suspected of having received another dose were excluded from all analyses. Such cases were identified based on either (i) amodiaquine concentrations exceeding 10 ng/mL on day 28, or (ii) higher drug levels on day 28 compared to day 7.

Statistical analysis

Primary and secondary efficacy outcomes

The primary outcome was malaria infection by day 28, defined as detectable parasitaemia by PCR on day 28 or any episode of intercurrent clinical malaria between day 0 and day 28, defined as fever (axillary temperature $\geq 37.5^{\circ}\text{C}$ or history of fever within 24 hours) with a positive RDT.

Cycle threshold (Ct) values from qPCR at day 28 were analysed as a secondary outcome to explore the association between post-treatment drug exposure and parasite burden. Lower Ct values indicate higher parasite density. Ct values were treated as left-censoring with PCR-negative samples assigned a Ct of 40, corresponding to the assay detection limit.

Exposure

Drug exposure was evaluated using post-treatment concentrations of sulfadoxine, pyrimethamine, amodiaquine, and desethylamodiaquine measured on day 7 and day 28. For pyrimethamine, due to its short half-life only day 7 concentrations were available (all concentrations on day 28 were below the LLOQ). For amodiaquine, concentrations were frequently quantifiable at day 7 but decline rapidly with most values below the lower limit of quantification by day 28, aside from small number of outliers. For sulfadoxine and desethylamodiaquine, additional pharmacokinetic metrics were derived for each child, including the area under the concentration–time curve (AUC) from day 7 to day 28 and the terminal elimination half-life. The AUC was calculated using the log-transformed concentrations at the two sampling times and the trapezoidal rule, representing the average exposure over the interval. Terminal half-life was estimated from the log-linear decline between the two timepoints. All exposure metrics were standardized (z-score) to allow comparison across drugs.

Primary exposure-efficacy analysis: malaria infection by day 28

To investigate the relationship between SMC drug exposure and the probability of malaria infection by day 28, we fitted generalized additive models (GAMs) with binomial link functions. Analyses were first conducted separately for desethylamodiaquine and sulfadoxine followed by a combined model incorporating the selected exposure proxies for both drugs and day 7 pyrimethamine concentration.

For each drug, multiple pharmacokinetic metrics were evaluated as potential predictors of protection (Figure S6) in a univariable GAM adjusted for study site, baseline malaria infection, nutritional status (z-MUAC), and smooth functions of age and enrolment time stratified by site. Akaike's information criterion (AIC) were used to identify the most informative exposure proxy for each drug.

Based on this selection process, the standardized day 28 desethylamodiaquine concentration, the standardized day 7 sulfadoxine concentration and the standardized day 7 pyrimethamine concentration were included together in a multivariable GAM to assess their combined influence on malaria risk. All models were fitted using restricted maximum likelihood (REML) with automatic smooth-term selection to penalize overfitting and shrink non-informative terms. To reduce undue influence from extreme values, analyses were restricted to observations with $|z\text{-score}| < 3$ for day 28 desethylamodiaquine and $|z\text{-score}| < 4$ for day 7 sulfadoxine.

To evaluate whether the protective effect of each drug varied with age, additional models were fitted separately for desethylamodiaquine, sulfadoxine, and pyrimethamine, each including an interaction term between the standardized drug concentration and age. These interaction models retained the same adjustment set as the primary analyses.

Secondary analysis: parasite density (Ct values)

For secondary outcome, Generalised additive models (GAMs) with Gaussian errors and identity link were used to flexibly identify dominant drug effects and potential interactions, adjusting for age, site, baseline Ct, mid-upper arm circumference, and enrolment time. To appropriately account for left-censoring at Ct = 40, Gaussian Tobit regression models [20] were fitted. Primary exposure variables included standardised (z-scaled) concentrations of desethylamodiaquine at day 28, sulfadoxine and pyrimethamine at day 7.

Sensitivity analysis

To estimate the causal effect of day 28 desethylamodiaquine exposure on malaria outcome, we applied inverse probability weighting based on the propensity score. Low versus high DAQ exposure was defined using a threshold of 24 ng/ml as the population mean exposure. The propensity score, defined as the probability of low exposure conditional on baseline covariates, was estimated using logistic regression including age, nutritional status (MUAC z-score), sex, and study site. Stabilised inverse probability weights were constructed and applied to create a pseudo-population in which measured confounders were balanced across exposure groups. The primary causal effect was then estimated using a doubling robust weighted regression model including the same covariates in the outcome model.

Software and code availability

All analyses were conducted in R (version 4.4.3). GAMs were fitted using the `mgcv` package [21] and left-censored Tobit models were fitted using the `survival` package [22]. Propensity score-based inverse probability weighting was used in sensitivity analyses, implemented using the `ipw` and `survey` packages [23]. All analysis code used to generate the results and figures in this manuscript is available at [GitHub/Chayaninss/SMC-Pooled-analysis].

Data availability

Pseudonymised participant data used in this analysis are available for access via the WWARN website (<https://www.iddo.org/wwarn/accessing-data>). Requests for access will be reviewed by a data access committee to ensure that use of data protects the interests of the participants and researchers according to the terms of ethics approvals and the principles of equitable data sharing. Requests can be submitted by email to malariaDAC@iddo.org via the data access form.

Authorship and contributions

Conceptualisation and study design: NJW, FHN, JT, MI, JAW; Data collection: CB, PS, AN, CN; Laboratory analyses: JT, KS, SP, MI; Statistical analyses and visualisation: CS, JAW; Writing first draft: CS, JAW; Supervision: NJW, FHN, JAW, SL, AMD; Reviewing and editing: all authors.

Conflict of Interest Disclosures

The authors declare no conflict of interest.

Acknowledgements

We thank all participating patients, and everyone involved in the original studies. NJW is a Principal Research Fellow funded by the Wellcome Trust (093956/Z/10/C). JAW is a Sir Henry Dale Fellow funded by the Wellcome Trust (223253/Z/21/Z). Mozambique team would like to thank Dr. Pedro Aide from CISM for the extensive work and support given in the field during data collection and management, and sample management.

References

1. World Health Organization, *Seasonal malaria chemoprevention with sulfadoxine–pyrimethamine plus amodiaquine in children: a field guide*. 2023: World Health Organization.
2. World Health Organization, *WHO Policy Recommendation: Seasonal Malaria Chemoprevention (SMC) for Plasmodium falciparum malaria control in highly seasonal transmission areas of the Sahel sub-region in Africa*, in *WHO policy recommendation: seasonal malaria chemoprevention (SMC) for plasmodium falciparum malaria control in highly seasonal transmission areas of the Sahel sub-region in Africa*. 2012.
3. World Health Organization. *Updated WHO recommendations for malaria chemoprevention among children and pregnant women*. World Health Organization [Internet]. [cited 11 Feb 2023]. Available: <https://www.who.int/news/item/03-06-2022-Updated-WHO-recommendations-for-malaria-chemoprevention-among-children-and-pregnant-women> 2023.
4. Chotsiri, P., N.J. White, and J. Tarning, *Pharmacokinetic considerations in seasonal malaria chemoprevention*. Trends in Parasitology, 2022. **38**(8): p. 673-682.
5. Flegg, J.A., et al., *Spatiotemporal spread of Plasmodium falciparum mutations for resistance to sulfadoxine-pyrimethamine across Africa, 1990–2020*. PLOS Computational Biology, 2022. **18**(8): p. e1010317.
6. Tarning, J., et al., *Population Pharmacokinetic and Pharmacodynamic Modeling of Amodiaquine and Desethylamodiaquine in Women with Plasmodium vivax Malaria during and after Pregnancy*. Antimicrobial Agents and Chemotherapy, 2012. **56**(11): p. 5764-5773.
7. White, N.J., *Intermittent Presumptive Treatment for Malaria*. PLOS Medicine, 2005. **2**(1): p. e3.
8. Bonnington, C., et al., *Pharmacometric Evaluation of Sulfadoxine-Pyrimethamine-Amodiaquine and Dihydroartemisinin-Piperaquine Seasonal Malaria Chemoprevention in Northern Uganda*. Clinical Infectious Diseases, 2025.
9. Bonnington, C., et al., *Parasitological efficacy of seasonal malaria chemoprevention in Nampula, northern Mozambique*. Transactions of The Royal Society of Tropical Medicine and Hygiene, 2025.
10. Ding, J., et al., *Population pharmacokinetics of artemether–lumefantrine plus amodiaquine in patients with uncomplicated Plasmodium falciparum malaria*. British Journal of Clinical Pharmacology, 2026. **92**(2): p. 589-605.
11. Ding, J., et al., *Adherence and Population Pharmacokinetic Properties of Amodiaquine When Used for Seasonal Malaria Chemoprevention in African Children*. Clinical Pharmacology & Therapeutics, 2020. **107**(5): p. 1179-1188.
12. Grais, R.F., et al., *Molecular markers of resistance to amodiaquine plus sulfadoxine–pyrimethamine in an area with seasonal malaria chemoprevention in south central Niger*. Malaria Journal, 2018. **17**(1): p. 98.
13. Beshir, K.B., et al., *Prevalence of Plasmodium falciparum haplotypes associated with resistance to sulfadoxine–pyrimethamine and amodiaquine before and after upscaling of seasonal malaria chemoprevention in seven African countries: a genomic surveillance study*. The Lancet Infectious Diseases, 2023. **23**(3): p. 361-370.
14. Wicht, K.J., S. Mok, and D.A. Fidock, *Molecular Mechanisms of Drug Resistance in Plasmodium falciparum Malaria*. Annual Review of Microbiology, 2020. **74**(Volume 74, 2020): p. 431-454.
15. Mousa, A., et al., *Impact of dhps mutations on sulfadoxine-pyrimethamine protective efficacy and implications for malaria chemoprevention*. Nature Communications, 2025. **16**(1): p. 4268.
16. Hatton, C.S.R., et al., *FREQUENCY OF SEVERE NEUTROPENIA ASSOCIATED WITH AMODIAQUINE PROPHYLAXIS AGAINST MALARIA*. The Lancet, 1986. **327**(8478): p. 411-414.
17. Ndiaye, J.L., et al., *Safety of Seasonal Malaria Chemoprevention (SMC) with Sulfadoxine-Pyrimethamine plus Amodiaquine when Delivered to Children under 10 Years of Age by District Health*

- Services in Senegal: Results from a Stepped-Wedge Cluster Randomized Trial*. PLOS ONE, 2016. **11**(10): p. e0162563.
18. Cairns, M., et al., *Amodiaquine Dosage and Tolerability for Intermittent Preventive Treatment To Prevent Malaria in Children*. Antimicrobial Agents and Chemotherapy, 2010. **54**(3): p. 1265-1274.
 19. Chandramohan, D., et al., *Seasonal Malaria Vaccination with or without Seasonal Malaria Chemoprevention*. New England Journal of Medicine, 2021. **385**(11): p. 1005-1017.
 20. Tobin, J., *Estimation of Relationships for Limited Dependent Variables*. Econometrica, 1958. **26**(1): p. 24-36.
 21. Wood, S.N., *Fast Stable Restricted Maximum Likelihood and Marginal Likelihood Estimation of Semiparametric Generalized Linear Models*. Journal of the Royal Statistical Society Series B: Statistical Methodology, 2010. **73**(1): p. 3-36.
 22. Therneau, T., *A package for survival analysis in S*. R package version, 2015. **2**(7): p. 2014.
 23. Lumley, T., *Analysis of complex survey samples*. Journal of statistical software, 2004. **9**: p. 1-19.

Supplementary Materials

Table S1: Summary of malaria infection in the analysis population. Percentage of malaria infection, any species, count over samples available for PCR, while species-specific infection percentage counted over PCR-malaria positive subjects

Species	Site	Baseline	Day-28 follow-up
All	Overall	478/935 (51.1%)	284/935 (30.4%)
All	Uganda	239/407 (58.7%)	109/407 (26.8%)
All	Mozambique	68/104 (65.4%)	44/104 (42.3%)
All	Burkina Faso	171/424 (40.3%)	131/424 (30.9%)
<i>Pf</i>	Overall	435/478 (91%)	238/284 (83.8%)
<i>Pf</i>	Uganda	223/239 (93.3%)	85/109 (78%)
<i>Pf</i>	Mozambique	63/68 (92.6%)	41/44 (93.2%)
<i>Pf</i>	Burkina Faso	149/171 (87.1%)	112/131 (85.5%)
<i>Po</i>	Overall	86/478 (18%)	29/284 (10.2%)
<i>Po</i>	Uganda	47/239 (19.7%)	22/109 (20.2%)
<i>Po</i>	Mozambique	25/68 (36.8%)	1/44 (2.3%)
<i>Po</i>	Burkina Faso	14/171 (8.2%)	6/131 (4.6%)
<i>Pm</i>	Overall	11/478 (2.3%)	1/284 (0.4%)
<i>Pm</i>	Uganda	10/239 (4.2%)	1/109 (0.9%)
<i>Pm</i>	Mozambique	0/68 (0%)	0/44 (0%)
<i>Pm</i>	Burkina Faso	1/171 (0.6%)	0/131 (0%)
<i>Pv</i>	Overall	0/478 (0%)	0/284 (0%)
<i>Pv</i>	Uganda	0/239 (0%)	0/109 (0%)
<i>Pv</i>	Mozambique	0/68 (0%)	0/44 (0%)
<i>Pv</i>	Burkina Faso	0/478 (0%)	0/284 (0%)

Table S2: Drug concentrations across sites, with values summarized as median (IQR).

	Concentration (ng/mL)		
	Uganda	Mozambique	Burkina Faso
Sulfadoxine (Day 7)	28 (22.2 - 38.7) ×10 ³	24.5 (16.1 – 32.7) ×10 ³	31.3 (21.0 - 41.5)×10 ³

Sulfadoxine (Day 28)	1.6 (<LLOQ - 3.7) ×10 ³	<LLOQ (<LLOQ - 1.7)×10 ³	1.0 (<LLOQ - 3.2)×10 ³
Pyrimethamine (Day 7)	91 (70 - 120)	46 (35 - 70)	49 (34 - 69)
Pyrimethamine (Day 28)	<LLOQ (<LLOQ - 6)	<LLOQ (<LLOQ)	<LLOQ (<LLOQ)
Amodiaquine (Day 7)	2 (<LLOQ - 5)	<LLOQ (<LLOQ - 2)	3 (<LLOQ - 5)
Amodiaquine (Day 28)	<LLOQ	<LLOQ <LLOQ)	<LLOQ <LLOQ)
Desethylamodiaquine (Day 7)	155 (117 - 205)	99 (69 - 148)	98 (69 - 140)
Desethylamodiaquine (Day 28)	30 (18 - 46)	14 (9 - 24)	26 (17- 39)

LLOQ: lower limit of quantification (840 ng/mL for sulfadoxine, 4.42 ng/mL for pyrimethamine, 1.87 ng/mL for amodiaquine, and 2.95 ng/mL for desethylamodiaquine)

Table S3: Distribution of molecular resistance markers at baseline.

Gene	Position	Genotype	Frequencies of genotype status		
			Uganda	Mozambique	Burkina Faso
pfdhps	A437	A	0(0%)	1(2%)	3(4%)
		A/G	1(3%)	9(20%)	7(8%)
		G	33(97%)	36(78%)	74(88%)
	K540	K	0 (0%)	1(2%)	83(99%)
		E/K	0 (0%)	10(22%)	0 (0%)
		E	34(100%)	35(76%)	1(1%)
pfdhfr	N51	N	0 (0%)	0 (0%)	9(11%)
		I/N	0 (0%)	0 (0%)	14(17%)
		I	34(100%)	46(100%)	61(73%)
	C59	C	1(3%)	2(4%)	7(8%)
		C/R	8(24%)	3(7%)	28(33%)
		R	24(75%)	41(89%)	49(58%)
	S108	S	0 (0%)	0 (0%)	4(5%)
		N/S	0 (0%)	0 (0%)	10(12%)
		N	34(100%)	46(100%)	70(83%)
Pfcrt	C72	C	117(100%)	46(100%)	88(100%)
	V73	V	117 (100%)	46(100%)	88(100%)
	M74	M	117 (100%)	46(100%)	83(94%)
		I/M	0 (0%)	0 (0%)	4(5%)

	N75	I	0 (0%)	0 (0%)	1(1%)
		N	117 (100%)	46(100%)	83(94%)
		D/E	0 (0%)	0 (0%)	3(3%)
		E/N	0 (0%)	0 (0%)	1(1%)
		D/N	0 (0%)	0 (0%)	0 (0%)
		E	0 (0%)	0 (0%)	1(1%)
	K76	K	117 (100%)	46(100%)	83(94%)
		K/T	0 (0%)	0 (0%)	4(5%)
		T	0 (0%)	0 (0%)	1(1%)
Pfmdr1	N86	N	29(100%)	46(100%)	75(96%)
		N/Y	0 (0%)	0 (0%)	1(1%)
		Y	0 (0%)	0 (0%)	2(3%)
	Y184	Y	9(31%)	21(46%)	28(36%)
		F/Y	10(34%)	15(33%)	15(19%)
		F	10(34%)	10(22%)	34(44%)
	D1246	D	38(95%)	46(100%)	77(99%)
		D/Y	1(3%)	0 (0%)	1(1%)




Figure S1: Distributions of demographic characteristics of the analysis population. Dashed lines represent median values in each subpopulation.




Figure S2: Plasmodium species prevalence




Figure S3: *Plasmodium falciparum*-specific infection prevalence by age. Children aged 3–11 months received a half-dose of SPAQ. Prevalence point estimates are shown with 95% binomial confidence intervals.




Figure S4: Drug concentrations in vomited VS non-vomited children. Individual plots colored by site are presented along with population median and interquartile ranges as error bars. Dashed lines are the lower limit of detection of each drug. Treatment-related vomiting was recorded in 31 children: 13/407 (3%) in Uganda, 3/104 (3%) in Mozambique, and 15/423 (4%) in Burkina Faso. All children who vomited were confirmed to have received a repeat dose. After adjustment for site, vomiting status was not associated with drug concentrations for most drugs and time points. For desethylamodiaquine at day 28, mean concentrations were slightly higher among children who vomited (p-value 0.03), however, the ranges substantially overlapped between groups. The median (IQR) concentrations of day 28 desethylamodiaquine were 34 ng/mL (IQR: 18–55 ng/mL) in the vomiting group and 27 ng/mL (IQR: 17–42 ng/mL) in the non-vomiting group.




Figure S5: Distribution of standardized (z-score) exposure metrics for Desethylamodiaquine (DAQ), Sulfadoxine (SUL) and pyrimethamine (PYM). Metrics include standardized day 7 and day 28 concentrations, area under the concentration–time curve (AUC), and terminal half-life (HL), shown across the study population.




Figure S6: Relationship between parasite density and post-treatment SMC drug concentrations.




Figure S7: Odds ratios for malaria infection by day 28 in relation to post-treatment SMC drug exposure. Drug concentrations were log-transformed and standardised to allow comparison across drugs.




Figure S8: Predicted probability of malaria infection and post-treatment drug concentration distributions. Predicted probability of malaria infection by day 28 is shown against concentrations of D28 Desethylamodiaquine calculated for a child of median age, weight, and MUAC who was malaria-negative at baseline. Histograms show concentration distributions by site. The black dashed line marks the overall mean, and coloured dashed lines denote site-specific medians and interquartile ranges. Shaded vertical stripes indicate one-standard-deviation intervals from the mean




Figure S9. Propensity score for low day 28 desethylamodiaquine exposure (<24 ng/mL) by age.




Figure S10: Individual-level genotyping of key SPAQ-resistance-associated mutations at baseline across study sites, ranked by the number of mutant codons. Colours indicate genotype status (dark blue = pure wild-type, pink = mixed mutant, and red = pure mutant). Markers include positions in *pf dhps*, *pf dhfr*, *pf crt*, and *pf mdr1*.




Figure S11: Amodiaquine dose per kilograms administered by age. Children below one-year-old got 75 mg of amodiaquine (half dose) otherwise got 150 mg (full dose). Colours indicate whether day 28 desethylamodiaquine concentration was above or below the predefined threshold.




Figure S12: Propensity scores for low versus high day 28 desethylamodiaquine exposure.