

Damiano Archetti^{*a}, Cesare M. Baronio^a, Elisa Caselani^b, Marta Magno^b, Stefano Calza^c,
Giovanni de Girolamo^b, Alberto Redolfi^a

SUPPLEMENTARY MATERIAL

IDENTIFYING ACTIVITY PATTERNS FROM DIFFERENT ACTIGRAPHY DEVICES IN PEOPLE WITH SCHIZOPHRENIA SPECTRUM DISORDERS: A COMPARATIVE STUDY

^a Neuroinformatics Laboratory, IRCCS Istituto Centro San Giovanni di Dio Fatebenefratelli, Brescia, Italy

^b Unit of Epidemiological Psychiatry and Digital Mental Health, IRCCS Istituto Centro San Giovanni di Dio Fatebenefratelli, Brescia, Italy

^c Unit of Biostatistics and Bioinformatics, Department of Molecular and Translational Medicine, University of Brescia, Brescia, Italy

Supplementary Table 1: Full socio-demographic characteristics of patients and healthy controls from DiAPASon

Variables	HC	SSD
Marital status, n (%) *		
<i>Single</i>	25 (25.3%)	92 (82.1%)
<i>Married/cohabiting</i>	68 (68.7%)	8 (7.1%)
<i>Divorced/widowed</i>	6 (6.1%)	12 (10.7%)
Education [years] *		
<i>Mean (SD)</i>	16.6 (4.8)	12.1 (3.1)
<i>Median (Min- Max)</i>	17 (6-27)	13 (2-21)
Working status, n (%) *		
<i>Working</i>	90 (90.9%)	34 (30.4%)
<i>Studying</i>	8 (8.1%)	7 (6.3%)
<i>Not working</i>	1 (1.0%)	71 (63.4%)
Weight [kg] *		
<i>Mean (SD)</i>	73 (15)	80 (19)
<i>Median (Min-Max)</i>	72 (44-115)	78 (42-127)
BMI [kg/m²] *		
<i>Mean (SD)</i>	24.4 (3.9)	27.0 (5.8)
<i>Median (Min-Max)</i>	24.0 (17.7-35.5)	26.1 (16.6-44.8)

Acronyms: BMI= body mass index; HC= Healthy Control; n= total number; SD= standard deviation; SSD= Schizophrenia Spectrum Disorders patient; *= significant differences between HC and SSD.