

Orthodontic analysis improves the accuracy of lip morphology prediction in facial reconstruction

Nanami Aoki^{1*}, Hajime Utsuno¹, Shuuji Namiki¹, Muhammad Garry Syahrizal Hanafi¹, Hitomi Nojima², Hisako Saitoh¹, Saki Minegishi¹, Sayaka Yamada¹, Hiroyuki Harada³, Hideaki Kamochi⁴, Eriko Marukawa⁵, Koichi Sakurada¹

¹Department of Forensic Dentistry, Graduate School of Medical and Dental Sciences, Institute of Science Tokyo, 1-5-45 Yushima, Bunkyo-ku, Tokyo 113-8510, Japan.

²Department of Dental Radiology and Radiation Oncology, Graduate School of Medical and Dental Sciences, Tokyo Medical and Dental University, 1-5-45 Yushima, Bunkyo-ku, Tokyo, 113-8510, Japan.

³Department of Oral and Maxillofacial Surgical Oncology, Graduate School of Oral Health Sciences, Institute of Science Tokyo, 1-5-45 Yushima, Bunkyo-ku, Tokyo 113-8510, Japan.


⁴Department of Maxillofacial Surgery, Graduate School of Medical and Dental Sciences, Institute of Science Tokyo, 1-5-45 Yushima, Bunkyo-ku, Tokyo, 113-8510, Japan.

⁵Department of Regenerative and Reconstructive Dental Medicine, Graduate School of Medical and Dental Sciences, Institute of Science Tokyo, 1-5-45 Yushima, Bunkyo-ku, Tokyo, 113-8510, Japan.

*Corresponding author:

Nanami Aoki

Email: aoki.nanami@tmd.ac.jp | ORCID ID: 0009-0002-9896-5224



Online Resource 2. Intra- and inter-observer ICC results for each measurement item.