






G 20°C MLM




G 20°C BLINK




G 16h 8°C



MLM



BLINK



MLMM

MLM

BLINK

MLMM






Fig. S1 Manhattan and QQ plots associated with several studied traits for the three models used.


TS 20°C




MLM BLINK MLMM




TS 16h 8°C



MLM BLINK MLMM



NS 20°C



MLM


BLINK

MLMM


MLM

BLINK

MLMM



NS 16h 8°C



MLM


BLINK

MLMM

MLM


BLINK

MLMM




d 20°C


MLM




BLINK




d 10°C



MLM BLINK MLMM




b 10°C




MLM

BLINK

MLMM




t10 10°C



MLM



BLINK

MLMM





t20 10°C

MLM




BLINK




t30 10°C

MLM




BLINK




t40 10°C

MLM




BLINK




t50 10°C


MLM



BLINK




T70 20°C




MLM

BLINK

MLMM




T70 10°C




MLM

BLINK

MLMM




T80 20°C



MLM

BLINK

MLMM




MLM

BLINK

MLMM

rb



MLM


BLINK

MLMM


MLM

BLINK

MLMM



rt20



MLM


BLINK

MLMM

MLM


BLINK

MLMM




rt30

MLM




BLINK



rt50


MLM



BLINK




MLMM



MLM



BLINK

MLMM





rt80


MLM



BLINK




TSW




MLM

BLINK

MLMM



HS



MLM

BLINK

MLMM

